

LECTURES
ON
NATURAL AND DIFFICULT
PARTURITION.

LECTURES

J. E. Clark M.D.
ON

NATURAL AND DIFFICULT

PARTURITION.

25M
M952

BY

EDWARD WILLIAM MURPHY, A.M., M.D.,

PROFESSOR OF MIDWIFERY, UNIVERSITY COLLEGE, LONDON; OBSTETRIC PHYSICIAN,
UNIVERSITY COLLEGE HOSPITAL; AND FORMERLY ASSISTANT PHYSICIAN TO
THE DUBLIN LYING-IN HOSPITAL.

ΕΠΕΑ ΠΤΕΡΟΕΝΤΑ.

740

NEW-YORK:

SAMUEL S. AND WILLIAM WOOD,

No. 261 PEARL STREET.

1846.

EDWARD O. JENKINS, PRINTER,
No. 114 Nassau Street.

TO
RICHARD CARMICHAEL, ESQ.

M. R. I. A.

PRESIDENT OF THE ROYAL COLLEGE OF SURGEONS, AND OF THE MEDICAL
ASSOCIATION OF IRELAND.

CONSULTING SURGEON OF THE RICHMOND SURGICAL HOSPITAL, THE WHIT-
WORTH AND HARDWICK MEDICAL HOSPITALS, AND COR-
RESPONDING MEMBER OF THE ROYAL
ACADEMY OF MEDICINE,
PARIS.

MY DEAR MR. CARMICHAEL,

Under your guidance I received the first principles of my Medical education, and as you have sown the seed, it is only natural that you should reap the fruit.

The quality, however, depends so much more on the soil than the cultivation, that there is too much reason to apprehend that these *first fruits*, which I now present to you, may prove to be a barren production. Nevertheless, if you accept the Dedication as a testimony of the respect and esteem of a former pupil, the work will fulfil at least one object, in giving me the opportunity of thus expressing sentiments with which I am deeply impressed, and which I have ever felt towards you. I remain,

My dear Mr. Carmichael,

Very faithfully yours,

EDWARD W. MURPHY.

P R E F A C E.

THE following lectures are the substance of those that have been given on the subject of Natural and Difficult Parturition in University College. The writer has published them in this form, in order that his pupils, recalling to their minds the lessons they have received, may be enabled to examine the principles and precepts delivered to them with more care and attention than could be done from the evanescent impressions of an oral lecture.

The order of the course has been slightly altered by the introduction of the first two lectures on the Obstetric Anatomy of the Pelvis; these are generally given in the course of the session, with the Anatomy of the Generative Organs; those on Gestation follow, and then Parturition is entered upon. The importance of a correct knowledge of the anatomy of the pelvis, to render instructive the mechanism and the difficulties of Parturition, induces the writer to deviate from that arrangement, and to adopt the order at present observed. He trusts that the reader will therefore pardon what may appear to be unnecessary repetition in the subsequent lecture, where the anatomy of the pelvis is alluded to, and descriptions previously given are briefly recapitulated.

EDWARD W. MURPHY.

12 Henrietta street,
Cavendish Square.

TABLE OF CONTENTS.

LECTURE I.

BONES OF PELVIS AND PELVIS COLLECTIVELY.—MEASUREMENTS OF PELVIS AND FŒTAL HEAD.

Obstetric Anatomy of the Pelvis—Coxal Bones divided by Brim into the Iliac or Abdominal portion and Ischiatic or Pelvic portion—Plane of Ischium—Double Inclination—Pubic portion—the Sacrum—Key-stone of Pelvis—Hollow and Promontory of Sacrum—Coccyx. *Articulations of the Pelvis*—Symphysis Pubis—Sacro-iliac Articulation—Sacro-vertebral Articulation. *Pelvis collectively*—*Brim of the Pelvis*—Iliac portion of Pelvis. *Pelvic Cavity*, with the Soft Parts—Influence of its shape on the Head of the Child. *Outlet*—Perinæum—Object of measuring the Pelvis—Measurements of the Brim, of the Outlet—Measurements above the Brim, of the Brim, of the Cavity, of the Outlet—Table of Measurements of Eighteen Pelves—Measurements of the Child's Head 13—32

LECTURE II.

MODES OF MEASURING THE PELVIS.

Varieties in the Pelvis. *Deviations* from the Standard Pelvis—Pelvis too large—too small—imperfectly developed—like the Male Pelvis—irregularly formed. *Deformities* of the Pelvis—the Ovate Pelvis of Infancy—the Cordiform Pelvis of the Adult—Manner in which these Deformities are produced—Influence of Gravity, of Muscular Forces—Rickets and Mollities Ossium considered as Causes of Distortion. *Modes of Measuring* the Pelvis—Baudeloque's Callipers—Contouly's Pelvimeter—Boivin's Pelvimeter—Digital Measurement . . . 33—54

LECTURE III.

MECHANISM OF PARTURITION.

Definition of Labour—Divisions of Labours by various Authors—Division into Natural, Difficult, Preternatural, Complex—Definition of Natural Labour—Stages of Labour. Arrangement of the Muscular Fibres of the Uterus, on the External and Internal Surfaces. The Effect produced by the Contraction of the different Sets of Fibres—Manner in which the Dilatation of the Uterus is accomplished—Use of Liquor Amnii. Order of Uterine Contractions—Wigand's views—Objections to them—Dewees' Eplanation of Uterine Action—Objections. Different Conditions of the Os Uteri—Dilatable—Rigid—Inflamed. Influence on the Action of the Uterus. 55—72

LECTURE IV.

MECHANISM OF PARTURITION, CONCLUDED.

Second Stage of Natural Labour—Bearing Pains—Passage of the Head through the Pelvis—Positions of the Head according to various Authors—Naegele's Views—Mode of distinguishing Vertex Positions—Varieties in Face-Presentations—Characters of—not so dangerous as had been supposed—Dilatation of the Perinæum—Expulsion of the Head—Direction of the Shoulders. *Third Stage*—Separation of the Placenta—Causes of Retention—without Hæmorrhage 73—89

LECTURE V.

MANAGEMENT OF NATURAL LABOUR.

Premonitory Symptoms of Labour—Evidence of the Commencement of the first Stage—Character of the Grinding Pains. *Obstetric Duties of the Practitioner*—When summoned to attend—Caution respecting his Patient—Objects of the Vaginal Examination in the first Stage, before the Membranes are ruptured—Signs of the first Stage being completed—Mode of preparing the Bed, and supporting the Patient 90—103

LECTURE VI.

MANAGEMENT OF NATURAL LABOUR, CONTINUED.

Second Stage—Obstetric Duties of the Practitioner—Vaginal Examination—Its Objects—Support of the Perinæum—Its Intention and Mode of Accomplishment. *Third Stage*—Removal of the Placenta—Support of the Uterus—The Abdominal Bandage—Its Object and Mode of Application—Management of retained Placenta without Hæmorrhage 104—116

LECTURE VII.

DIFFICULT LABOURS.

Difficult Labours divided into Tedious and Laborious Labours—Causes of Tedious Labours—Over-Distension of the Uterus—Extreme Obliquity—Gradual Escape of Liquor Amnii—Hysterical Excitement—Mental Despondency—Rigidity of Os Uteri from Inflammation—Toughness of Os Uteri—Extreme Rigidity, etc. 117—132

LECTURE VIII.

LABORIOUS LABOURS.

Causes of delay in the Second Stage of Labour: Head of the Child misplaced—Too large, and too much Ossified—Hydrocephalic—Pelvis like the Male Pelvis—Deformed—The Ovate Pelvis—The Cordiform Pelvis—Head of the Child above the Brim of the Pelvis—Fixed in the Brim—Arrested in the Cavity—Characters of Arrest and Impaction of the Head 133—147

LECTURE IX.

LABORIOUS LABOURS.

Symptoms of Inflammation—Of Exhaustion—Treatment—Different Degrees—Disproportion in the Pelvic Cavity.—Case in which the Head moves very slowly through the Pelvis—Opposite Opinions as to its Management—Discussion of the Question—Attempt to determine it by Statistical Results . . . 143—153

LECTURE X.

LABORIOUS LABOURS.

Management of Cases when the Head of the Child is impacted—Opposite Opinions as to the Application of the Forceps—Examination of the Question—Forceps as an Instrument of Compression—Baudeloque's Experiments—Perforation—Advantages of Auscultation—Case when the Head is retained at the Outlet—Accidental Obstructions—Ovarian Tumours—Polypus—Fibrous Tumours—Osteo-sarcoma . . . 166—185

LECTURE XI.

OBSTETRIC OPERATIONS.—THE FORCEPS.

Instruments employed for preserving the Lives of the Mother and the Child—The Vectis—Of limited Application—Mr. Gaitskill's manner of using it—Proposed mode of Operation—Objection to use it as a Lever—The Forceps—Operation with the Short Forceps when the Head is resting on the Perinæum—Operation when the Head is in the Pelvic Cavity—Operation when the Head is fixed in the Brim of the Pelvis . . . 186—203

LECTURE XII.

OBSTETRIC OPERATIONS.

Operations to save the Mother only—Perforation—Importance of Auscultation—Conditions that authorise Perforation where the Child is alive—Mode of operating when the Head is fixed in the Brim or impacted in the Cavity—Comparative Merits of the Crotchet, and the Craniotomy Forceps—Operation when the Head is above the Brim, or can with difficulty be forced into it—Elizabeth Sherwood's case—Dr. Davis's Osteotomist—Baudeloque's Cephalotribe—Sigaultian Operation—Cæsarian Section—Induction of Premature Labour. . . 204—223

LECTURE XIII.

OBSTETRIC INSTRUMENTS.

THE VECTIS: History of—Invented by Roonhuysen, De Bruyn—Secret purchased by Visscher, and Van de Poll—Denman, Bland, Aitken, Lowder, Gaitskill.—THE FORCEPS: Invented by Dr. Paul Chamberlen, and used extensively by him

and his Sons, Hugh and Peter—Forceps contrived by Giffard, Chapman, Gregoire—Smellie's Improved Forceps, the Parent of those at present used—Varieties in the Construction of different Forceps—Objects intended by them—Difference in the Principle adopted—Dr. Davis's Forceps, Dr. Denman's, Dr. Beatty's, Dr. Ziegler's.—THE PERFORATOR: Smellie's Scissors, Denman's, Naegele's—Holmes's Perforators—Crotchet—Craniotomy Forceps—Conclusion.
224—241

DESCRIPTION OF INSTRUMENTS, WITH ENGRAVINGS 243—251

SUMMARY OF THE PRINCIPLES AND RULES, ARRANGED AS APHORISMS . . . 255—274

ERRATA.

On page 19, 12th line from bottom, for *two* wide, read *too* wide.

On page 23, 4th line from bottom, for *axis* of the pelvis, read *axes* of the pelvis.

On page 48, note, for † Vide p. 29, fig. 2, read † Vide p. 42, fig. 2.

LECTURES

ON

NATURAL AND DIFFICULT PARTURITION.

J. E. Clark M.D.

LECTURE I.

BONES OF PELVIS AND PELVIS COLLECTIVELY.—MEASUREMENTS OF PELVIS AND FETAL HEAD.

Obstetric Anatomy of the Pelvis—Coxal Bones divided by Brim into the Iliac or Abdominal portion and Ischiatic or Pelvic portion—Plane of Ischium—Double Inclination—Pubic portion—the Sacrum—Key-stone of Pelvis—Hollow and Promontory of Sacrum—Coccyx. *Articulations of the Pelvis*—Symphysis Pubis—Sacro-iliac Articulation—Sacro-vertebral Articulation. *Pelvis collectively*—*Brim of the Pelvis*—Iliac portion of Pelvis. *Pelvic Cavity*, with the Soft Parts—Influence of its shape on the Head of the Child. *Outlet*—Perinæum—Object of measuring the Pelvis—Measurements of the Brim, of the Outlet—Measurements above the Brim, of the Brim, of the Cavity, of the Outlet—Table of Measurements of Eighteen Pelves—Measurements of the Child's Head.

GENTLEMEN,


THE first subject to which I shall direct your attention, is the anatomy of the pelvis, so far as it is connected with the process of parturition. It would be unnecessary to enter minutely into its descriptive anatomy. We shall dwell only on those points that are connected with practical midwifery. In this sense, it will require your particular attention. The difficulties, sometimes almost insuperable, which occur in the process of parturition, arise most frequently from the disproportion that exists between the head of the child, and the space it has to pass through. The source of many of these difficulties is in the pelvis; and the irregularities in its shape are often the most frequent causes of difficult labour. Independently of this, the beautiful mechanism which Nature adopts in the passage of the

child through this bony cavity, requires an accurate knowledge of its anatomy, in order to understand the perfection of the contrivance.

The bones of the pelvis are three: *two coxal or hip bones, and the sacrum* with its appendage *the coccyx*. In the process of ossification, the coxal bones are not completed until a later period, consequently, the older anatomists have been in the habit of describing it, as consisting of three bones—the ilium—the ischium—the pubis; but this is evidently incorrect; we shall therefore consider them as parts of the same bone, consisting of an iliac, ischiatic and pubic portion.

The iliac portion is much the largest, and is divided into two parts, by a very well-defined line of demarcation, which is a continuation of the linea-ileo-pectinea, and terminates at the sacro-iliac articulation. Superior to this line is the iliac fossa, being the internal surface of a broad irregular portion of the bone sometimes called the *ala* of the pelvis: it belongs to the abdominal cavity. This portion is completely surrounded by very powerful muscles. The iliac attached to its internal surface, the gluteal to its external; and to its crest are inserted the muscles of the abdominal parietes. Being, therefore, a common point of attachment to muscles of such power, it is necessarily very irregular both in its shape and thickness, the crest rough and waving, the centre of the bone thin, smooth, and sometimes even diaphanous. In the female pelvis it is more expanded than in the male, gives a greater breadth to the hips, and, being sufficiently conspicuous, it is often taken as an index of the proportion of the pelvis itself; here, however, an error may be very readily committed, inasmuch as the cavity may be narrow, and yet the alæ of the pelvis very much apart. This greater breadth is obviously very advantageous, when the increasing uterus begins to occupy the abdominal cavity. In the well-formed pelvis, the distance from one antero-superior spinous process to the opposite is about ten inches.

The inferior portion forms a part of the true pelvic cavity, and principally consists of the ischium. Its internal surface is bounded by the obturator foramen on the one side, and the ischiatic foramen on the other: it is smooth, and corresponds to the acetabulum on the external surface. This surface is called (in obstetric language) *the plane of the ischium*, because the head



Vertical Section of Pelvis,
showing the anterior and posterior surface of the plane of the Ischium.—
The lines represent the inclined plane of the promontory—plane of the
brim—of the cavity—of the outlet. Vide p. 24.

of the child glides upon it in its descent, and passes forward under the arch of the pubis; but if carefully examined, it will be found to consist of two planes very slightly inclined in opposite directions, and divided by a line passing from the pectineal eminence to the spine of the ischium. In some pelves, this is more obvious than in others; but when the soft parts are attached, this will be found nearly corresponding to the reflections of the peritoneum which form the broad ligaments. Thus, the internal surface of the ischium before the soft parts are removed presents two broadly curved surfaces, one anterior, the other posterior; these greatly contribute to facilitate the rotation of the head of the child in passing through the pelvic cavity. *The anterior plane* terminates at the obturator foramen, an opening nearly filled with membrane, giving support to the internal and external obturator muscles, and offering less resistance to the advance of the head forwards, than if it consisted of bone. The obturator foramen is bounded by the ischio-pubic ramus, the pillar of the arch of the pubis. It presents a smooth surface, bevelled off towards the arch; and when the head passes from the obturator foramen upon it, this inclination greatly assists its exit under the arch of the pubis. *The posterior plane* terminates in the ischiatic foramen, and the portion of the head which

comes upon it, glides in a similar manner upon the coccygæus and pyramidal muscles, and shorter sacro-ischiatic ligament towards the hollow of the sacrum. Thus, the inclinations of these surfaces oblige the head to pass through the pelvic cavity in a spiral direction. The pubic portion of the bone is smooth on its internal surface, which also greatly favors its advance. The only remaining points connected with this bone worthy of your notice, are *the spine* and *tuber of the ischium*. To the former is attached the inner short sacro-ischiatic ligament, upon which the head glides into the hollow of the sacrum: it is, therefore, important that it should present no impediment to this motion; and being short, rather rounded and smooth in the normal pelvis, it does not do so; but if much acted upon by the muscles attached to it, new osseous matter is deposited; the spine increases in length, becomes rough, and sometimes is drawn inwards, and thus will present an obstacle that at once arrests the head. From a similar cause, the tubera-ischii often oppose the escape of the head from the outlet of the pelvis, when it is enlarged by increased deposition of bone. These causes of delay are only met with, when the pelvis is surrounded by strong and constantly exercised muscles; and therefore you may expect to find them more frequently among a healthy, vigorous, rustic population, than in towns or manufacturing districts. But in the latter class, a similar impediment may arise from a cause of a perfectly opposite character, wherein the pelvis becomes diseased, its osseous matter diminished, and the spines and tubera-ischii are pressed too close together.

The next bone to which I shall direct your attention, is *the sacrum*. This bone is placed between, and is very firmly united to the two other bones: it forms a kind of key-stone to the pelvic arch, upon which the spinal column rests. Its external surface is extremely rough, and gives insertion to the most powerful muscles of the back: the internal is smooth, and forms what is called, in obstetric language, *the hollow of the sacrum*. The curvature of the sacrum, which forms this hollow, is worthy of your attention: it varies very much in different pelvis; if it be too straight, the antero-posterior space of the pelvic cavity is diminished; if too abruptly curved, the coccygeal extremity resists the progress of the head.

That portion of the bone, however, which has received most

attention, is distinguished by the remarkable title of the *Promontory of the Sacrum*, a term used by the older authors, and is a sufficient evidence of the frequent instances in which the difficulty of labour has been attributed to this projecting point. The passage round the promontory, which was supposed to be the leading difficulty, became a kind of doubling the Cape of Good Hope; and if it were well formed, or that the head advanced beyond it, all was well. It is rather the intervertebral cartilage which unites the sacrum to the last lumbar vertebra, that forms the projection, than the superior surface of the sacrum; and therefore, the promontory lies a little above the sacrum, or rather the sacrum is its inferior boundary. The opposite extremity of the sacrum terminates in the coccyx, which, in the female pelvis, is generally moveable, and by its mobility contributes to increase the outlet of the pelvis, when necessary to parturition. If unfortunately ossification should take place between it and the sacrum, great difficulty is necessarily produced; but this accident is very rare in the healthy pelvis during the parturient period.

Such are the bones that constitute the pelvis. Let us now consider the manner in which they are united together, and you will perceive every provision to secure strength, and at the same time to avoid the effects of concussion. In this respect, the union of the coxal bones to each other affords a very perfect example. The *symphysis pubis* consists of a mass of highly elastic fibro-cartilage, arranged in a series of concentric laminæ, the outer layers firm and resisting, while those within are softer; and in the centre of them is placed a small arthrodial articulating surface, moistened with a portion of synovial fluid. The shocks to which the pelvis is liable in the more violent motions of the body, as in leaping, especially downwards, are all more or less concentrated upon the symphysis pubis, and hence a provision of this kind is necessary. In the female pelvis, by its greater breadth, the space between the pubic portions of the coxal bones is increased, and at the same time, a perfectly smooth surface is presented posteriorly to the head. If, unfortunately, the reverse should take place, if the symphysis were narrow, and still more, if it were made rough from ossific depositions, serious injury might be done to the soft parts lying between it and the head, and as the urethra lies in this position, you can readily

perceive the risk that might occur; hence it is necessary that the symphysis pubis should be broader and smoother in the female, than in the male pelvis.

The sacro-iliac articulation is remarkable in its contrivance to preserve immobility. If you examine carefully the articulating surfaces of the sacrum and ilium, you will find them so adapted to each other, or if I might use the expression, so dove-tailed on one another, as when pressed together not to admit of the least motion: such is the case even in the dried bones; but when in the recent state we find a firm cartilage intervening, the articulation surrounded by the strongest ligaments, and additional strength given by the tendinous expansion of the neighbouring muscles, we at once perceive, by the provision that is made to prevent it, the importance of preserving the union of the sacrum and coxal bones undisturbed. When we recollect the relative position of the sacrum, the key-stone of the arch we have described, the centre upon which the spinal column rests, the wedge which keeps the coxal bones apart, and of course the point of resistance to any force tending to compress these bones, as the lower extremities necessarily would, if not in this way prevented, we can understand why a compact and firm articulation is so essential.

The inferior extremity of the sacrum is united to the coccyx by a fibro-cartilaginous lamina similar to the intervertebral cartilages, and is supported by anterior and posterior ligaments. This articulation, as well as those connecting the small bones of the coccyx together, admits a certain extent of motion of one bone upon the other, so that the coccyx from being curved may be rendered nearly straight, a highly essential advantage in the female pelvis.

The superior surface of the sacrum is united to the last lumbar vertebræ, by an intervertebral fibro-cartilage, which differs from the others in being much deeper before than behind; the aspect of the two articulating surfaces are consequently oblique to each other, and the cartilage presents a broad surface anteriorly which, strictly speaking, forms what is called the promontory of the sacrum. It is the most prominent point of the spinal column anteriorly, the whole weight of which rests upon it; and this part would necessarily be pressed forward, if the pelvis were weakened by disease.

Having thus given you a detailed description of the several bones composing the pelvis and the manner in which they are united together, let us proceed to consider the pelvis collectively. You perceive that it is divided into two portions by the line already alluded to, a continuation of the linea-ileo-pectinea on either side, passing along the lower margin of the iliac fossa, and terminating at the sacrum: this is the brim of the pelvis, which in the older language of midwifery was called by the English term, “basin” and “brim of the basin;” a term which included not merely the line described, but the parts of the ilia above it. These divisions have been called by different names,—“the greater and lesser pelvis,” “the true and false pelvis.” Sometimes the whole portion above the line is still called “the brim”—and that below it “the cavity” of the pelvis.

It is more important, however, to recollect that the superior portion belongs to and forms part of the inferior boundary of the abdomen, and must be taken in connection with it: the axis of this part of the pelvis is therefore the same as that of the abdomen. The pelvis, and consequently the abdomen, are wider here in the female than in the male, in order to accommodate the uterus, when it becomes an abdominal viscus. It may, however, be too wide or too narrow: if the ilia be too open, they give no support to the uterus when it enters the abdomen; the natural obliquity of the uterus is therefore greatly increased—it falls too much to one side, and so may remain until labor begins—then the action of the uterus becomes irregular and inefficient, and labour is delayed from this cause alone;—if they are very upright, the uterus rises into the abdomen too much in the middle line of the body; and if the brim be too wide, the weight of the uterus presses down on the soft parts beneath it—it may descend even into the vagina and give rise to the disease called prolapsus uteri: if too narrow, the uterus has not room to pass between the pubis and the promontory of the sacrum, irritation takes place, premature action of the muscular fibres is induced, and miscarriage is the result: thus you see that even here exactness of proportion is important.

But this is still more remarkably the case, when we consider the “pelvic cavity,” or “true pelvis,” through which the head of the child has to pass: it will therefore require an attentive examination.

This cavity, which is destined to contain the genito-urinary organs, is bounded above by the brim of the pelvis, below by the tubera-ischii and coccyx. The smooth posterior surfaces of the bodies of the pubis with their connecting symphysis, the obturator spaces, and the ischio-pubic rami, form its anterior wall ; the sacrum and ischiatic notch, the posterior boundary ; and the planes of the ischium already described, constitute its sides. A very imperfect idea of the cavity is formed if it be confined to the dried pelvis : it is necessary to consider the empty spaces left in the bones as being filled up with the soft parts which belong to them in order to obtain an accurate notion of it. *In the recent pelvis*, therefore, we find, in the anterior wall, the obturator spaces occupied by fibrous membrane to which the obturator muscle is attached, having a small opening above for the transmission of the obturator nerves and vessels. This muscle is concealed by the levator ani, and both are enclosed in fibrous membrane, so that in this space a kind of muscular cushion is formed for the head, as it advances into the cavity of the pelvis. Immediately below the symphysis lies the subpubic ligament, and beneath it a continuation of the same fibrous membrane, giving passage to the urethra.

In the posterior wall, the ischiatic notch is converted into a foramen by the internal sacro-ischiatic ligament which unites the sacrum to the spine of the ischium ; and beneath it a second opening is formed by the greater sacro-ischiatic ligament which passes from the sacrum to the tubers of the ischium. The larger sacro-ischiatic foramen is occupied by pyriform muscle, the thick branches of the sacral plexus of nerves which converge to form the great sciatic nerve, and the gluteal and ischiatic vessels. The smaller foramen is filled by the tendon of the obturator muscle, and the pubic vessels ; but, lying in a plane posterior to the former, it is rather withdrawn from the cavity of the pelvis. The sides of the posterior wall of the pelvic cavity present also to the advancing head of the child, a muscular cushion similar to the anterior. The planes of the ischia form the sides of the cavity, and, if you recollect the aspect which they present, their double inclination anteriorly and posteriorly, you will perceive the effect which this must have on a body passing along their surface,—that it cannot preserve the same direction, but must necessarily be rotated slightly as it advances. If, for instance,

we assume that the head is so placed that it enters the pelvic cavity nearly transversely with the occiput, corresponding to the plane of the left ischium, and the sinciput to that of the right, if the occiput descend on the anterior plane, it is directed forwards toward the obturator space, while the sinciput, gliding along the posterior plane of the opposite side, is directed backward to the ischiatic space; thus slightly altering from the transverse to the antero-posterior direction. As the head descends still lower in the cavity of the pelvis, it meets anteriorly the ischio-pubic ramus, and posteriorly the inner short sacro-ischiatic ligament; the former is so bevelled off, that the occiput glides from it under the arch of the pelvis, the latter forms a smooth inclined plane upon which the sinciput passes into the hollow of the sacrum. When this is accomplished, the head will lie in the antero-posterior direction; so that if it enters in the direction which has been supposed, it cannot pass through the pelvic cavity without performing a rotation. Let us now consider whether, when the head enters the pelvic cavity, it is placed in the manner which has been assumed: this leads us to examine what is called the brim of the pelvis—the shape of the brim in the dried female pelvis is rather elliptic, the long axis being its transverse measurement; consequently, the brim affords more room to the head when it enters it in the transverse, than in any other direction. But this applies only to the denuded bone: if we examine the brim in the recent state, we find it not elliptic but triangular; the psoæ muscles form the sides; the promontory of the sacrum the apex; the base being the anterior portion of the pelvis, lying between the pectineal eminences on either side; hence the greatest space of the brim would not be in the transverse but in the oblique direction: the head therefore enters the pelvis in the oblique measurement of the brim, and when the occiput lies anteriorly (its usual position) it at once meets the anterior plane of the ischium, and is rotated in the manner stated.* The outlet of the pelvis also requires some attention; but there is so

* In the description here given of the progress of the head through the pelvis, it is assumed that the pubic angle is 90° , sufficient to allow the occiput to escape completely under it; but many pelves, in other respects well formed, have the pubic angle less than this: the occiput therefore descends along the ischio-pubic ramus still further, and the head is expelled in the oblique direction. The latter is the more frequent course.

much more of the soft parts than of bone entering into its formation, that here also the dried pelvis gives but an imperfect conception of it. As we should wish, however, to reserve a more particular description of the perinæum, it will be sufficient at present to assume that it is a firm resisting structure, partly closing up the lozenge-shaped space which the outlet forms. The sides of the quadrangle are constituted by the greater sacro-schiatric ligaments and the ischio-pubic rami, the angles are antero-posteriorly at the pubis and coccyx, and laterally at the tubera ischii; the perinæum closes up the posterior half; the vulva lies in the anterior: the sides are not in the same plane; those formed by the ligaments run downwards and forwards, while the ischia-pubic rami run downwards and backwards, both meeting at the tubera ischii, as if the lozenge were bent in the middle. If we suppose the head advancing still onward, and endeavouring to escape from the outlet, we find that a change in its motion takes place—the occiput passes down along the anterior sides until it has sufficient space to emerge under the pubic arch, it then becomes a fixed point, a centre of motion, round which the head rotates, from behind forwards, passing along the curve of the sacrum, straightening the coccyx and pressing upon, distending, and forcing open the soft parts which form the perinæum, with a force which can only be understood by recollecting the kind of power employed. If you please to class it with the mechanic powers, you may consider it a lever of the third order. This force of distension is so great, that the perinæum would seldom be preserved from laceration if nature did not adopt other provisions to prevent such an accident. Even in the best-formed pelvis, the outlet is narrow, and affords but little space for the head to pass; you can therefore readily perceive the difficulties which must arise, if there be any diminution of its transverse measurement.

The passage of the head through the pelvis shows mechanical contrivance in the construction of the latter. The more it is examined, the more perfect this mechanism will be found: its necessity will be obvious, if you reflect on the very close adaptation of the head to the pelvis. The human head is larger in proportion to the size of the offspring than that of any other animal, and in consequence of man's erect position, the cavity and outlet of the pelvis is more closed in, for the purpose of


supporting the weight of the viscera above it. The head being large and the pelvis narrow, every contrivance that nature can adopt, is essential to accomplish her purpose; and also the proportions between both must be so exact, that the slightest deviation becomes an obstacle. Hence obstetricians, even from an early period, knowing the importance of accuracy in these proportions, have endeavoured to reduce them to a standard of measurement. They have sought to ascertain the dimensions of the perfectly formed pelvis; and having fixed upon that as the normal standard, it was supposed that every deviation from it would explain one or other of the difficulties which may be met with. How far this is possible, we shall have again to consider; at present let us observe the manner in which it has been measured. Every impediment to the passage of the head seems to have been referred to two sources—either irregularity of the brim, or narrowing of the outlet. In nearly all the popular works on Midwifery, these are the only parts that are measured;* and, consequently, all difficulties are attributed to their irregularities. As the brim seems to be most commonly at fault, it has received a proportionate share of attention. It has been measured over and over again in the dried pelvis; very ingenious instruments have been contrived for the purpose of measuring it in the recent state, and during life; and every attempt has been made to determine by such means, beforehand, when it is possible for the head to pass, and when it cannot.

Measuring the pelvis in this way, and with this intention, may serve our purpose in detecting pelvic deformities where the disproportion is great; but it is very questionable whether it will enable us to discriminate those slighter degrees of disproportion which are so often the causes of delay and difficulty in parturition. It is not, therefore, with this object in view, that we would direct your attention to these measurements, but rather for the purpose of still further illustrating the mechanism already alluded to, as well as to point out to you that irregularities in the brim and outlet are not the only difficulties met with in the pelvis. Let us first consider the axis of the pelvis. If the brim of the pelvis be supposed to be a plane surface, and that a straight line passes perpendicularly through its centre and

* Dr. Churchill's valuable little work is, however, an exception.

is continued on both sides, this line would touch the hollow of the sacrum near the coccyx in one direction, and would pass out below the umbilicus in the other. This is called the axis of the brim. This line intersects and forms an angle with the perpendicular of the body more or less acute; the angle is generally 45° , but it may be much greater; sometimes it is less. The outlet has a different axis, and would be represented by a line passing downwards and forwards from a point below the promontory of the sacrum nearly in the direction of the vagina, but as this is curved, it would be more correct to describe it as touching the floor of the vagina, and passing out through the centre of the perinæum. These lines intersect each other; and in order that the head should pass from one axis into the other, it must describe a curve from the above downwards, which takes place when the occiput rests upon the arch of the pubis.


In place of confining your attention to the measurements of the brim and outlet of the pelvis, we would wish you to consider the pelvis in another point of view, as consisting of a series of planes taken from above downwards, having different aspects and different measurements. The first of these is above the



Horizontal section of pelvis made at the brim of the pelvis.—*The superior surface* showing the antero-posterior and lateral measurements of *inclined plane of the promontory*—The dotted lines represent the transverse and oblique measurements of the brim.


brim posteriorly, but meets it anteriorly at the symphysis pubis: it is therefore an inclined plane, forming an acute angle with the plane of the brim. The antero-posterior measurement of this plane is taken from the centre of the lumbo-sacral interver-

tebral cartilage to the top of the symphysis pubis; it is generally 4 inches, and is of importance, because it gives the projection of the promontory of the sacrum much more accurately than the same measurement of the brim: in fact, the true promontory is above the brim. The lateral measurements are taken from the same point to the pectineal eminences on either side; these are about $3\frac{1}{2}$ inches, but are very seldom found exactly equal. It is necessary to recollect these distances, because of the effect sometimes produced by their inequality. For instance, if the promontory be very much directed towards the right pectineal eminence, the head would be at once prevented entering the brim if it preserved its usual position, that is, with its anterior part opposite the right sacro-iliac synchondrosis: but if it took the other direction, so that the same part was applied in the same way to the left side, it would pass quite easily. Hence, in the same patient, one labour may be difficult and another easy, merely from the accidental position of the head. The




Inferior surface of same section, showing the antero-posterior transverse and oblique measurements of the brim of the pelvis.—The dotted lines represent the lateral measurements of the inclined plane.

second is the plane of the brim, of which the antero-posterior measurement, from the centre of the upper edge of the sacrum to the top of the symphysis pubis, is about four inches. The transverse, from the centre of one ilium to the other, is $5\frac{1}{8}$ inches, and the two oblique measurements, passing from each pectineal eminence to the sacro-iliac articulation of the opposite side, about 5 inches.



Remaining section of pelvis, showing the antero-posterior, transverse and oblique measurements *of cavity*.

We have, then, the plane of the cavity, one most generally omitted, and perhaps the most important of the three. The antero-posterior measurement of this plane passes immediately below the symphysis pubis parallel to that of the brim, directly backwards to a point above the middle of the sacrum; it is about $4\frac{1}{4}$ inches. The transverse passes from the centre of one plane of the ischium to the other, and is $4\frac{1}{2}$ inches; and the oblique lies between the centres of those muscular masses which fill up the obturator and ischiatic spaces. Their measurement is uncertain; but it is more than 5 inches.



Outlet of Pelvis.

The outlet cannot be considered as a plane. Its antero-posterior measurement is taken from below the symphysis pubis to the extremity of the coccyx, and when the coccyx is extended is about $4\frac{1}{2}$ inches. The transverse measurement between the tubers of the ischia lies above this, and is about the same; so that when the head is pressing through the outlet, it forms a pretty accurate circle round it.

If you study attentively these proportions, you will at once perceive how much they contribute to the rotation which has been


explained to you. Comparing the plane of the brim with the plane of the cavity, you find the transverse measurement of the cavity diminished, while the antero-posterior and oblique distances are increased. You may also observe that the oblique lines of the cavity approach nearer to the antero-posterior direction than those of the brim of the pelvis; so that when the head passes from the brim into the cavity, always seeking the widest space, it first rotates from the oblique of the brim into the oblique of the cavity, and as it descends, is obliged, from the convergence of the planes of the ischium, still more to assume the antero-posterior direction, until, the occiput escaping under the pubic arch, it becomes fixed in this position: then the second rotation of the head from behind forwards commences, the transverse measurements of the head corresponding to the transverse of the outlet, and the longitudinal passing successively out in the antero-posterior measurement of the outlet.* In stating to you these measurements, as being those of the standard pelvis, I am very far from wishing to convey to you that they are constant or immutable; on the contrary, you will find, when you examine these points for your own satisfaction, that the pelvis is no exception to natural objects in general, and that it agrees with them perfectly in this principle, that two of the same kind are never exactly alike. When you study it, and are accustomed to observe it, you will find as much difference in the pelves as you would in the faces of those to whom they belonged; and therefore, where no two pelves exactly agree, it would be impossible to fix a standard; consequently, the measurements given must only be considered as a kind of mean, to which there are numerous exceptions. In order to demonstrate this in a clearer light, you have before you a table of measurements of several pelves, all well formed, and through which the head of the child would readily pass, but no two of them agree. You have also before you the measurements of the pelvis given by different popular authors.† The transverse space of the outlet is sometimes measured in a different manner, in order that the whole space within the ischio-pubic rami may be included; that is, by making the symphysis pubis the centre of a circle, of which

* It is here assumed, that the pubic angle is sufficiently wide to allow the occiput to pass completely under the arch, and to place the head in the antero-posterior direction.

† Vide Tables No. 1 and 2, placed at the end of the Lecture.

these pillars (as they are sometimes called) are the radii ; the arc of the circle between these is measured in degrees, which of course gives the angle at the pubis. The pubic angle in the standard pelvis is 90° . Other measurements are given, which are not of equal importance. The depth of the cavity, anteriorly, posteriorly and laterally, is stated thus by Dr. Burns : Depth of symphysis, $1\frac{1}{2}$ inches ; depth of sacrum, 5 to 6 inches ; depth of ischium, $3\frac{3}{4}$ inches.


In the description given of the pelvis, it has been explained to you that, from its construction, the head must rotate first laterally, and then in the antero-posterior direction, before it is expelled ; but these are not the only motions of the head in its passage through that cavity—there are others, which still further illustrate mechanical contrivance, and which deserve your attention. Before alluding to these, however, we would wish you to understand, that when we speak of the *head* passing through the pelvis, and that in a certain direction, we do not mean you to suppose that this is *constantly* the case ; on the contrary, the head may enter the pelvis in a different position ; and that of the child itself is sometimes reversed, so that the breech or foot passes first. It is only for the purpose of illustration that we would assume the head as the presenting part, as it is called obstetrically, and its position as uniform. The variety of these positions, and the mode of ascertaining them, will again come under our consideration in future lectures.



Longitudinal Measurement of the Child's Head.

Like the pelvis, so the child's head has been measured in different directions. There are three measurements given of its longitudinal axis. One from the occiput, just above the neck to

the middle of the open membranous space between the frontal and parietal bones, called the "bregma," or "anterior fontanelle." This is generally about $3\frac{1}{2}$ inches. Another passes horizontally from the most projecting point of the occiput to the centre of the frontal bone, above and between the superciliary ridges. This is usually $4\frac{1}{2}$ inches. The third lies between the same point of the occiput and the centre of the chin. This is 5 inches.



Transverse Measurement.

There are also three transverse measurements. One between the parietal protuberances, which is $3\frac{1}{2}$ inches; another between the temporal fossæ, which is from $2\frac{1}{2}$ to 3 inches; a third between the zygomatic arches, which is from $3\frac{1}{2}$ to 4 inches. Sometimes three distances are given from the chin, as a central point: one gives the length of the face and forehead (mento-frontal) $3\frac{1}{2}$ inches; a second, from the chin to the bregma, is 4 inches; a third, from the chin to the occiput, is rather more than 5 inches. We have arranged them thus in a tabular form:

LONGITUDINAL	IN.	MEASUREMENTS.	IN.	TRANSVERSE MEASUREMENTS.	IN.
Occipito-Bregmatic . . .	$3\frac{1}{2}$	Mento-Bregmatic . . .	4	Bi-parietal	$3\frac{1}{2}$
Occipito-Frontal . . .	$4\frac{1}{2}$	Mento-Frontal . . .	$3\frac{1}{2}$	Bi-temporal	3
Occipito-Mental . . .	$5\frac{1}{8}$	Mento-Occipital . . .	$5\frac{1}{8}$	Bi-zygomatic	$3\frac{1}{2}$ -4

The shortest of these measurements in the longitudinal axis is the occipito-bregmatic; and when the head enters the pelvis, which it does obliquely, this is made to correspond to the oblique measurements of the brim, by the anterior part of the head being so pressed up that the chin rests upon the chest of the child: but

as the head descends into the cavity, and gains more space in the oblique and antero-posterior direction, the forehead advances more than the occiput, so that the occipito-frontal measurement corresponds nearly to the oblique of the cavity. A little lower down, at the short sacro-ischiatic ligament, the forehead becomes a resting-point, and the occiput again descends obliquely along the ischia-pubic ramus until it emerges, with part of the parietal bone, under the pubic arch. The head therefore, in its descent, seems, as it were, to oscillate upon its transverse axis.

The bi-parietal measurement of the head is generally stated to correspond to the conjugate or antero-posterior axis of the brim; and as the former is $3\frac{1}{2}$ inches, the latter 4, only half an inch is allowed for the soft parts, even in the best-formed pelvis; consequently, the least diminution of the conjugate axis causes a difficulty, a greater one becomes an obstruction. Hence, among accoucheurs, it has been an anxious problem to determine the smallest conjugate diameter (as it is improperly called) through which the head can pass without destroying the child.

Its solution has been attempted by comparing the bi-parietal measurement of the head with the conjugate of the pelvis, and it has been stated by Dr. Joseph Clarke (a high practical authority), that if the conjugate be less than $3\frac{1}{2}$ inches, a living child cannot pass the brim of the pelvis. But in all these discussions, it has been too confidently assumed that these two measurements of the head and the pelvis exactly coincide. They do not do so: on the contrary, as the head is entering the pelvic cavity, the parietal protuberance next the pubis descends lower than that next the sacrum, so that the bi-parietal axis lies obliquely downwards, and so it remains more or less until the occiput escapes under the pubis. The part of the head, therefore, which would be felt lowest in the pelvic cavity, is this parietal bone. By this means nature avoids the difficulty which would often arise, if both parietal protuberances descended in the same plane.

Thus you perceive that the head slightly rotates on its longitudinal axis also; and, in order to effect its passage through the pelvis, combines four distinct movements: two of them upon the vertebral column, one in the lateral and one in the antero-posterior direction; two on the head itself, one on its longitudinal and a second on its transverse axis. By the combination of these motions the passage of the head is ultimately effected.

MEASUREMENTS OF THE PELVIS STATED BY VARIOUS AUTHORS.

[illegible]

LECTURE II.

MODES OF MEASURING THE PELVIS.

Varieties in the Pelvis. Deviations from the Standard Pelvis—Pelvis too large—too small—imperfectly developed—like the Male Pelvis—irregularly formed. *Deformities of the Pelvis*—the Ovate Pelvis of Infancy—the Cordiform Pelvis of the Adult—Manner in which these Deformities are produced—Influence of Gravity, of Muscular Forces—Rickets and Mollities Ossium considered as Causes of Distortion. *Modes of measuring the Pelvis*—Baudeloque's Callipers—Contouly's Pelvimeter—Boivin's Pelvimeter—Digital Measurement.

HAVING, in the preceding lecture, explained to you the structure of the pelvis, its obstetric characters, its normal proportions and the manner in which the head of the child passes through it, let me now direct your attention to those numerous exceptions to the standard pelvis which are so often met with, and which become causes of delay or difficulty in parturition. Some of these are only *deviations* from the just proportions of the pelvis; others are *deformities* the consequence of disease. We shall therefore consider each class separately. 1st. The *deviations* or irregularities in the pelvis are various. It may be altogether too large, or too small; it sometimes retains its infantile shape although increased to its full size; it may resemble the male pelvis; and, again, some one, or perhaps all, of its proportions may be irregular; all these *deviations* may be met with in the healthy pelvis, and therefore should be considered separately from the next class. 2nd. The *deformities or distortions* of the pelvis which are produced by disease.

The first exception, then, to the normal pelvis is that which is *altogether too large*: here is an example of it, (No 9 and 17 in the table). Such a pelvis as this could never be a cause of delay in parturition; but it may cause danger notwithstanding. For instance, when the uterus is increased in its weight and size during

pregnancy, and it occupies the abdomen, the pelvis is its great support below ; but if the latter be too large, the uterus presses into the vagina, gradually inverting it ; there are instances in which it has passed quite through the vagina and appeared at the vulva before labour commenced. But although this may not happen ; yet when the vagina is at all distended and inverted in this manner, the foundation is laid for that troublesome disease, prolapsus uteri. Hence, the accoucheur, in such cases, uses every precaution after delivery to prevent the uterus pressing upon the vagina, until this organ returns to its original size. Another danger of a very large pelvis is, that the child may be too suddenly expelled. In most cases of parturition, the action of the uterus is continued a certain time before delivery takes place ; and without entering into any enquiry as to the cause of that action, we know by experience that when once set up, it does not suddenly cease, even when the immediate exciting cause is removed. If you watch the uterus after labour when the placenta is detached, you will find contractions and relaxations still going on, although slight in their degree ; these contractions sometimes increase so as to become “after-pains” (as they are called). Now, when the child, meeting no resistance from the pelvis, is suddenly expelled, the uterus may still continue to relax and contract, although the stimulus of the child be withdrawn : hence there is a danger of hæmorrhage taking place ; and if the uterus (as is very probable) be thrown into irregular contractions, a stricture may be formed at the cervix uteri. The blood may only flow into the uterine cavity, being retained by a clot formed above the stricture ; and thus a case of internal hæmorrhage may arise ; which, if you are not prepared for, will escape your observation. Again you must recollect, that during pregnancy the abdominal vessels are more pressed upon than usual, by the addition of the uterus in the abdominal cavity ; that it is the inelastic coats of the veins that yield to this pressure ; and that the column of blood in the cava and large iliac veins is of necessity diminished ; nevertheless the same quantity of blood ascends through other channels to the heart. What then must be the effect when that pressure is suddenly removed ? You may have a most dangerous syncope, the circulation being suspended in consequence of the right side of the heart being nearly emptied of its venous blood. Another accident, which is


sometimes the result of this kind of pelvis, is inversion of the uterus: the fundus of the uterus, meeting no opposition from the pelvis, sinks within itself as the child is being expelled; it is sometimes turned completely inside out, but more commonly the depressed fundus forms a kind of cup-shaped cavity at the top of the uterus; this irregularity immediately excites the fibres of the uterus into an action somewhat resembling intus-susception of the intestines; the inversion of the fundus is increased; and it is ultimately forced in this state through the vagina. You perceive, therefore, that a very large pelvis is not quite so advantageous as it may appear to be.

The pelvis may be too small in proportion to the size of the head of the child. You can perfectly understand that a female who is well-formed may have all the bones small, she is consequently low in stature, but not disproportioned; and, although the pelvis bears the same relation to the rest of the skeleton that the standard pelvis does, still all the measurements of the former are less than the latter. The pelvis is diminished, but diminished equally in every part. In such a case a child of the average size would pass with great difficulty. Instances of this kind are rare, although sometimes met with; but it would be quite incorrect to assume that the pelvis must be small in females of low stature, it is much more frequently the reverse, the bones of the extremities, though diminished in length, are large, and so is the pelvis; but in the instance referred to, the bones are not only short but proportionately small, and the pelvis is of corresponding dimensions; hence you will find in this, as in other instances, that an accurate observation of the bones of the extremities will assist you in forming an opinion of the pelvis. Again, there are instances where the same kind of small pelvis is met with in women of the average height. Dr. Rigby quotes three such cases, from Busch's Berlin Reports, in all of which labour terminated fatally.*


The development of the pelvis may be arrested: pregnancy may take place before it is perfectly completed, as in young girls. In order that the effects resulting from this circumstance be made intelligible to you, it is necessary to direct your attention to the

* Rigby's Midwifery, p. 185.

growing pelvis. If you examine the pelvis at the time of birth, you at once perceive that it is imperfect, its ossification is incomplete, the viscera that properly belong to it are in the abdomen, because there is no pelvic cavity. During childhood its development is only slowly advancing, and the form of the pelvis is constantly undergoing alteration to the period of adolescence, when it at length attains a tardy maturity. Thus the osseous covering which protects the generative organs is not completed, until the time when they are prepared to enter upon their proper function. It is important, therefore, to attend to the changes going forward. In the infant pelvis the unfinished ilia are short, rounded, very patulous, and without any foss. The brim of the pelvis looks almost directly forward, in consequence of the pubis lying so much below the sacrum, that a line passing horizontally backwards from the symphysis pubis would pass through the extremity of the coccyx. The antero-posterior measurement of the brim is the longest, the transverse is the shortest, just as in the lower animals. There is scarcely any pelvic cavity; the ischia are closed in; and consequently the tubers approximate, and the arch of the pubis is contracted: hence the transverse measurement of the outlet is very small, the antero-posterior equally long, and, being almost parallel with that of the brim, it resembles in this respect also the pelvis of quadrupeds. From this extreme, a gradual change goes forward, until the pelvis



Child's Pelvis.



No. 12.


No. 12. Large Female undeveloped Pelvis compared with young Pelvis.

assumes its permanent character. Now it may happen that this alteration of the pelvis is so arrested in the middle of its progress, that no farther change of shape takes place, but, nevertheless, that the pelvis continues to increase, just as you see in monstrous fœtuses an arrest of development at the sixth week of gestation magnified into a monstrosity in the full-grown child. Compare, for instance, the above pelvis, which is rather a large one, with the young pelvis, and you will observe the resemblance between them; the antero-posterior of the brim ($5\frac{1}{2}$ inches) longer than the transverse, the transverse of outlet proportionately diminished, and the cavity rather shallow: yet this pelvis is above the standard size, any difficulty or delay in the passage of the head could only arise from the outlet being incomplete. This also shows that the development of the pelvis may be arrested although its size continues to increase. But an arrest of development may take place, and the growth also be retarded: you may have in the adult woman the pelvis of the girl, with all its proportions below the standard, but not irregular or deformed. This had been pointed out by Mr. Shaw* some years ago; and the manner in which it occurs is very clearly explained. The rate of growth of the whole body is not equal. In the infant, the head, thorax, and upper extremities, are much more advanced in their formation than the pelvis and lower limbs. In the adult, it is the reverse, the latter exceed the former in their proportionate size. But if, from any cause, (as rickets,) the general growth of the bones be retarded, the pelvis and lower limbs will not increase so rapidly as they should do; they will still retain something of their immature character; consequently the pelvis may be too small although not deformed. It is possible such may have happened in the cases reported by Busch and quoted by Rigby. The development of the pelvis may be almost completed, and yet be too small for the passage of the head; and, as the difficulty may happen just at the time of puberty, it becomes an objection to early marriages. During the growth of the pelvis, the transverse and oblique measurements of the brim are constantly increasing and the outlet becoming wider; but they do not begin to exceed the antero-

* Med. Gazette, Vol. xvi. p. 45.

posterior until after puberty, as may be readily perceived in the altered shape and carriage of the female at that time. At puberty, therefore, these measurements may only equal the antero-posterior, the outlet and perhaps the cavity being still in diminished proportions. Thus the young girl, although perfectly well-formed but pregnant at too early an age, may be the victim of a difficult labour, simply from this cause.

Another deviation from the standard pelvis is *when it assumes the character of the male pelvis*; and, as this circumstance is much more frequently the cause of severe labour than is generally supposed, or at least than is stated, it deserves your particular attention. The contrast between these two pelvises must at once strike your attention. This resembles the male—



No. 17. The Male Pelvis.



No. 18. The Female Pelvis.

this, the female pelvis. Comparing them together, you find

the iliac bones of the male more upright, the crista ilii rougher and more waving, and the iliac fossa rather deeper.

The brim of the pelvis is more triangular than in the female, in consequence of the transverse measurement being less. Its axis is also directed more upwards. The cavity is much deeper; anteriorly, the symphysis pubis is narrower and longer, often ossified; laterally, the planes of the ischia are closer to each other; and, posteriorly, the sacrum is longer, narrower, and rather straighter, hence it is something like an obliquely truncated cone, inverted. The ischio-pubic rami form a more acute angle, which measures generally between 60° and 70° . The tubers of the ischia are closer, and if the coccyx be much curved, the outlet of the pelvis is very much closed in. The male pelvis is also much more ossified, and is consequently heavier than the female pelvis.

The cause of this difference must be at once obvious to you, if you recollect the different circumstances in which each pelvis is placed, and the law which seems to be observed in the growth of bone, viz. that it bears a strict relation to the purpose which it is intended to fulfil. If it be for an osseous covering, its size corresponds exactly to the development of the organs it protects: thus the alterations of the cranium keep pace with the varying development of the brain; a deformity from the sinking of the thorax is produced when the lungs are compressed. If it be as a centre of muscular action, it is strengthened and increased directly as the action of the muscles attached to it. Apply this rule to the pelvis, and you find one organ, and that an important one, absent in the male which exists in the female: hence the pelvic cavity is narrower in the former. The pelvis is also the centre of the most powerful muscular actions in the body; and where those muscles are stronger, and are called more frequently into action, they exert a proportionate influence upon the pelvis; therefore the male pelvis is more ossified and heavier than the female, and its shape is such as will give its muscles points of insertion the most favourable to their action: hence the distance between the opposite attachments of the same muscle is, as far as possible, lessened; consequently, in the male pelvis the ilia are more upright and nearer to the linea alba; the acetabula are closer, in order to diminish the distance between the pelvis as a centre of motion


and the thigh-bones ; and thus the arch of the pubis is narrower and the tubera ischii closer than in the female pelvis. If we have sufficiently explained this principle, let us consider for a moment how far the female pelvis might be influenced by it. It is true that the uterus is here contained ; and therefore we cannot explain by its absence any narrowness of the pelvic cavity which may exist : nevertheless, if we suppose the muscles connected with the pelvis to be large, strong, and constantly exerted, the effect would be nearly the same ; the acetabula would be drawn closer to the centre, the planes of the ischia would converge more, not, as in the diseased pelvis, protruding into the cavity, but by the bone in its growth adapting itself to the diminished distance. In a similar manner the ilia would be more upright, and the pelvis of the female would gradually assume many of the characters of the male. Such is frequently the case with women in the rural districts, who are strong, healthy, and constantly employed from early youth in carrying weights, and in other active muscular exertions. The difficulties offered by such a pelvis, are altogether different from those of the diseased pelvis, which, if you study what is written on the subject, would seem to be the only source of all the embarrassments that are met with. The obstacles, therefore, which the female pelvis may present to the passage of the head, when it approaches the character of the male pelvis, deserve your attentive consideration. The triangular shape of the brim is not generally an impediment, because, although the transverse measurement is diminished, the oblique is sufficiently wide, and the head will usually pass into the cavity. But here all the difficulties seem to centre : anteriorly, the symphysis pubis is narrower and more unyielding ; even a deposition of bone is sometimes found behind it, which may be extremely dangerous, if the intervening soft parts are pressed against it by the head : posteriorly, the promontory of the sacrum offers no opposition ; but the sacrum itself being straighter, there is less facility in the head performing the lateral rotation which has been already described to you ; and this difficulty is still more increased by the convergence of the ischio-pubic rami : the head is obliged to descend much lower in the pelvic cavity before it can escape under the arch of the pubis, and it is prevented from doing so in consequence of the space being so much lessened by its funnel-

shape. In addition to this, the tubers and spines of the ischia are more ossified, the one thicker and rougher, the other larger and more projecting; thus as the head advances, its passage becomes more and more impeded, until it is ultimately arrested, perhaps close to the outlet. In women of this description, it is possible, also, that the head of the child may be more than usually ossified, and the action of the uterus is always strong; so that a most disadvantageous combination of circumstances may take place in a healthy pelvis of this kind. On another occasion we shall have to refer to it; at present, we would only request you to notice its anatomical peculiarities, as it is important thoroughly to understand them; and here again we would observe, that the bones of the extremities will be a useful guide. The wrists and ankles are large, the phalanges thick and short: hence the old popular opinion amongst midwives, that "a thick, short hand is a bad sign when a woman is in labour," has a more just foundation than what, at first sight, might appear reasonable.

The last of the deviations in the pelvis to which I would direct your attention is, *where there is an irregularity and a want of correspondence between its different proportions*. The effect produced when the ilia are too patulous or too upright, has already been explained to you. In the brim of the pelvis there is a great variety in the direction of its axis. It may be too upright, and if the pelvic cavity be wide, it will cause prolapsus uteri in the manner that has been stated: but the axis is more generally in the opposite direction, and approaches too much the horizontal line. When this is the case, the weight of the gravid uterus is thrown very much upon the lower part of the abdomen, its parietes gradually yield to the pressure, and the uterus projects much more forwards than it should do. Sometimes, after several pregnancies, the abdomen has become so weak as to give it no support whatever; and it has been reverted over the pubis so as to rest on the thighs. We shall have again to point out how such a deviation may cause considerable delay in parturition. The greater inclination of the promontory of the sacrum to one side than the other has been already alluded to. The cavity of the pelvis, although sufficiently well-formed, often varies very much in shape and depth: it may be round, oval, triangular, deep or shallow, and yet cause little alteration in the

passage of the head. One variety of this kind, however, deserves notice, as it forms a pretty accurate contrast to the pelvis resembling the male. It is also funnel-shaped; but the funnel is reversed. There is rather less space in the brim than in the standard pelvis, it is a little more oval, having its short axis (antero-posterior) less than 4 inches; but the cavity is wider, the planes of the ischia more apart, and the outlet much more open than the normal pelvis. It is almost doubtful whether a pelvis of this character may not be *slightly* diseased, and consequently that it was beginning to assume something of that shape, the extreme of which forms the distortion of rickets.

Fig. 1.



Ovate Pelvis of Elizabeth Sherwood.


Fig. 2.



Cordiform Pelvis of Elizabeth Thomson.

From these varieties in the healthy pelvis, let me now direct your attention to those which are *diseased*, and which form the second class of exceptions to the standard pelvis. A very great difference may be observed in the shapes of the distorted pelvis. In one variety (Fig. 1) you find the brim not only oval, but inclining to an hour-glass shape by the close approximation of the promontory of the sacrum to the symphysis pubis: at the same time the cavity is shallow and open, and the outlet very wide; it is the extreme of the pelvis above alluded to. In another example (Fig. 2), you find the ilia very upright and almost doubled on themselves. The brim is called cordiform, that is, it resembles the ace of hearts, but when the distortion is great, it approaches much nearer the letter Y. The promontory of the sacrum and the pectineal eminences are quite close, and the bodies of the pubic bones are doubled back upon each other. The cavity is quite contracted in consequence of the planes of the ischia being pushed into it by the heads of the thigh-bones.

The sacrum looks as if it were broken ; it is bent up so abruptly at the coccygeal extremity. The tubers of the ischia are scarcely two inches apart ; and the ischio-pubic rami nearly parallel ; hence the pubic angle is, in some cases, only 10° .




The obliquely ovate pelvis of Naegele.

In a third instance, one side of the pelvis is of its usual shape ; and the opposite seems to run almost in a straight line from the sacro-iliac synchondrosis to the pubis, as if while soft, it had fallen on that side and were flattened. These three are the leading varieties we meet with in the *deformed* pelvis ; but from the two former there are numerous deviations, still, however, preserving their specific characters. The first is generally described as the deformity from *rickets*, and as being caused in infancy ; the second, as deformity from *mollities ossium*, and produced in the adult by that peculiar disease. The third variety has been known only within the last few years ; and we are indebted for our knowledge of it principally to the distinguished Naegele. Let us first compare the ovate and cordiform pelvises, and consider the reason why the shape of one differs so much from that of the other ; we shall then inquire into the nature of the disease, to which these changes are attributed ; and, lastly, consider the new variety of Naegele. In order to understand the cause producing the difference of shape observed in either of the pelvises alluded to, it is not necessary to enter upon the consideration of the disease that is said to be the cause of it. Rickets and malacosteon agree in the one result—softening of bone ; and it is sufficient to assume this condition of the

pelvis to explain, from other causes, the different appearance of each. Let me direct your attention therefore to some of those causes.


There are two forces constantly acting upon the pelvis, which, in its healthy state, it is always able to resist by the mechanical perfection of its structure : one force is caused by the weight of the body from above, and the resistance of the thigh-bones from below ; the tendency of the former is to press the promontory of the sacrum inwards and downwards ; that of the latter, to press the acetabula inwards, upwards, and backwards, towards the sacrum. Another force is the action of the muscles attached to the pelvis. A careful examination of the effect which these forces must have on either pelvis, in relation to the circumstances in which it is placed, will sufficiently explain the cause which modifies their form. They act very differently on the adult and on the infant pelvis : in the former, a line passing through the centre of gravity would fall rather within ; in the latter, in consequence of the straightness of the spine, it would fall outside and before the pelvis. In the one case, if the pelvis yielded, the spine and femurs would press in towards the centre—the cavity of the pelvis ; in the other, the weight of spine would fall in front of the pelvic cavity, while the acetabula would press up behind it. Hence the effect on the cavity would be, that when the line of gravity fell within it (as in the adult pelvis), it would be pressed inwards ; when beyond it (as in the infant pelvis), it would be drawn outwards by the divergence of the acetabula, the sacrum pressing down between them. Of necessity, therefore, the softened adult pelvis would take the shape called cordiform ; while the infant pelvis would be transversely lengthened. In this explanation, it is assumed that the spinal column preserves, in either case, its proper direction, and that it is not altered by disease. But it frequently happens (although by no means constantly) that the spine is softened and bent, as well as the pelvis ; this circumstance would modify the shape of the pelvis, by altering the line of gravity. You have here (Fig. 2) the pelvis of a child distorted like the adult pelvis—cordiform ; but there is a posterior curvature of the lumbar vertebræ, by which the weight of the body is thrown more upon the pelvic cavity.

Fig. 1.




Ovate Pelvis with lateral curvature of spine.

Fig. 2.



Cordiform Pelvis with curvature backwards in a child.

Anterior Curvature of Spine, with a singular Deformity of the Pelvis. (*Moreau.*)

Moreau, in his Atlas of Plates, gives a remarkable instance of the opposite kind: an adult pelvis, in which the spine just above the promontory of the sacrum is bent so much forwards, that the brim of the pelvis looks rather downwards. In this you see clearly that the weight of the body falls in front of the pelvic cavity, while the acetabula are pressed up behind it, consequently it so far resembles the oval pelvis, that its transverse measurement is increased. It differs from the oval pelvis in the conjugate measurement not being lessened: but the cause of this is obvious; the pubis lies completely behind the promontory of the sacrum, almost opposite the coccyx, which nearly rests upon

the tubers of the ischium. Therefore, although the distance between the pelvis and sacrum, measured directly backwards, is greatly diminished, that between the pubis and promontory is, if anything, increased, in consequence of the latter projecting so very much before the former. These exceptions, therefore, seem rather to prove the rule. Let us now consider the effect of muscular forces. In the motions of the body there are two sets of muscles connected with the pelvis to be considered, each having a distinct office to perform: one set, passing anteriorly and posteriorly between the pelvis and the thigh-bones, keep the pelvis fixed in its position; these, therefore, would act very powerfully in distorting the softened bone to which they are attached, but would manifestly produce a much greater effect when the body is upright and the pelvis is made a centre of motion, as in the adult pelvis, than when the body is bent forward and moves less upon the pelvis, as in the child. Such we find to be the case: the lower portion of the sacrum and the coccyx is bent nearly at a right angle by the great gluteal and pyramidal muscles, and closes up the outlet. Anteriorly, the effect is not so apparent in the adult pelvis, because it is counteracted by the acetabula and ischio-pubic rami being pressed in towards the centre; but you may still perceive the edges of these rami more everted, and the pubic arch itself immediately beneath the symphysis, wider than it ought to be. The other set of muscles are those that maintain the body in its erect position: posteriorly, the dorsal; and anteriorly, the abdominal muscles. The tendency of the former is to draw the sacrum towards the spine, and thus to increase the projection of the promontory: the effect of the latter is to draw the ilium more upright, and to render it more irregular. The action of these muscles will therefore explain to you the character of some of the distortions in the adult pelvis.

In the infant pelvis, their influence is modified by the altered position of the body. In this case, the weight from above presses down upon and tends to separate the thigh-bones more from each other; the muscles, therefore, passing between them and the pelvis, will draw that portion of the pelvis to which they are attached outwards: hence the ischio-pubic rami are more separated, and the tubers of the ischium more apart than natural; but the distance of the thigh-bones being increased,

the coccyx must still be drawn forwards by the muscles attached to it; consequently, the outlet is much more open than it ought to be, and the abruptly curved sacrum becomes the only impediment to the escape of the head. In this explanation of the distortions of the pelvis, we have confined our remarks to mechanical causes alone, and have made no allusion to the nature of the disease that gives rise to the softened state of the bone which prepares the pelvis for these alterations. We did so, in order that you might separate in your minds, the deformity of the pelvis and the disease to which the distortion is attributed, and that you might not suppose, as sometimes has been imagined, that the deformed pelvis is oval because it is rickety, or that its cordiform shape is the necessary consequence of mollities ossium. Let me now direct your attention *very briefly* to these diseases; and perhaps we may be able to remove another error almost as popular, that rickets is alone met with in childhood, mollities ossium in the adult. The term *rickets* has its origin in *ραχις* the spine, because spinal distortions form so prominent a feature in the disease; but the term *mollities ossium* might be equally well, if not better, applied to it. It is met with generally at that period of infancy (dentition,) when there is a formation of new bone going forward, and arises when the demand for ossific matter is not sufficiently supplied. Whatever be the cause that deranges the health of the child, imperfect nutriment, impure air, or hereditary disease, the effect is the same, the blood does not supply the want that is felt; the teeth are always very late in their appearance; the bones have not firmness to resist the forces that act upon them, and hence the deformity. Under proper management, the child generally recovers from the disease, but not from the effects of it; and the pelvis, distorted in infancy, is never restored to a perfect state. Now this softened state of the bones can scarcely be considered as an essential disease; rickets is only one of the effects of a general derangement of the health in which other structures than bone are equally affected, and then the question arises, whether the same causes, acting at a later period in adult life, may so derange the health as to produce rickets? Whether, in fact, healthy girls brought into large factories, or other confined situations, may have their health ultimately so deranged as to have rickets, and consequent deformities of the pelvis? If

such were the case, the ricketty pelvis in these instances would be cordiform and not oval. We have every evidence, if we call to mind the number of instances in which spinal deformities occur about the same period, that these distortions of the pelvis are likely to take place, in the same manner as they do in infancy, from a deficient supply of osseous matter; that rickets is the consequence of a general derangement of the health; and therefore, you must not confound it with *mollities ossium*, which is a distinct and very rare disease. *Mollities ossium* is accompanied by much more urgent symptoms, startings, restlessness, pains like rheumatism, more or less through the body, and general fever; there is also a white sediment found in the urine. The disease is seldom arrested, but pursues its course more or less rapidly, until the bones are so altered that they consist only of thin shells covering a grumous liver-like substance, they bend in all directions, and may be cut through with a knife. One of the most striking cases of this kind, you will find in Professor Cooper's Surgical Dictionary, the case of Madame Suppiot. Another equally remarkable, is recorded by Mr. Solly.* If you read them with attention, and consider their true characters and the course of the disease, you may perhaps arrive at the conclusion, that a pelvis softened by *mollities ossium* is not likely to offer a very great resistance to the head of the child. Such instances, however, have occurred when the progress of the disease has been arrested; and those *extreme cases of distortion* which have rendered the Cæsarian section necessary are generally caused by this disease. You have here an instance, in the pelvis from which this cast is taken, of the extent of the deformity. The case which it illustrates, was that of Elizabeth Thomson,† upon whom the operation was performed, by Mr. Wood, of Manchester.‡ But such cases taken in comparison with the general number of deformed pelvises, are only remarkable exceptions "*rari nantes in gurgite vasto*," which by no means authorises us to consider the cordiform shape of the adult pelvis, as an indication that *mollities ossium* was its cause. If, therefore, we may assume that derangement of the general health, arising from other causes than *mollities ossium*, close

* Med. Gazette, 1842-43, p. 510.

† Vide p. 29, Fig. 2.

‡ Memoirs of Med. Society (Manchester)? Vol. 5.

confinement, poor diet, impure air, or hereditary disease, may produce deformities in the adult pelvis, just as in infancy, you have the source of a class of cases (unfortunately too large) where these deformities are met with. You will find them in the large manufacturing towns and districts, perhaps also among the poor needle-women of London, sometimes even in the higher ranks of the aristocracy, and in all these, the deformity of the pelvis will present those characters more or less, of which this cast is the extreme. Such patients are a perfect contrast in their appearance, and in their pelves, to those strong active women having pelves like the male, and as both are equally liable to difficulty in parturition, it is important to remember the distinction between them. This we shall consider in a future part of the course; at present, we would merely direct your attention to some of the external characters, which accompany this condition of the pelvis. The peculiar and well-known aspect of scrofula may often be observed; but this is not always the case. It is rather in the osseous system you will find the safest guide; the extremities of the bones are large, the teeth imperfect and uneven, the hands fine, but the points and joints of the fingers thick, the nails are short and easily broken, the ankles are large and generally bent in towards each other. Along with this, you may sometimes have a slight curvature of the spine, but you must be cautious in assuming that the pelvis is deformed because the spine deviates from its proper direction. Here, as in the former instances, the extremities will be the safest indices of the character of the pelvis.

It is not necessary to dwell upon the varieties that are met with in these two forms of distortion. It seldom happens, for instance, that the promontory of the sacrum projects forwards exactly in the middle in the antero-posterior measurement; it generally inclines to the right or left side. This has been explained by Baudeloque, on the supposition that "it is because the rickets has not equally affected all the bones of the pelvis, nor equally hurt all their junctions: and because the attitude which the child takes in walking or sitting, may change a little the direction of the compressing power which I have just mentioned.* This explanation seems to me quite unnecessary. If


* Baudeloque, by Heath, vol. i. p. 60.

you take a softened pelvis, and try with a moderate force to press the promontory forwards in the middle line towards the pubis, you will find it much more difficult to do than to give it an inclination to either side. When the softened pelvis is compressed between the opposite forces, its tendency is to become slightly twisted on itself, and therefore to alter the direction of the sacral promontory.

The next variety of deformed pelvis to which we would direct your attention, is the pelvis of Naegele, which he calls "the obliquely ovate." One side is quite normal, but the opposite is not at all expanded; on the imperfect side, the sacro-iliac synchondrosis is ossified. It might therefore be explained by supposing that absorption of bone, the consequence of disease, had taken place at that symphysis, and thus produced the deformity; but we have no evidence, in the history of these cases, of any such previous disease; the assumption is consequently gratuitous, and the pelvis still remains a problem to be solved. We are indebted to Dr. Knox, of Edinburgh, for a very rational attempt to do so, which further observations may, perhaps, render successful. Dr. K. attributes it to arrest of development at the period when the ossification of the sacrum is incomplete; that one side of the pelvis advances to completion, while the opposite remains stationary; and that in the adult pelvis you have one side perfect, but on the opposite, the undeveloped pelvis magnified, just as if the lateral halves of the full-grown and infant pelvis were joined in the middle. This view seems to be supported by the fact, that the sacrum on the affected side is incomplete; but it is right to state that the remaining portion of the bone on that side does not show the same deficiency of development.

The concluding subject to which I would direct your attention, is to the different modes of measuring the pelvis, and to the variety of pelvimetres: these need but a brief description.

Contouly's pelvimetre is a straight, graduated rod, upon which slides a smaller one; each has an upright, just like a shoe-maker's rule, and the pelvis is measured by placing one upright against the sacrum, and sliding down the other, until it rests against the symphysis pubis; the distance between each is given on the scale. How this can be managed in the living pelvis, with all the soft parts attached to it, and the uterus, or perhaps,



Contouly's Pelvimetre.

the head, in the way, I must leave you to find out. Baudeloque employed a different instrument, a pair of callipers. He expected that by measuring between the antero-superior spines of the ilium in one direction, and from above the spinous process of the sacrum to the symphysis pubis in another, he could calculate the pelvis. We have already explained to you the uncertainty of any conclusion as to the pelvis, derived from the distance between the ilia. Little dependence, therefore, can be placed upon the transverse measurement. You are indebted to your late respected Professor Davis for a proof, that just as little reliance can be placed upon the antero-posterior. Baudeloque assumed that the thickness of the base of the sacrum was always three inches, therefore if the callipers gave seven as the antero-posterior distance, the true measurement of the brim would be four inches, and so on. Dr. Davis put this to the test. He measured the distance between the promontory of the sacrum and the middle point of the spinous ligament passing from the last lumbar vertebra to the sacrum, in seventeen pelves, taking them indifferently, well formed and distorted. The following are his results, in which you must recollect that the soft parts being absent, an additional source of error was removed (vide table*); you see

* In Fig. 1, pelvis, the distance was 3 inches.

" 2,	"	"	2 $\frac{7}{8}$	"
" 3,	"	"	"	"
" 4,	}	"	2 $\frac{5}{8}$	"
" 5,		"	"	"

that there is a full inch difference in the thickness of the base of the sacrum, which would be no trifling matter, if added or taken from the conjugate measurement of the brim. These two instruments are the best known and most frequently employed, especially on the continent. But there are others also employed.

The late Mdme. Boivin, aware of the objection to Contouly's pelvimetre, thought that it might be improved, and she contrived an instrument by which the antero-posterior distance was taken differently. One curved rod was passed into the rectum to the promontory. Another, sliding upon it, but capable of being raised to a given angle, was passed into the vagina; when there, it was pressed up to the pubis, and the angle formed between both rods measured on a scale, gave the antero-posterior distance. If you recollect the direction which the rectum takes, you will perceive that the rectal-rod either passed to the left side of the promontory, or dragged the rectum out of its position in front of it. Much dependence could not therefore be placed upon this.

Besides these modes of measuring the pelvis, none of which can be depended upon, there are what have been called "digital measurements;" or, in other words, the experienced accoucheur, from constant habit, when he passes the fingers or hand into the vagina, will form a very accurate estimate of the space in the pelvis. This is done in different ways: if one or two fingers are pressed towards the promontory of the sacrum, and if they at all approach it, it is certain the promontory projects too much, otherwise this never could happen, as you will find, if you try the experiment on the dried pelvis, sometimes as much of the hand as the vagina will admit, is introduced; if the sacrum below the

In Fig. 6, pelvis, the distance was $2\frac{1}{6}$ inches.

" 7, }	"	"	"
" 8, }	"	"	$2\frac{1}{4}$ "
" 9, }	"	"	"
" 10, }	"	"	"
" 11, }	"	"	$2\frac{3}{8}$ "
" 12, }	"	"	"
" 13, }	"	"	"
" 14, }	"	"	$2\frac{1}{2}$ "
" 15, }	"	"	"
" 16, "	"	"	$2\frac{1}{4}$ "
" 17, "	"	"	2 "

promontory and the pubis are only touched, when the fingers are separated, it indicates sufficient space; if it be impossible to separate them, the contrary: and from the degree to which the fingers are compressed, the amount of disproportion is estimated. In some instances it was impossible to get more than two or even one finger within the pelvis.

TABLE OF MEASUREMENTS IN NINE DISEASED Pelves.

No. in Mu- seum.	Inclined Plane of Promontory.				Plane of Brim.				Cavity.		Outlet.		Arch of Pubis.	Character of Pelvis.
	Prom. to Pubis.	Prom. to LPE	Prom. to RPE		Ant. Pos.	Tr'ns- verse.	Left Ob- lique.	Right Ob- lique.	Ant. Pos.	Tr'ns- verse.	Ant. Pos.	Tr'ns- verse.		
23	4½	3	2⅞		4½	4	3⅞	4	5½	3⅞	4¼	3¼	40	Cordiform.
24	3⅝	2¼	2¾		3⅞	3½	4½	4½	4⅝	3⅝	4⅝	3½	40	Ditto.
25	4½	2⅝	2⅝		4½	4	4½	4½	5½	3⅞	4⅝	1½	10	Ditto.
26	2	¾	⅝	⅝	2	2½	3¼	3¼	3½	2⅞	3½	2¾	10	Cast of Elizabeth Thomson's Pelvis, de- livered by Cæsarian Section, by Mr. Wood.
27	3	2⅝	2¾		3¼	5	4½	4½	3½	4½	4½	3⅝	70	Ovate.
28	2⅞	2¾	2½		2⅞	5½	4¾	4¾	3½	5	4	4⅝	85	Ditto.
29	2½	2½	2⅝		2¾	5¼	4¾	4¾	3½	5½	4⅝	4¾	90	Ditto.
30	1⅞	*1⅞	*1⅞		1⅞	4⅝	4¼	4⅝	2⅞	4⅝	2⅞	4⅝	85	Wooden Model.
31	1⅝	*1¼	*⅞		1¼	5	4¼	4½	2¼	5	2⅞	4½	100	Cast of Elizabeth Sherwood's Pelvis, de- livered by Crotchet, by Dr. Osborne.

* The oblique measurements of Pelves 8 and 9 are taken from the side not from the centre of promontory.

LECTURE III.

MECHANISM OF PARTURITION.

Definition of Labour—Divisions of Labours by various Authors—Division into Natural, Difficult, Preternatural, Complex—Definition of Natural Labour—Stages of Labour. Arrangement of the Muscular Fibres of the Uterus, on the External and Internal Surfaces. The Effect produced by the Contraction of the different Sets of Fibres—Manner in which the Dilatation of the Uterus is accomplished—Use of Liquor Amnii. Order of Uterine Contractions—Wigand's views—Objections to them—Dewees' Explanation of Uterine Action—Objections. Different Conditions of the Os Uteri—Dilatable—Rigid—Inflamed. Influence on the Action of the Uterus.

WE are now prepared to enter on the consideration of Parturition; a subject which embraces the most important questions we have to bring before you, and which will require your most serious attention, in order properly to understand the principles of midwifery.

At the termination of the period required for the complete development of the ovum, a new series of operations are entered on, for the purpose of giving birth to the fœtus which has been matured; these are included under the term Labour. It usually commences at the completion of the ninth month of gestation; in some instances it occurs before that time, when it is called Premature Labour. In this comprehensive sense, therefore, we would define Parturition to be—*the action of the uterus to expel its contents when the fœtus is sufficiently mature to sustain respiratory life.*

There are many circumstances depending either upon constitutional peculiarities, irregular formation, or upon accident, which may endanger parturition or render it dangerous; hence, labours have been divided and subdivided to meet these different conditions. Some adopt only two divisions. The first includes these labours which proceed regularly to their termination without in-

terruption. The second embraces those which do not do so. The one is the rule, the other the exception; but as the exception includes several varieties, this second class is subdivided into corresponding heads. Others place the most usual form of labour, termed *Natural*, in the first, and then add separately two, three, or four subsequent divisions, according to the importance which they attach to these deviations. In this manner, from two to seven divisions have been made, as you will perceive in the table which is placed before you.

DIVISIONS OF LABOURS BY VARIOUS AUTHORS.

Natural	Difficult	With 15 subdivisions or	.	.	.	Merriman
Eutochia	Dystochia	orders.	.	.	.	
Natural	Abnormal	With 6 subdivisions	.	.	.	Rigby
Eutochia	Dystochia		.	.	.	
Natural	Laborious	Preternatural	.	.	.	Deweese
Natural	Unnatural	with 6 orders.	Complex	.	.	Churchill
			with 6 ord.	.	.	
Natural	Difficult	Preternatural	Complex	.	.	Denman
Natural	Instrumental	Preternatural	Complex	.	.	Davis
Natural	Laborious	Preternatural	Anomalous	Flooding	.	Blundell
Natural	Premature	Preternatural	Tedious	Laborious	Impracticable.	Burns
					Complicated.	

These numerous divisions, to which many more might be added, are a sufficient reason for not wishing to complicate the subject by proposing another. It is preferable to adopt that which is most generally known, quite as distinct, and is in many respects more simple and practical, than some of those before you. Denman's division is sufficient for our purpose; we propose it to you for your selection, and shall consider labour under the several heads of *Natural*, *Difficult*, *Preternatural*, *Complex*. Denman defines labour to be *natural*, "if the head of the child present; if the labour be completed in twenty-four hours; and if artificial assistance be not required." Labour is called *preternatural*, when some other part than the head of the child presents. It is called *difficult labour*, when it exceeds twenty-four hours; and *complex labour*, when some accidental cause of danger occurs which may render interference necessary.

It is of the highest importance that you should have a clear view of the whole series of phenomena which constitute parturition; for, unless you perfectly comprehend the changes which are going forward in the uterus, and have an accurate knowledge of the means adopted by nature to accomplish her purpose,

you can never understand the principles of midwifery: your practice must be empirical; and however indebted to chance you may be for success, you will always be exposed to the risk of committing some fatal mistake. It, therefore, becomes our duty to explain to you the series of operations, by which the delivery of the child is effected. We would desire, at the same time, to direct your attention to the beautiful application of mechanical contrivance employed by nature to effect her object.

In order to study parturition efficiently, it is necessary to divide it into certain stages. The means by which the uterus is opened is not the same as that by which the child is forced through the pelvis; and again, the manner in which the placenta is separated and expelled is different from either; hence, labour has been divided into three stages, sometimes into four, and even five. The most usual division is that of Denman—the *first stage* being the dilatation of the os uteri; the *second*, the expulsion of the child; and the *third*, the separation of the placenta. Other authors subdivide the first into premonitory and dilating stages, and some divide the second stage into two, as you perceive in this table.

STAGES OF LABOUR BY DIFFERENT AUTHORS.

1st Stage.	2nd Stage.	3rd Stage.	4th Stage.	5th Stage.	Authors.
Premonitory	Dilating ...	Expulsive	Placental	Denman.
	Dilating ...	Expulsive	Placental	Velpeau.
	Dilating ...	Passage through the pelvis.	Expulsion	Placental	Jno. Clark.
	Dilating ...	Occiput under the arch of the pubis.	Expulsion	Placental	Merriman.
Premonitory	Dilating ...	Rupture of membranes and passage of the head.	Expulsion	Placental	Naegele.


Of these divisions, we would propose to you that of Denman; it is the most generally adopted, and seems to be that which is marked out by nature herself.

The *first stage* is dated from the opening of the os uteri to its complete dilatation. Denman adds to this definition, the rupture of the membranes and the discharge of the liquor amnii. This, however, renders the division imperfect, because the membranes often are not ruptured until late in the second stage, and occasionally remain unbroken when the head is expelled. This addition is therefore unnecessary, and causes confusion.

The second stage commences when the os uteri is perfectly dilated, and terminates in the expulsion of the child.

The third stage is occupied with the expulsion of the placenta.


Before describing the manner in which the dilatation of the uterus takes place, it is necessary to recapitulate very briefly the arrangements of the muscular fibres of the uterus, which has been detailed in a former lecture.




Muscular fibres on the external surface of the uterus, terminating in the round ligaments *a b*.

The fibres *on the external surface* form two broad fan-shaped muscular layers, spreading from the round ligaments over the fundus uteri. *On the internal surface*, there are three distinct sets of fibres: two of these surround the Fallopian tubes in a concentric arrangement. The third set pass circularly round the body of the uterus, and the outer fibres of the two former layers gradually pass into and intermix with those of the latter. The mass of fibres lying between the external and the internal layers have no determinate direction, but may be supposed to give increased power to those we have described. Sir C. Bell, in his valuable paper "On the Muscularity of the Uterus," has mentioned fibres which pass in a "vortiginous" direction from the fundus to the mouth of the uterus. Such, then, is the arrangement of the muscles or muscle of the uterus, as far as it has been demonstrated. Let us now consider their action. *The external muscular layer* slowly contracts for some time before labour has actually commenced, and draws the uterus gradually towards

the pelvis. By this means, also, the fundus is maintained in its proper direction, and prevented from inclining too much to either side. This gradual contraction is unaccompanied by pain, and therefore is not taken notice of; but its effect in altering the size of the abdomen, and making it less prominent, has always been observed, and noted as a premonitory sign of labour. These fibres also serve a useful purpose when the dilatation of the os uteri commences: the fundus being thus supported, the fibres on the internal surface contract more efficiently.




Exaggerated view of the fibres on the internal surface of the uterus:
a a, the orifices of the Fallopian tubes.



Uterus inverted, to show the *natural* appearance of the fibres on its internal surface.

The action of the *internal sets of fibres* requires a more careful examination, inasmuch as different and, it appears to me, incorrect explanations have been given both of the arrangement of these fibres and of the manner in which they contract. Besides this, it is necessary for you to have a correct view of the uterine action, and the order observed in these contractions, to enable you to notice the deviations from it that occur. We shall therefore consider each separately. *First, the effect produced by the contractions of the different sets of fibres. Secondly, the order in which these contractions take place.*

When the fibres surrounding the Fallopian tubes contract together, the fundus uteri is equally diminished on all sides, and their combined effect, conveyed through the medium of the liquor amnii, is precisely the same on the mouth of the uterus as if the fibres passed down vertically and acted directly upon it. In order to render this intelligible, I shall have to ask your patient attention for a few moments.



The lines *a, b*, represent the direction of the force of the fundal muscles: *c d*, the direction of that of the circular muscles of the body of the uterus: *d e*, the combined force of these muscles.

The dotted right lines represent the direction of the force reflected by the liquor amnii; the dotted curved lines, the direction of the circular fibres of the body of the uterus.

Let us suppose a line (*a b*) to pass from the opening of the Fallopian tube of one side of the uterus to the opposite, in such a manner as to represent the direction of the force of the fibres

surrounding it. This line would pass obliquely downwards to the opposite side. If, therefore, these fibres alone acted, the fundus on that side would be diminished in its size, and the contents of the uterus pressed toward the lower section of the opposite side, but not against the os uteri. If, further, two such lines (*a d*), passing from the orifice of each Fallopian tube represented the force of each set of these concentric fibres, the intersection of these lines would be the common point where these forces meet, and to a certain extent, are opposed. The combined force must, therefore, take an intermediate direction equi-distant from both lines, which would be represented by a line (*d e*) passing in the axis of the uterus, and through the os uteri. It follows, therefore, that when the fundus of the uterus contracts equally, the resulting force must be communicated to the os uteri, as perfectly as if the fibres passed vertically from the fundus to the mouth of the uterus.

When the circular fibres of the body and cervix contract, their tendency is to render the uterus more and more cylindrical, according to the degree of their contraction, at the same time that they close in the cervix. Again, if the uterus were emptied of its contents, the simultaneous action of all these different sets of fibres would be, to draw the parietes equally towards the centre of the cavity. But when the uterus is gravid, and makes an effort to expel the foetus, they cannot all contract in this manner.

The fundal muscles are those which chiefly effect the dilatation of the os uteri and the expulsion of the child, the fibres of the body and the cervix remaining comparatively passive. It has already been explained to you, that their united action is in the direction of the os uteri; but there is still a necessity for a means by which the result of that action should be perfectly conveyed to it. This is accomplished by the fluid enclosed within the amnion, which acts with a distending power upon the os uteri exactly equal to the combined force of these muscles. The muscular bands described by Sir C. Bell, must also have the effect of expanding the os uteri by drawing it upwards. The circular fibres of the body and cervix resist the efforts of the fundus to distend them; and the force of their resistance is also communicated to the contained fluid. This force is therefore, as it were, reflected upon the os uteri, so that the whole uterus

might be said to act as one muscle in dilating the mouth. The os uteri has been generally considered to possess circular fibres, its action to resemble, in some degree, the sphincter muscles in other situations, and its dilatation to be the effect of their relaxation. The existence of these circular fibres at the os uteri has never been proved. Hunter could not demonstrate them; Sir C. Bell could not trace them out.* If a sphincter muscle exist in that situation, why should not its fibres be as distinct and as easily made out as the sphincters of other hollow muscular viscera? Neither does the manner in which the os uteri dilates support this assertion. Its expansion is very gradual, it yields slowly to the power described, and does not resemble the comparatively sudden relaxation of a sphincter muscle. Were it similar, rapid labours (at least so far as the dilatation of the os uteri is concerned) would be the rule, not the exception. It would be incorrect, then, to attribute the resistance offered by the os uteri to muscular contraction. It seems more probable, that the firm, highly-condensed tissue that forms the cervix of the virgin uterus, still retains, in its altered state, many of its original characters; that this tissue, although more unfolded, is still sufficiently compact and elastic to offer a great degree of resistance, and that its dilatation is effected by the incessantly repeated efforts of the uterus slowly overcoming and expanding it. Such a view seems to be fully supported by the observation of labour in its first stage, where we can perceive the os uteri first

* "I have not succeeded in discovering circular fibres in the os tincæ corresponding in place and office with the sphincter of the other hollow viscera; and I am therefore inclined to believe, that, in the relaxation and opening of the orifice of the uterus, the change does not result from a relaxation of the muscular fibres surrounding the orifice. Indeed it is not reasonable to conceive, that the contents of the uterus are to be retained during the nine months of gestation by the action of a sphincter muscle. The loosening of the orifice, and the softening and relaxation which precedes labour, is quite unlike the yielding of a muscular ring." Sir C. Bell, on the Muscularity of the Uterus.—*Med. Chir. Trans.* vol. iv. p. 346.

Dr. William Hunter, whose perfect accuracy of description is now established, states, "The cervix uteri, where the penniform rugæ are situated, had not such regular nor so large fasciculi as the rest of the uterus. In the body of the uterus, the fibres are very regularly circular. The fundus was made up of two concentric circular planes of fibres, at the very centre of which was the orifice of the Fallopian tube."—*Anatomical Description of Gravid Uterus*, p. 27.

becoming thinner, and then progressively (not suddenly) opening. It has been stated that the sudden relaxation of the mouth of the uterus, after a continued contraction, can only be explained on the assumption of its muscularity. The same effect, however, may be observed in other tissues besides muscle; the perinæum, for instance, often resists, for a long time, the action of the uterus, and then yields suddenly to its full extent, so that the head is most unexpectedly delivered, and yet the distended portion of the perinæum is not muscular. Besides, these sudden dilatations are only exceptions to the general rule. In ordinary cases, dilatation occupies a certain period of labour (often a very long one), and the mouth of the uterus yields very gradually to the power employed.

We would now direct your attention to the means provided by nature, to prevent the danger which might arise during this process. If the uterus exerted its full power upon the undilated os uteri, and if the unyielding head of the child were driven forcibly against it, the almost certain consequence would be, that the irritation would excite increased resistance, and ultimately terminate in inflammation of the mouth of the uterus. To obviate such an effect, nature interposes a *fluid medium* between the power and the resistance. The liquor amnii contained within the membranes, occupies the cavity of the uterus, and when its parietes contract upon it, the force exerted is (as we have explained) by this means, accurately conveyed to the os uteri. When the latter dilates in the slightest degree, the fluid insinuates itself within the smallest opening, and expands it by a direct lateral pressure against its edges. The power of the uterus is thus made to act in the most favourable manner for distending its mouth.

Other advantages are also gained. The os uteri may dilate irregularly; but any attempt to overcome forcibly the undilated portion, is prevented when the force is conveyed through a fluid, which, while it readily yields to an undue resistance, still maintains an equable pressure upon the edges of the os uteri. Any irregularity in the action of the uterine fibres, is also, to a certain extent, obviated, because these contractions, although irregular, being still conveyed by the fluid, are thus equally communicated to the os uteri. Further, so long as the tissue of the uterus intervenes, it is necessary to moderate the great power

which the uterus is capable of exercising to dilate it : this is effected by the liquor amnii. The force conveyed by a fluid, you are aware, does not act in one direction only, but is distributed to every part of the surface to which the fluid is applied. The force, therefore, which is exerted to expand the mouth of the uterus, being communicated by a fluid, is not only directed against the os tincæ, but against the fundus and sides of that organ. The fundus, consequently, is opposed, not only by the os uteri, but by its own action reflected by the liquor amnii. Hence, so long as the fluid remains and the os uteri is undilated, the more powerful the action of the fundus, the greater is the resistance to it. The actual force employed, is therefore very moderate, and any sudden or violent effort at distension is altogether obviated. You may observe this in the character of the pains during this stage. You will find, that, however severely they may commence, they last but a short time, and the effect on the os uteri is comparatively slight. If these short, though severe pains, be contrasted with the long-continued and powerful pains which follow them when the liquor amnii is discharged, and the os uteri dilated, the difference in the effect will be sufficiently obvious. As a means, therefore, of conveying the whole muscular power of the uterus upon the os uteri—of moderating and equalizing the force employed—of dilating the mouth of the uterus without exciting irritation—the liquor amnii is of essential importance.

We shall now consider *the order* observed by the uterus in the contractions which take place. This may be ascertained experimentally. For instance : when the hand is passed into the uterus after delivery, to remove the placenta, we find that it may remain for some time in the cavity without exciting its contraction, but the moment the hand is being withdrawn, the fundus instantly contracts, and as it passes along the vagina, the contractions are continued from above downwards ; so also, in other instances, when the os uteri is only irritated by the fingers of the hand introduced into the vagina, and an attempt is made to dilate it, the fundus immediately contracts, not the os uteri. You have thus a very favourable illustration of the reflex nervous function. Hence we infer that the order of uterine contractions is from the fundus downwards, and that the action is commenced there.

In opposition to this view of the order of uterine contractions, there is the authority of Wigand, who gives an explanation altogether different from what we have stated. In order to place his opinion clearly before you, I shall quote the following passage from Dr. Rigby's work (p. 99), in which Wigand's views are faithfully given:—

“In examining the course of a true pain, we shall find the contractions of the uterus *do not begin in the fundus, but in the os uteri*, and pass from one to the other. *Every pain which commences in the fundus is abnormal*; and either arises from some derangement in the uterine action, or is sympathetic with some irritation not immediately connected with the uterus, as from colic, constipation, etc. We very seldom find that a contraction of the uterus which has commenced in the fundus, passes into the cervix and os uteri, and becomes a genuine effective pain; usually speaking, the contraction is confined to the circumference of the fundus, without detruding the fœtus at all. When a genuine pain comes on, so far from the head being pressed against the os uteri, it at first rises upwards, and sometimes gets even out of reach of the fingers, *whilst the os uteri itself is filled with the bladder of membranes*; if it had commenced in the fundus, instead of the inferior segment of the uterus, so far from the head being drawn up at the first coming on of the pain, it would have been forcibly pushed down against the os uteri. In the course of a few seconds, the contraction gradually spreads over the whole uterus, and is felt especially at the fundus; the head, which has been raised somewhat from the uteri, is now again pushed downwards to it, and seems to act as a wedge for the purpose of dilating it; it is not until the whole uterus is beginning to contract, that the patient has a sensation of pain. We may therefore, consider that a genuine uterine contraction consists of certain phenomena which occur in the following order:—First, the os uteri grows tight, and the presenting part rises somewhat from it, then the rest of the uterus, especially the fundus, becoming hard, the patient has a sensation of pain, and the presenting part of the child advances.”* Now, if we desired an additional evidence to prove that the fundus was the first part

* Wigand, op. cit. vol. ii. p. 197.

of the uterus to contract, and not the os uteri, we could not have a stronger proof than that advanced by Wigand to support a contrary opinion—viz., the head, when the contractions commence, getting, “even out of reach of the fingers, whilst the os uteri is filled with the bladder of membranes.” In Wigand’s explanation, the influence of fluid pressure seems to be altogether forgotten. The immediate effect of contraction commencing at the fundus would be to compress the liquor amnii which of necessity forces its way before the head, on to the mouth of the uterus. The fluid in this position re-acts against the head with the same power that it is compressed, and therefore pushes it up until the increasing contraction of the fundus forces the head down again, so that you perceive the phenomena quoted are quite consistent with the statement that uterine contraction begins at the fundus; in fact, it could not be otherwise, so long as the waters remain in the uterus. But if the contraction commenced from below, the fluid must be driven upwards towards the fundus, and that portion between the os uteri and head pressed aside, at least in the first instance, so that the head might be easily felt when the pain commences, although not so afterwards; but the reverse is the case, and you will find that in those cases where the liquor amnii is in large quantity, that it is difficult to feel the head at all, except in the interval of the pains. “The tightening of the os uteri,” alluded to by Wigand, seems to be another source of error on this point, it being generally confounded with muscular contraction of the os uteri. It seems to me to be produced by the pressure of the fluid downwards against the sides of the uterus, combined with the increased determination of blood towards the os uteri, which arises from the vessels at the fundus expelling a portion of their blood during its contraction. The os uteri is rendered fuller, and the lips are more closed than before; hence the opinion that it is muscular contraction, the evidence for which does not seem to me sufficient to establish so important a fact. Having placed before you the opinion of Wigand upon one point connected with the action of the uterus, it would be improper to omit the explanation given by another eminent authority, of the mode in which the uterus contracts. Dewees adopts a different view of the mechanism of dilatation from what has been stated to you, not a difference as to the order of uterine contraction, but as to the manner of it.

Dewees assumes that the longitudinal fibres which pass from the fundus to the os uteri, are the antagonists of circular fibres passing round the body of the uterus. The contraction of the former shortens the uterus, and distends the circular fibres, which, stimulated by this distension, contract, in opposition to the longitudinal fibres; that "the effect of this compound action is to direct the body to be moved towards that part of the organ which offers the least resistance to it, and this is the small opening called the os uteri; the fibres which surround this opening and oppose its immediate dilatation gradually become weakened by the superior strength and persevering action of the longitudinal fibres; and after a struggle of greater or less severity and duration, are obliged to yield; and in their quiescence the dilatation of the uterus consists. In the whole of this arrangement, we [Dewees] can see no necessity for the mechanical agency of the membranes acting on the circle of the os uteri like a wedge, which Dr. Denman speaks of as important to its dilatation, for every day's experience proves that the most perfect and most speedy relaxation of the mouth of the uterus takes place without any such influence."* In this explanation, Dewees has assumed what yet remains to be proved. We have no sufficient evidence that the arrangement of the fibres of the uterus into longitudinal and circular is as antagonist muscles;† and if we assume such to be the case, and

* Dewees, p. 181.

† When these Lectures first appeared in the "Lancet," it had been stated inadvertently that there was no evidence of the *existence* of longitudinal and circular fibres. The mistake was not perceived until they were again revised for publication. It was not intended to deny the existence of these fibres, but the kind of action assigned to them by Dewees,—that of antagonist muscles. The description given by different observers of the situation and arrangement of the uterine fibres being dissimilar, it was our wish to confine the attention of the student to those only which were the most obvious, and which are quite adequate to explain the action of the uterus, viz., the external layers which draw down the fundus of the uterus, the concentric fibres round the orifices of the Fallopian tubes, and the circular fibres of the body of the uterus, both of which may be observed on the internal surface. The explanation we have given of their action nearly corresponds with that of Jobert, who says:—"The direction of the uterine fibres shows how they act in freeing the uterus from its contents. The longitudinal layer of fibres, which originates at the fundus, and is inserted in the neck and vagina, tends to diminish the length of the uterus, whilst the semicircular fibres, by their action, diminish its cavity in every sense."—*Lancet*, vol. i. p. 732. The semicircular fibres that Jobert alludes to, are the concentric fibres at the fundus, and those round the body

that they act in opposition to each other, the effect of the collision must be communicated to the liquor amnii; but how the membranes could resist the effect of this struggle it would be difficult to understand, when we know that change of position, walking across the room, or other such trifling causes, are sometimes sufficient to rupture them, from the gravitation of the fluid alone, and therefore the greater force arising from the action of the fibres of the uterus against each other, must break them much more frequently than what we know to be the case. Besides this, Dewees' explanation proves too much. It would make the membranes not only useless as a means of dilatation, but rather a difficulty in its accomplishment. The head of the child, directly applied to the cervix would overcome the resistance of the circular fibres much more efficiently than the liquor amnii, so that the most favourable kind of dilatation would be that which occurs when the membranes are ruptured at the commencement of labour. It is very well known that this does not happen; and Dewees seems to be quite aware of the difficulty which this fact presented. He attempts to get rid of it by another assumption, which is, that the uterus is surprised in action by this circumstance, before the natural stimulus is given to it. "In consequence of this, the uterus is made to embrace the child closely, by virtue of its tonic power, and is by this thrown into the most violent and painful contractions, very often by the unequal surface which the child's body presents, the evacuation of the waters preventing the lower part of the uterus from being fully stretched, by the contractions of the body and fundus forcing them (the waters) down against it."* But in place of the waters, there is the large, round, and unyielding head of the child forced down upon the lower part of the uterus, which one would suppose more efficient for the purpose of mechanical distension.

We have felt it our duty to place before you those opposite views as to the dilatation of the mouth of the uterus, lest you

of the uterus, which he states extend down to the neck. The only difference between this explanation and that given in the text, is that the circular fibres of the body are supposed to remain passive, merely resisting distension, during the passage of the child, and that the simultaneous contraction of all the fibres does not take place until afterwards, when the uterus is empty.

* Dewees, p. 182.

might hereafter be confused by finding explanations of the action of the uterus, supported by respectable authorities, which are in contradiction to what we have stated to you, and which we had passed by unnoticed. We have already pointed out the advantage of the liquor amnii; hence you will find that the time at which the membranes are ruptured often makes a material difference in the effect produced upon the mouth of the uterus. If they are ruptured when the dilatation is very slight, the suddenly increased power of the fundus, forcing the head of the child against the os tincæ, soon excites irritation, prevents expansion, and sometimes causes inflammation. But if they are broken when the uterus is sufficiently open to allow the membranes to protrude into the vagina, and the contractions of the fundus to increase, it is probable that the dilatation will be advanced more rapidly, because of the diminished resistance from below, and the increased force from above. This is not, however, invariable. It occasionally happens that, even under these circumstances, the os uteri becomes irritated and retarded in its dilatation.

If we have succeeded in making you acquainted with the mode in which the uterus acts in dilating the os tincæ, and the means provided for controlling the great power it is capable of exercising, we would now direct your attention to the nature of the resistance presented to it—viz., to the os uteri itself. This part of the uterus varies greatly in its density and firmness; it consequently offers different degrees of resistance to the fundus uteri. You will recollect that the os uteri, during gestation, has gradually undergone certain changes, preparatory to its dilatation. The highly condensed cellular tissue of which it consists has become looser, and is traversed by more numerous vessels. The cervix is nearly, if not altogether, obliterated, and the circular orifice of the os uteri alone remains. Its edges may be either thick, full and soft, or extremely thin, according to the degree to which its cellular tissue is unfolded. They are always moistened with the viscid mucus, which is secreted so abundantly at this time. If the fingers be passed within the os uteri, and separated, the edges yield readily to a moderate pressure, there is a very slight increase of temperature, and there is no tenderness or pain produced when the os uteri is touched. This is the most favourable state of its dilatation; it is quite prepared to yield to the action of the uterus, and is called, in obstetric language, *the dilatable os uteri*.

There are, however, many exceptions to this condition, varying with the degree to which the density of structure in the os uteri may be increased. The cellular tissue is never so loose and permeable in the first instance as it becomes afterwards; the mouth of the uterus is therefore more resisting in first than in subsequent pregnancies. Its structure retains more of its elasticity and firmness in young women pregnant for the first time, and consequently much more time is occupied in unfolding it; hence the first stage of labour is always longer in primiparæ than in those who have had many children. The os uteri is still more firm and resisting, if, in addition to a first pregnancy, the woman be advanced in years: the cervix and os uteri remain close, compact and impermeable to the moment of parturition, which may be attributed to the increased firmness and diminished vascularity which age produces in the tissues generally. It then obtains the name of *rigid os uteri*. But there are different degrees of rigidity. Sometimes the structure is only *tough*. It gives way very slowly to the action of the uterus—nevertheless it yields, although, as it were, reluctantly. In such cases, the os uteri may remain cool and free from tenderness, but opposes a firm resistance to the pressure of the finger, and always requires a long time before the dilatation is accomplished. There are, however, a certain class of cases in which this condition of the uterus is in the extreme. It might almost be called *the undilatable os uteri*. In this state its structure is unusually dense, and feels like cartilage. The edge of the os uteri is perfectly unyielding; when thick, it might be compared to the feel of Gimbernat's ligament. If very thin, it still offers the same resistance, and is to the touch like a hole made in parchment. Instances of this extreme rigidity are met with not only in women who are advanced in life, but in those who have been all their lives accustomed to much bodily exertion, and exposed to the vicissitudes of laborious occupations. They are generally hard-featured, coarse-skinned, muscular women, of low stature, with thick short fingers, large wrists, and the bones generally prominent. It is in these cases you meet with that form of pelvis that I have described to you, as possessing many of the characters of the male pelvis. If you should, unfortunately, meet with a case of this kind, you must be prepared for difficulties from the commencement to the termination of labour, and therefore the consideration of it deserves your closest attention.

All these varieties are included under the term "rigidity." But beside, there are cases where the os uteri becomes *rigid*, *although previously dilatable*. If the os uteri become inflamed, rigidity is the result of it; the os tinæ grows hot and tender, is swollen, and becomes rigid. This alteration may arise from any irritation: premature rupture of the membranes for instance, by which the head is brought into direct contact with the undilated os uteri. It is also often induced, not by accidental causes, but by too much meddling, making too frequent examinations, attempting to dilate the os uteri artificially, etc. You cannot, therefore, be too cautious in this respect. Sometimes the head of the child presses so unequally upon the os uteri as to excite inflammation in it. The head may not be directed exactly in the axis of the brim, but may rather rest upon the pubic portion of it, compressing the anterior lip of the uterus with every pain. While the remaining portion of the mouth of the uterus expands, this remains undilated, and forms a band in front of the head. When the membranes are ruptured, the pressure is so much increased that the anterior lip often inflames and grows quite rigid. Again, there are cases where the os uteri is driven down with the head into the pelvic cavity, and the whole circle of the os tinæ compressed so tightly against the pelvis as to produce inflammation; further dilatation is arrested, the os uteri is rigid, and if it remain long in this condition, slough may be the result; the whole os tinæ has been completely separated in this manner, and expelled with the head of the child.

In conclusion, we would direct your attention to the difference in the action of the uterus, when it has to overcome an unusual opposition, arising from this state of rigidity. The contractions take place continuously for a certain time; but when the period which is usual to effect the dilatation is exceeded, or when the os uteri becomes irritated, the pains grow feeble, and the uterus often suspends its action altogether. By this means an interval of rest is gained, the irritation may subside, the patient may get some sleep, and recover from her fatigue, which otherwise might end in exhaustion. When the action of the uterus is renewed after a suspension of this kind, the dilatation is often rapidly completed. Much confusion has arisen as to the duration of labour, in consequence of neglecting this fact. Its commencement is generally dated from the sanguineous discharge (*the show*)

which marks the first opening of the os uteri. But if the first stage occupy a very long time, including these intervals of suspension, some altogether discard the previous irregular labour, and date its commencement from the time that the pains return regularly and continuously. Thus, a labour which one author would describe as being very much prolonged, another might bring within the usual limit of twenty-four hours, meaning by this twenty-four hours' continuous labor; and hence you will find much contrariety on this point amongst obstetric authorities. At a future time we may have to refer to this; at present we would desire to point out to you the conclusion which is deduced from this irregularity in the uterine action. It affords an additional illustration of the principle which Nature seems to observe in the dilatation of the os uteri—*to do nothing by violence*. In all ordinary cases, the liquor amnii moderates the action of the uterus; but if there be an unusual resistance offered to it, and the waters are discharged, the increased action does not continue; it is suspended, and again renewed; so that the object is obviously to accomplish by time what Nature avoids effecting by force.


LECTURE IV.

MECHANISM OF PARTURITION, CONCLUDED.

Second Stage of Natural Labour—Bearing Pains—Pains—Passage of the Head through the Pelvis—Positions of the Head according to various Authors—Naegele's Views—Mode of distinguishing Vertex Positions—Varieties in Face Presentations—Characters of—not so dangerous as had been supposed—Dilatation of the Perinæum—Expulsion of the Head—Direction of the Shoulders. *Third Stage*—Separation of the Placenta—Causes of Retention—without Hæmorrhage.

At our last meeting, we considered the means by which the dilatation of the mouth of the uterus was effected; we have now to examine the manner in which the child passes through and is expelled from the cavity of the pelvis. This is the *second stage of labour*. So long as the tissue of the uterus was interposed to the advancing head, the design of nature was evidently to moderate the action of that organ, and to prevent too violent a distension of its structure. But when this no longer arrests its progress, and the pelvis becomes the impediment, the full power of the uterus is exercised to force the head through the osseous cavity which resists its advance. You will therefore observe a marked difference in the character of the uterine contractions. Not only is the entire force of the uterus employed, but it is aided by the muscles that bound the abdominal cavity. This change will explain to you a difference in the character of the pains which are the effect of these contractions. In the first stage, they are sharp, severe, but short in their duration. They are called, in obstetric language, "*grinding pains*." In the second stage, they are less acute, perhaps dull, but are steady and long-continued; a full inspiration is taken previous to their commencement, and the pain is expressed by a gradual expiration, accompanied by a deep tone of voice. These pains are called "*bearing pains*," and their deep continuous groan forms

a strong contrast to the shrill and almost agonizing cry that accompanies the grinding pains. When the action of the uterus is so much increased, you must perceive the obvious hazard that would arise if the adaptation of the head to the pelvis were not exact; if the former were too large, or the latter too small, serious danger might be the consequence. Hence, the second stage of labour, and the passage of the head through the pelvis, requires your closest attention. It is necessary for you to un-



The head above the brim of the Pelvis; the anterior and posterior fontanelles being nearly on the same level.

Note —In order to render the relation of the Head and Pelvis more intelligible, the pubic side of the pelvis is represented as being transparent in this and the following views.


derstand all its varieties, not in theory only, but also to take every opportunity to make yourself *practically* acquainted with them.

We would urge this upon you with the greater earnestness, because it too frequently happens that the practitioner is satisfied if he can distinguish the head, without caring much about its position, and hence decides upon the necessity for interference, not by his knowledge of the cause of the difficulty, but the length of time this stage may occupy. We would therefore recall to your recollection the observations which we made when

describing to you the obstetric anatomy of the pelvis ; we then pointed out to you the influence of the varieties of its shape on the progress of the head of the child, and showed you that its passage was accomplished by a combination of four distinct motions harmonizing in one effect. These we shall briefly recapitulate.

1st. When the head is above the brim of the pelvis, the forehead and occiput are nearly on the same level, but when the head enters the brim, the occiput descends lower than the sinciput, and a certain distance along the plane of the ischium against which it rests. The forehead then advances more rapidly at the opposite side of the pelvis, until it is arrested by the convergence of the ischium and shorter sacro-ischiatic ligament. The occiput again descends obliquely along the ischio-pubic ramus, and emerges with part of the parietal bone beneath the pubic arch. The head, therefore, might be said to oscillate on its bi-parietal measurement.


2nd. Simultaneous with this motion, there is a very slight rotation on the longitudinal axis of the head, by which that side of the head which is next to the pubis descends lower than that near the sacrum, so that the parietal protuberance of the pubic side becomes the presenting part.



The head within the pelvic cavity. The occiput and right parietal bone being the most dependent parts, the occiput resting against the plane of the ischium and obturator space, and the ear to the right of the symphysis pubis.

3rd. As the head is so advancing through the pelvic cavity, the shape of the pelvis obliges it to pass in a spiral direction; hence the head, which may enter the pelvis in the oblique or transverse measurement, is turned as it descends, towards the conjugate axis.

4th. When the head escapes from the outlet, the occiput rests against the ramus of the pubis and ischium, and becomes a fixed point, round which the remaining portion of the head successively passes out. In some instances, the occiput rests directly against the arch of the pubis, and the head is expelled in the conjugate axis of the outlet. According to its more usual course, it passes out obliquely.



Further advance of the head through the pelvic cavity, previous to emerging from the arch of the pubis. The parietal bone, and part of the occiput presenting at the arch of the pubis—the head more in the antero-posterior direction, and the anterior again descending to the level of the posterior fontanelle.

The head, however, does not descend always in the same position, and there are accidental displacements that may retard its progress. It is necessary, therefore, to understand these deviations. Formerly, the varieties in the position of the head received but little attention. The older writers only observed the manner in which the child was expelled (Sir Fielding Ould, however, is a noble exception.) They found the occiput generally towards the pubis, when the head escaped from the

vulva, but sometimes the face; hence they made only two divisions, or, to speak more correctly, they considered the former to be the rule, the latter an accidental exception to it. Baudeloque, however, observed the progress of the head while it was within the pelvis, and determined the position by touch, not by sight. He described six different positions; and, since his work appeared, every division that has been made, whether into four, six, or eight positions, has been formed on the basis which he has laid down—that of determining the position while the head was in the pelvis, and not when it was expelled from it. Authors have not agreed upon the number of these positions, nor upon their order of frequency. It will be necessary, therefore, in order to prevent confusion, to arrange their divisions in a tabular form, before describing the positions, and the mode of ascertaining them. The terms employed require a brief explanation. The position of the head is generally determined by that part of it which corresponds to the plane of the ischium, and therefore is opposite to the cotyloid cavity. If the occiput be in that situation, it is called the occipito-cotyloid position; if the forehead, the fronto-cotyloid. In the same sense you may have the occipito or fronto-pubic, and occipito-iliac; when the occiput, or forehead, is applied to the pubis in place of the ischium, or when the occiput is opposite the centre of the ilium. The occiput, also, may be to the left or right side of the pelvis; and thus you have the right occipito-cotyloid position, the left occipito-cotyloid position, and so on.

In the table before you, the numbers are placed in the order adopted by each author.

Left occipito- cotyloid	Right occipito- cotyloid	Occipito- pubic.	Left fronto- cotyloid.	Right fronto- cotyloid.	Fronto- pubic.	Left occipito- iliac.	Right occipito- iliac.	
1	2	3	4	5	6	} Baudeloque, Dewees, etc. Lachapelle Naegele Ramsbotham.
1	2	...	3	4	...	5	6	
1	3	...	2	4	
3	4	7	5	6	8	1	2	

Nearly all authors agree that the most usual position for the head to pass is with the occiput corresponding to the left plane of the ischium. It is then called *the first position (or left occipito-cotyloid.)* When the head is in this situation, the occiput has the relation to the ischium which we have stated. The forehead

corresponds to the right sacro-iliac synchondrosis. The right parietal bone is lower than the left, and its protuberance the lowest part of the head, which presents. As the head advances, the occiput becomes gradually more anterior, and the forehead dips down a certain distance in the pelvis, but the parietal bone remains unaltered. As it approaches the outlet, the forehead ceases to advance, and the occiput and parietal bones press down upon the perinæum to the utmost extent, descending along the ischio-pubic ramus, in order to emerge under the arch of the pubis. In making a vaginal examination to ascertain this position, the finger first touches the parietal protuberance; the sagittal suture is felt close to it, directed obliquely backwards. Anteriorly, and to the left side, this suture terminates in the posterior fontanelle; but the anterior fontanelle cannot yet be felt at the opposite side. The first position is chiefly determined by this situation of the posterior fontanelle; you should, therefore, recollect its distinctive characters, which we shall presently point out to you.

In the *second position* of Baudeloque (or right occipito-cotyloid), the occiput is opposite to the right plane of the ischium; the posterior fontanelle is, therefore, in the same relation to the right side of the pelvis that it was to the left in the first position, being anterior to its transverse axis. The sagittal suture passes obliquely backwards, from right to left; the left parietal bone is on the pubic side, and descends lower than the right. Dr. F. Ramsbotham places the transverse position of the head as first and second in the order of frequency, and considers the occiput more frequently opposite the ilium, at the right or left extremity of the transverse axis, than the plane of the ischium. If such were the case, the ear would be felt directly behind the symphysis pubis, in place of the right or left groin. It is certainly true that it is found more frequently near to the former than in the latter situation, which gives a plausible support to his opinion. Authors seem to have placed the oblique position of the head before the transverse, in frequency, from the circumstance, that in the recent pelvis the psoas muscles diminish the transverse measurement without affecting the oblique. It seems to me, however to be doubtful, whether, in a pelvis of moderate width transversely, these muscles interfere much with the head in its progress, especially as they are always as much

relaxed as possible during labour, and therefore yield to the pressure of the head. Judging from the opportunities of observation I have had, the head seems to be placed most frequently in the left oblique occipito-cotyloid position, but not so much so as to remove the ear from the neighbourhood of the pubis. In the third and sixth position of Baudeloque, the sagittal suture corresponds nearly to the conjugate measurement of the brim of the pelvis, only that in the former the posterior, and in the latter the anterior fontanelle, are opposite the pubis: these positions are very rare, and are omitted by Lachapelle and others, who consider *the third position (or left fronto-cotyloid)* as the converse of the first. The frontal bone and anterior fontanelle correspond to the left plane of the ischium; the sagittal suture passes backwards from left to right; the posterior fontanelle is opposite, and near the right sacro-iliac synchondrosis; the left parietal protuberance is the most dependent point, and the ear is situated as in the second position, only more withdrawn from the pubis, and nearer the groin. This is Baudeloque's fourth position.

The *fourth position (or right fronto-cotyloid*—fifth of Baudeloque) has the frontal bone opposite the right plane of the ischium; the sagittal suture passes backwards, from right to left; the right parietal bone presents, and the right ear is opposite the left groin. Such are the principal positions, as they are given in the different works on midwifery. The diagnostic marks by which they are distinguished are, first, *the fontanelles*. In order to discriminate between occipito and fronto-cotyloid positions, you must understand the character of the anterior and posterior fontanelles, judging of them by touch, not by sight. In the dried bone, the posterior fontanelle is a triangular space. In the head of the child, the finger will often detect no space, but only a point, the centre in which three lines of suture terminate. The anterior fontanelle, being so much larger, lozenge-shaped, and having four lines of sutures terminating at its angles, can be easily distinguished from the posterior, if the eye were to decide the question; but, as we must judge by the finger alone, these characters are not always obvious. The anterior fontanelle is higher up, and more removed from the finger in fronto-cotyloid positions than the posterior fontanelle is in the first and second positions; consequently, the finger cannot reach

sufficiently far to trace its exact shape. The divergences of the parietal bones may be felt, and perhaps the coronal sutures, but the two remaining sides of the lozenge formed by the frontal bone are too distant, neither can the frontal suture be felt; hence there is some resemblance to the triangular space of the posterior fontanelle, the only difference being in the greater size of the triangle in the anterior fontanelle: but in cases where the ossification of these bones advances slowly, the posterior fontanelle is large, and might resemble the anterior. You perceive, therefore, that it is possible to confound the one with the other, and that it requires practice in examining by touch, in order to discriminate them in all cases. As a general rule, however, you will find the posterior fontanelle like a central point, in which three lines meet, the anterior more distinctly a membranous space, but usually undefined. If there be any difficulty, it may be removed by a careful examination of the second diagnostic mark — *the ear*, which may be felt on the pubic side of the pelvis, except in cases of great disproportion. As the lobe of the ear is always nearest to the occiput, you can by it determine the side of the pelvis where the occiput lies, and therefore you can distinguish the two positions which have the occiput on the left side, from the two that have the frontal bone in the same situation; but you cannot so easily decide between collateral positions the first and fourth, or the second and third. It has been stated to you, that authors have not agreed as to the second position. Baudeloque placed the right occipito-cotyloid position second in the order of frequency, and was followed by several other writers, until Naegele made these positions the subject of his observation. The result of his examinations led him to doubt the accuracy of the description which had been given, and ultimately to deny it altogether. He found, that although the head was expelled in Baudeloque's second position, it did not enter the brim of the pelvis in that direction, but that it passed down first in the third position, with the occiput towards the right sacro-iliac synchondrosis, and, as it descended, rotated gradually into Baudeloque's second position, in which it was expelled. Thus the occiput, and of course the head, might be said to describe a kind of spiral curve from right to left, as it passed through the pelvis. Naegele's explanation has been since confirmed by other observers, and is

in the main correct—a conclusion which I have arrived at from personal observation. Being anxious to determine this question, I availed myself of the opportunities afforded me in the Dublin Lying-in Hospital, of putting it to the test, and found that in nearly an equal number of cases the head entered the brim in the third position as in the second, that of those which descended in the third, the majority passed without any difficulty into the second, and were so expelled, while a very few remained in their original position. The whole evidence establishes Naegele's accuracy of description, and it may be admitted that, as a general rule, the head rotates from the third into the second position, when it is passing through the pelvis; but there are exceptions to it. The fourth position also passes into the first; but sometimes, although very rarely, preserves its original direction.*

From these varieties of the vertex presentation, we would now direct your attention to some deviations from them which are occasionally met with. It has been stated to you, that in one of the motions of the head of the child, the frontal bone descends to the level of the occiput. In some instances, however, from an accidental cause, the forehead is driven down too far, that the head becoming fixed in the pelvis transversely, its progress is thus arrested. It is essential for you to understand this cause of delay in the second stage, because it is very easily corrected; and any ignorance respecting it might lead you into the erroneous impression that the head should be delivered by instruments, because it was so long fixed in its position. When this accident takes place, the anterior fontanelle may be observed to be remarkably distinct; you readily trace out its lozenge shape, and feel the four sutures distinctly at each of the angles. The finger also passes very easily between the pubis and the head, so that there appears to be rather more space in that situation than usual. Finding, therefore, this evidence of room in the pelvis, the anterior fontanelle perfectly within

* Of seventy-four cases recently examined the following results were noted:—

First Position.	Fourth into First.	Second Position.	Third into Second.	Third Position.	Irregular.	Face.
43	3	11	9	2	4	2

reach, and, at the same time, the head not advancing, you have sufficient proof of this deviation. It is described by many authors as the *premature separation of the chin from the chest of the child*. The mode of correcting it is sufficiently simple. The head should be dislodged from its position in the interval of a pain, and the fingers pressed against the frontal bone until the uterus again contracts; the occiput will at once descend, and the labour proceed without further difficulty. There are rare instances, in which the forehead descends completely into the cavity of the pelvis, and becomes the presenting part; when this occurs, the displacement cannot be corrected, as in the former instance, and the position is so unfavourable, that the head soon becomes arrested in its progress. It acts upon the pelvis something like a wedge, the forehead being the narrow end, and the occiput and neck of the child its broad base. The more the head advances, the more the difficulty is increased. When this accident takes place, the exploring finger feels the frontal bone traversed in the centre by the frontal suture; there is, therefore, some resemblance to the vertex position, but it may be easily distinguished, because the forehead is smaller than the occiput, and the anterior fontanelle being near, can be very easily traced. The inexperienced observer would imagine that there was more space than usual in the pelvis for the passage of the head, and might not be able to explain why its progress should be delayed; but if the finger were passed sufficiently high, he would soon perceive that the head was completely wedged in the pelvic cavity.

Sometimes there is a complete rotation of the head on its transverse axis, so that the forehead rests upon the brim of the pelvis, and the face descends into the cavity. This constitutes the *face presentation*. Two varieties of this position are generally met with. In one, the chin corresponds to the right plane of the ischium (the *right mento-cotyloid position*). In the other, the chin has the same relation on the left side (*left mento-cotyloid*). When the former position presents, the *right malar* bone and cheek are the most dependent parts, the mouth and nose being rather posterior on the sacral side. In the latter the *left side of the face* presents. In both instances, there is the same obliquity observed as in vertex positions. Two other varieties are described, where the chin is directed backwards towards the

sacro-iliac synchondrosis ; but I believe that they are very rarely, if ever, met with.

If we are correct in considering face positions as the result of a rotation of the vertex, the manner in which they occur might be explained in this way : so long as the head is above the brim of the pelvis, its position can always be very easily altered, and it frequently changes from one into another. If it should happen that in any of these alterations, the occiput is placed upon the brim of the pelvis, it may glide from it towards the ilium, the forehead take its place, and the face descend as described. If this were true, the first position of the face would be the result of a rotation of the fourth position of the vertex, which rarely enters the pelvis ; and the second of the face, a rotation of the third of the vertex.

It is more important, however, to consider the mode of distinguishing these positions, because the situation of the face is such that it is exposed to great tumefaction. The face is much more vascular than the vertex ; and being the most dependent part, and very much compressed, the cheek, the eye, and the portion of the mouth which presents, are always very much swollen. Even with the most careful management, the features of the child are more or less disfigured, but this may be greatly increased by frequent examinations. Unfortunately, the infant is too often a severe sufferer from this cause. The unusual characters of the position lead the inexperienced practitioner to make several fruitless attempts to determine what it is ; the tumefaction is much greater than before, and the outline of the features more confused ; thus serious injury to the child is sometimes the result : the eye has been destroyed in this way.

In examining a face presentation, the first object that attracts your attention is, the *irregularity of the soft surface* which presents itself. It may seem to be very easy to determine the features of the infant ; but you will not find it so. The soft cheek and the malar bone have some resemblance both to the breech and shoulder, neither can you define the outlines of the features with the same accuracy by touch as by sight : it is this difficulty that leads to the frequent examinations that are made in these cases. You generally first feel the nose like a soft fleshy elevation : if this be cautiously traced, the finger passes on one side to the bridge of the nasal bones, and on the opposite, to the lips

and mouth. This may be taken as a diagnostic mark of the position. When the finger passes from the lips to the gums, and from thence to the tongue, the sensation communicated by the firm ridge of gums, cannot be mistaken.

These face presentations were formerly considered to be very unfavourable, and to give rise to great delay and difficulty in the progress of labour; but this impression seems to have been created much more from the novelty of the position than from experience. In the extensive hospital at Vienna, Boer noted eighty face presentations, and observes, "there were three, or at the most four, where the children were born dead. None of the patients suffered in the slightest degree from any of these labours; and, *except one case*, all were left entirely to nature. In one case only, *on account of the weakness of the pains* and doubtful character of the symptoms, I deemed it necessary to terminate the labour by the forceps.—(Rigby, p. 130.) In the Dublin Lying-in Hospital, Dr. Collins gives thirty-three cases in his report,* and states, "Four of the thirty-three were still-born. With the first, the labour lasted thirty-six hours (the only instance); the second was an acephalous fœtus; with the third, the labour lasted eight hours; with the fourth seven hours: all were delivered without assistance." Of these thirty-three cases, thirty-one were delivered within twelve hours; and Dr. Collins further adds, in explanation of the small proportion thirty-three cases bear to the total number in the report, 16,654: "Some cases of face presentation, I am disposed to think, were not noted, *delivery having taken place so very speedily* as to excite little attention, and to cause it to be overlooked." (p. 34.) These statements coincide with my own observations in the same hospital. The total number of cases which came under my notice during three years, and the results of which had been noted, were 5,699; of these there was not a single face-position which required aid in the delivery, nor did the labour in any occupy twenty-four hours. The only danger, therefore, which might arise from these positions is the danger of meddling too much with them.

When the head (which we will suppose to be in the first posi-

* Collins' Report, p. 83

tion) approaches the outlet of the pelvis, and descends along the ischio-pubic ramus, it is opposed by *the perinæum*. When this takes place, every provision is made to obviate the effects of the struggle about to ensue. The mucous secretion flows much more abundantly from the vagina; the presenting part at first just touches the perinæum and then retreats from it, and the action of the uterus seems again to alter its character; the pains are not so vigorous as before, but for some time are shorter in their duration, and apparently less efficient. The same caution is exercised to guard against violence here as in the first stage, and you will often find the head or the membranes advancing and retreating for a long time before it descends so far as to allow the occiput and parietal bone to pass between the pillars of the pubic arch. During this period the structure of the perinæum becomes more and more unfolded, so that the delay which usually takes place in the labour at this point seems to arise much more from the feeble character of the pains than the resistance of the perinæum. At length, when the head is enabled to pass down so far that the occipital portion can emerge from the outlet, the perinæum suffers its greatest degree of distension. The occiput then first rests against the ischio-pubic ramus, and as it is expelled, rises upwards towards the pubic arch. It becomes a fixed centre, round which the remaining portion of the head moves; the whole force of every pain is, therefore, spent upon the perinæum; and if these return with much rapidity or violence, there is a danger that this part may be lacerated. In the majority of cases, however, the head retreats in the intervals of the pains, even where it seems to be almost expelled. At this time, also, the pains become much stronger, and are the severest and most trying to the patient of any that she has experienced. At the conclusion of this stage, one or two very strong pains take place, by which the head is first protruded so far that it does not retreat, and it is then delivered. As soon as the head is born, you are given a favourable opportunity of confirming your diagnosis as to the position. In this, the first position, the face is directed obliquely upwards towards the right thigh of the mother, because when the shoulders and body of the child pass into the pelvic cavity, they enter the brim of the pelvis in the oblique measurement, opposite to that in which the head has passed, and therefore as the child goes through the cavity of the

pelvis, the thorax and abdomen correspond to the right or superior side of the pelvis, and the face looks in the same direction.

When the shoulders are being expelled, the perinæum is again put upon the stretch, and is sometimes unequally protruded by the arms of the child, by which, if care be not taken, it might be lacerated—but as soon as they escape there is no further danger. The uterus also ceases to act with the same force, the remaining pains are weaker, so that the body and limbs of the child are very slowly expelled. Such is the manner in which the child, in its most usual position, is delivered, and this stage completed. But before entering upon the consideration of *the third stage*, we would direct your attention to the mode in which other positions of the head are expelled, and their influence on the perinæum. When the head passes out in the second position, the back of the child is to the right side of the pelvis, in the same relation as the thorax and abdomen are in the first position; the face, therefore, looks downwards to the left thigh. When the shoulders are passing the perinæum in this position, great caution is necessary, because you do not feel the hands and arms coming out, as in the first position; they might therefore press on the perinæum unequally, and lacerate it without your knowledge. When the face is towards the pubis, the perinæum is very much endangered, in consequence of the greater distance to which the head must descend before it can pass under the pubic arch; and in order to do so, the direction of its motion is downwards and backwards, the reverse of what usually occurs; consequently, the perinæum suffers a much greater degree of tension than in ordinary cases, and there is a proportionate risk that it will give way. In face-positions, from a similar cause, the perinæum also suffers, but not to the same extent, because the tension upon it is only continued until the chin passes under the arch, when the pressure is at once relieved.

As soon as the expulsion of the child is accomplished, the uterus ceases to contract for some time, the interval varying from five to fifteen minutes; then the contractions are renewed, for the expulsion of the placenta. This constitutes *the third stage of labour*. These can scarcely be called pains; they bear no resemblance to those which preceded them, and are but little noticed by the patient. The manner in which the placenta is separated depends

very much upon the mode in which the uterus contracts while expelling the body and limbs of the child. If the fundus receive due support from the abdominal muscles, so as to ensure a uniform and efficient contraction from above downwards, it often happens that the same pain which extruded the limbs of the child from the vulva, expels the placenta from the uterus into the vagina, where it may remain for some time. The same will occur if artificial support be given by pressure with the hand on the fundus; in either case, the uterine contractions have no further effect upon it. It would continue for some time in this situation if left to itself, and either be gradually removed from it by the slow action of the fibres of the vagina, or suddenly, by some shock from above driving the uterus down upon it; as, for instance, coughing, retching, or much forcing with the slight pains which occur at this time. It is seldom, however, suffered to remain, but is generally drawn away by the funis. *Retention of the placenta* may arise from different causes. Sometimes the sphincter of the vagina closes upon it, and the placenta is thus retained until removed by the hand, or by firm pressure on the fundus of the uterus. In other instances the *placenta remains in the uterus* after the delivery of the child, until it is expelled by its subsequent contractions, rendered efficient by similar contractions of the diaphragm and abdominal muscles. This additional aid is required, inasmuch as the action of the uterus alone is not sufficient for the purpose. Hence, when the abdominal muscles are feeble, so that the uterus can derive no support from them, *the placenta is retained in this cavity*. This cause of retention has been generally mistaken for *inertia* of the uterus; and under this impression the placenta has been, very unnecessarily, withdrawn from the uterine cavity. It is therefore necessary for me to direct your attention especially to this point, in order that you may understand the principle upon which the management of this stage depends. When the child leaves the uterus, a very powerful stimulus to its action is removed; and this stimulus the placenta is quite inadequate to supply. The uterus, therefore, first ceases to act for a certain time, and when the action is renewed, it is weak, and continues only for a short time. If the uterus fail in discharging the placenta by a few of these efforts, it becomes accustomed, as it were, to its presence, and it no longer acts as a stimulus, but remains with the uterus imperfectly contracted around it. A

very efficient means of supplying this want of irritation to the uterus, is the pressure of the abdominal viscera which surround it. When the abdominal muscles are strong, they contract upon the retiring uterus, compressing the intestines, and consequently the uterus, on all sides. These weak pains, therefore, are greatly assisted and rendered effectual by the straining efforts of the patient acting as a stimulus to the uterus from without. But the abdominal muscles are not always strong; on the contrary, in most instances, they are extremely weak, in consequence of our civilized habits. They are too often reduced almost to a state of atony from the constant pressure of the corset; hence it follows that the uterus derives little or no support from them, and the placenta is retained, not from any want of power in the uterus to expel it, but from a want of sufficient stimulus to cause the uterus to contract. There is no *inertia* of the uterus, but only a suspension of its action. It is for this reason, and to supply this deficiency, that the pressure of the hand on the fundus of the uterus during the expulsion of the child, is found so useful; and on the same principle, as we shall have again to explain to you, the application of a bandage round the abdomen is always necessary, in order to give it proper support. This *suspended action* of the uterus, as a cause of retained placenta, must be carefully distinguished from *true uterine inertia*, which we shall allude to under the subject of hæmorrhage.

The placenta may also be *retained from irregular contraction* of the uterus, either during the expulsion of the child or subsequently. One of the fundal muscles may contract and not the other; or the fibres of the body may draw the uterus into a cylindrical shape, leaving the fundus relaxed; or lastly, there may be a spasmodic contraction of the fibres at the cervix, forming a kind of stricture.

These irregular contractions may arise, either from the too rapid delivery of the child not allowing the uterus time to follow its usual order of contraction, or from that order being inverted, in consequence of too great relaxation of the fundus, the result of deficient irritation. The effect may be, that the cervix or body contracts first, and therefore retains the placenta. Sometimes an irregular contraction of the fundus may exist and not be perceived, especially if frictions be used over the abdomen for the purpose of exciting uterine action. The anterior wall of the

uterus remains firm and contracted, and the fundus seems to be so too; but if the hand be passed down towards the sacrum, and along the posterior wall of the uterus (where the placenta is often situated), it will generally be found relaxed, and, when excited to contract, often expels the placenta, which had been perhaps for a long time retained. All these irregular contractions must be corrected by endeavouring to restore the order of uterine contraction from the fundus to the os uteri. The stricture at the cervix is, however, an exception; it must be overcome, in the same manner as strictures in other places, by direct dilatation. Another cause of retained placenta is adhesion, either partial or general; but this, like uterine inertia, is so constantly accompanied with hæmorrhage, that we must defer its consideration to that part of our subject.

When none of these causes operate, and the placenta is expelled in the usual time, the uterus does not altogether cease its contraction for some time afterwards. If left altogether to itself, without being properly supported, there is a danger that there may be too great relaxation of the uterus; consequently, hæmorrhage and severe after-pains, from the attempts made by the uterus to discharge the coagula formed in its cavity. This is one of the most frequent causes of after-pains. You have abundant evidence that it arises from the circumstance stated, because these after-pains occur far more frequently with women who have had many children, than with those who are only pregnant for the first time. In the latter case, the abdominal parietes being only once distended, retain a certain degree of tonic contractile power, which is altogether destroyed by frequent pregnancies.

If we have succeeded in placing before you in a clear manner the phenomena which take place in parturition, the contrivance employed by nature for the safe delivery of the child, and the principle which she seems to follow for the accomplishment of her purpose, we shall be enabled to enter upon the consideration of the management of labour, and you will have but little difficulty in determining the value of those rules which are laid down for your guidance. *The management of labours*, therefore, shall form the next subject for our attention.

LECTURE V.

MANAGEMENT OF NATURAL LABOUR.

Premonitory Symptoms of Labour—Evidence of the Commencement of the first Stage—Character of the grinding Pains. *Obstetric Duties of the Practitioner*—When summoned to attend—Caution respecting his Patient—Objects of the Vaginal Examination in the first Stage, before the Membranes are ruptured—Signs of the first Stage being completed—Mode of preparing the Bed, and supporting the Patient.

HITHERTO we have considered the process of parturition as an interesting subject of observation, one of a thousand illustrations of the perfect adaptation of means to the end proposed, by which nature accomplishes her purpose. We have now to enter upon the more practical inquiry; the symptoms that attend the phenomena we have described to you, and the treatment required to secure the safety of the patient. We would therefore direct your attention to the symptoms of labour, and to what are popularly called “the duties of the obstetrician.”

During the period that the uterus is descending towards the pelvis, the size of the abdomen diminishes, and the patient feels much less inconvenience and distress than she did previously. Her respiration is less impeded, and she has less anxiety. In some instances, the woman seems almost to forget what has been the constant object of her thoughts for many previous months. As the time of labour approaches, some monitors present themselves to remind her of the event; sometimes the nerves are very much pressed upon, so as to cause sensations of numbness or tingling, down one or other of the limbs; both are seldom affected. Occasionally they are slightly paralyzed, and cause lameness. As the uterus begins to press more on the pelvis, the patient becomes awkward in her carriage, and unable to walk even a short distance without fatigue. The neighbouring organs soon show the influence of the change going forward in the

uterus; the bladder becomes irritable, so that constant micturition is the result. We may, however, find it in the opposite state, and the urine retained: the rectum is also affected in a similar manner, and a fæculent diarrhœa frequently precedes labour. This is always salutary, because it secures the important object of having the intestines unloaded; so also constipation sometimes takes place, especially in hysterical habits. The length of time that the rectum may remain without an evacuation, and the woman feel no inconvenience from it, is often far beyond what you might suppose; a fortnight, and even three weeks, is not unusual. In these instances, the patient is sometimes deceived, and imagines that her bowels are quite regular, because there is a daily inclination to go to stool; there is then a scanty discharge of loose fæculent matter, leaving the mass of scybalæ undisturbed. You cannot, therefore, pay too much attention to this point of practice, because when the rectum is in this state, the whole of the large intestines become distended with flatus, which causes spasmodic pains that are often mistaken for labour-pains. Should the patient escape this, and labour actually commence, the action of the uterus goes on imperfectly: the same influence which is exerted by the uterus on the rectum in suspending its ordinary action is, as it were, reflected on itself, producing a similar suspension of the uterine contractions. For some time previous to labour, the mucous secretion from the vagina begins to increase in quantity, and often amounts to an abundant viscid discharge. This change may appear even for three weeks before labour, and consequently is occasionally mistaken by the young and inexperienced mother for the well-known symptom of labour called "the show."

When this latter evidence presents itself, the discharge is tinged with more or less blood, arising from the ruptured vessels of the os uteri when it first dilates. As soon as this takes place, labour properly begins; *the patient has now entered upon the first stage.* The pains which accompany the first contraction of the uterus are slight, short in their duration, and return at long intervals. The French describe them by the expressive term, "*les mouchettes.*" They generally also (like the mosquitos) attack the patient at night, and are sufficient to prevent her obtaining her usual sleep, although not so severe as to call for assistance.

Towards morning, however, they increase in frequency and severity ; all inclination to sleep is dissipated ; the woman is anxious to rise in order to change her posture, and will move from place to place, and try every alteration of position that may seem to allay her suffering.

Unless the patient has more than usual fortitude, and has great command of her feelings, she cannot help giving loud and almost agonizing expression to the pain she endures. She generally avoids to take a deep inspiration, or to use any straining effort. A short inspiration is followed by a shrill cry, which she cannot suppress. When you are accustomed to obstetric practice, you will learn to distinguish these *grinding pains* (as they are popularly called) from the *bearing pains* that follow them, by the peculiar cry that accompanies each. During the first stage, the irritability of the bladder and rectum generally continues, so that the patient has a frequent desire to evacuate them. A slight rigour may be observed at the commencement, or she may be seized with sudden vomiting. The latter is often very serviceable in those who are of a plethoric habit, because the nausea and sense of exhaustion that follow are often a means of controlling excitement, either in the mind or in the circulation. When any of these evidences prove that labour has commenced, the practitioner is generally hastily sent for. Sometimes, however, it is avoided until the pains seem to say, "He can no longer be dispensed with." If labour be preceded by false pains, he is often summoned unnecessarily, these spasms being mistaken for labour. The medical attendant, knowing that such a mistake is possible, and that even if labour have actually commenced, it will occupy some time before he can be of use, might procrastinate : he might be disposed to think that he has quite sufficient time before him, and give but a dilatory attention to the message he receives. *A greater mistake cannot be committed.* Any summons from one who has placed entirely in your hands her own safety and that of her offspring, equally dear to her, should receive instant attention ; neither are you to consider the urgency as measured by your own, but by her impressions of the case. You may be called upon without there being the least necessity for haste ; but by your promptitude you will gain a considerable advantage in strengthening the confidence she has already given you. The

necessity, however, may be real, and not fictitious. Cases have occurred in which the patient has been delivered before the practitioner arrived; and if such an accident arose from any negligence on his part, he must have a much greater influence over his patient than is usual, if he can recover from the effects of it.

If it be a first pregnancy that you are summoned to, it is advisable that your introduction be not too abrupt: caution in this respect is of still greater importance if called to a patient to whom you had not been previously introduced; the mere circumstance of a stranger entering the apartment of a parturient woman has caused a total suspension of her labour. Some preparation, by way of announcement, is therefore necessary. For the same reason, it would be prudent, when introduced, to direct your patient's attention as much as possible from contemplating the character in which you appear before her, to draw her away from the subject that brought you there, and to lead her to forget the office that you have to fulfil. You would not, therefore, catechise her too strictly about herself, or remind her of what is going to happen by too busy a display of preparation. A few minutes' conversation with the nurse is generally sufficient to learn every particular of importance; but your patient should only receive from you the words of comfort and encouragement. The nurse, however, does not require the same forbearance. It will be your duty to ascertain from her every point upon which you desire to be satisfied. When the pains commenced? Their character? If accompanied by much, or by little excitement? The state of the bowels, and whether the bladder has been relieved? If your patient has any constitutional peculiarity? You should also examine the bandage, pins, ligature, and every trifling matter which might inconvenience you, if not prepared according to your views.

Having satisfied yourself, you can then return and engage your patient, if possible, in general conversation. You may thus form your own opinion of the character of her labour. If it be in the commencement, when the pains are short and the interval between them long, you can engage her attention with facility; but if the pains are severe, these attempts at conversation had better be dispensed with. The patient should be left in charge of the nurse, because your presence may become un-

pleasant to her as a witness of her suffering, and would be embarrassing, if it interfered with the necessary evacuations: you may therefore retire to the neighbouring apartment, until you have determined upon the time for taking the first important step in your professional capacity.

Making a Vaginal Examination.—Some have considered it as a matter of the first importance, that this should be done as early as possible in the labour, in order that any correction which might be required in the presentation should be effected before it was too late; as, for instance, the funis or hand coming down with the head, the head descending in a wrong position, or perhaps the shoulder presenting. In order to interfere with any of these complications, the labour must be more than in its commencement; the os uteri must be dilated, although the dilatation may be far from being complete. It would be precipitate, therefore, to require an examination before the steady frequency of the pains gives evidence that this was likely to have taken place; and therefore a vaginal examination very early in the labour, when the pains only return slowly, would be unnecessary. Neither is the object for which such an examination is stated to be made, of that importance which seems to be attached to it. The funis cannot be interfered with, or suffer any very dangerous pressure, so long as the membranes are entire. We shall presently point out to you the danger of meddling too much with supposed wrong positions of the head, as well as when the hand comes down with it. The only position that should be ascertained before the waters are discharged, is the shoulder presentation; but if you examine for this too soon, when the os uteri is only slightly dilated, and the presenting part above the brim, you might fail to detect any presentation, and even if you did, may, after all, be mistaken in the result; the hand and arm might be felt, and yet the head afterwards descend. It is certainly very satisfactory to find out that the head presents, as early as you can; but if you cannot do so in the beginning of labour, it is no proof that the position is preternatural. If you cannot, therefore, gain a decided advantage by making a very early examination, there is one strong reason for a little delay. When the patient is only enduring the first short pains that attend this stage, she has always a great reluctance to be examined; all her natural feelings and

prejudices are in full play against you : she submits very unwillingly, and complains loudly of the least pain or inconvenience you may cause her. You will not, therefore, find it easy to make a satisfactory examination ; and if you fail, you may not so readily obtain her consent to its repetition. When there is *no absolute necessity*, therefore, it is better to wait until these grinding pains increase in strength and frequency ; her mind is then absorbed in her present suffering ; she is willing to submit to anything which may be of use to her, and is often very anxious to know whether the labour will be safe : you have also the great advantage of being able to make the examination *perfectly*, because the dilatation of the uterus has made some advance. Having determined upon the proper time for making a vaginal examination, the nurse should communicate your wishes, which are now readily acceded to : she may then place her in the most favourable position for the purpose. The patient, loosely attired in her night-dress, should lie on the bed on her left side, as near to the edge as possible, having the knees drawn up towards the abdomen. You should then wait until the pain returns ; and when it is about to cease, pass the fore-finger of the right hand, anointed with cold cream, bear's grease, or any unctuous substance, within the vagina ; examine carefully its condition, if it be relaxed and moistened with the usual secretion, or if dry and rigid. *Examine the rectum through the posterior wall*, which, if empty, or nearly so, feels like a thick band ; but if loaded, it projects into the vagina a surface so firm and irregular, that if it were your first case you might imagine it was the distorted sacrum or perhaps the back of the child. You should also *examine the anterior wall* of the vagina : if the fundus of the bladder should press into it ; advancing slowly, *the middle finger* might then be introduced, in order that *a careful examination of the os uteri* be made. One finger is not sufficient for this purpose, unless the uterus is low in the vagina. If it be above the brim of an ordinary sized pelvis, the fore-finger, or rather the nail of the fore-finger, just touches the anterior part of the mouth of the womb, and may also touch the head if it lie near to it, but scarcely any other information is gained ; knowledge quite sufficient for those who only examine to find if the position be natural, but not at all adequate for the practitioner who is anxious to form a correct judgment of the

time this stage might occupy, or of the causes which might retard its progress. The middle finger is longer, and reaches higher in the vagina: with two fingers, also, you have a double advantage, from your sense of touch. When you have acquired, from long experience, a perfect tact, then one finger may be sufficient. Both fingers being introduced, pass them first along the sacral side of the vagina, and when you cannot advance them further, direct them forwards towards the pubis. If the pains have ceased and the os uteri be dilatable, having the membranes lying loosely within it, you feel what seems to be the irregular folds of a flaccid bag projecting into the vagina; examining this with caution, the edge of the os uteri, soft but more resisting, may be traced, and if the finger be passed within it, you will sometimes feel, towards the pubis, the head firm and round; you may, however, often fail, and yet the head present.

The finger should not be withdrawn until you have ascertained the state of the os uteri, as to its dilatability and density, its exact direction, as nearly as you can judge, and the degree to which it is opened. The act of withdrawing the fingers sometimes induces a pain; you may then cautiously observe what takes place,—the fulness and slight closing of the os uteri, the membranes descending as a tense bag, and the mouth of the womb tightly stretched over it: but take care of roughness, lest the membranes give way. Before the fingers are removed from the neighbourhood of the os uteri, examine the distance of the sacrum; and as they are being withdrawn, ascertain, if possible, the space in the pelvic cavity. It would be well, also, in the commencement of your practice, when it is so necessary to educate your sense of touch, to take this opportunity of examining the orifice of the urethra. For this purpose, when the fingers are being withdrawn, let the forefinger follow the course of the canal to its termination in the trumpet-shaped, semi-cartilaginous opening. Be careful, at the same time, to avoid the clitoris. In order to make a sufficiently careful examination, a little time may be required, during which the pains may return; you should then cease until they subside, noting only those points which have been mentioned. But make it a rule not to withdraw from your examination until you have perfectly satisfied yourself as to the character of the labour. Having accomplished

this, a second examination during this stage, unless it be prolonged, would be unnecessary.

Exploring the vagina and uterus in this way, is called, in the popular language of midwifery, "trying a pain," or "taking a pain." And the object assigned to it by the patient and her friends, is to learn whether the labour is safe, and how long it may last. You are, therefore, generally asked, and expected to give a distinct answer to both these questions. The first is usually easy, and may be replied to in the affirmative; but if you have any doubt, you should not precipitately sound an alarm. If you ascertain that it is a cross-birth, as shoulder-positions are called, you may state the nature of the case, and the kind of assistance required, *to the friends* of your patient, in order that they may select a consultant in case you find it necessary; but be cautious not to alarm the patient herself: it is quite time enough to inform her of her situation when you have determined to deliver her. In all other cases, the difficulty must be decided by time and further observation of the case; you should not, therefore, give way to apprehensions which may be unnecessary, or commit yourself to an opinion which the result might not justify.

The second question, as to the duration of labour, cannot be too cautiously answered. The friends may ask, "Is everything right?"—but, "How soon shall I be well?" is invariably your patient's anxious question. In order to tranquillize and encourage her, it might seem pardonable to state a period for the termination of her sufferings earlier than what you know will be the case. But such a practice would be extremely injudicious, because when the time had passed in which she expected a relief to her anguish, her disappointment would lead to impatience of further suffering, if not to a secret dread that the delay arose from some cause dangerous to herself. The result might be a suspension of the uterine action, and a still further prolongation of her labour. Along with this, being proved, as it were, a false prophet, your patient may lose all confidence in your opinion and judgment.

It is better, therefore, to err on the other side; and if you are obliged to give a distinct and positive answer, to state a period beyond that in which you expect the delivery will take place. No doubt this will give but little comfort to one who requires

every consolation, but it will induce her to summon that resolution and patience under suffering, which is the peculiar attribute of her sex; and it will enable her to go through her trial more favourably, because more patiently, than if she expected more prompt relief. Her confidence, also, in your judgment, would be rather increased if your prediction prove to be true; and if not, she would much more willingly forgive a deception which rendered her delivery an agreeable surprise. If, from any accident, the membranes are broken during this stage, a second examination is necessary, in order to determine the presentation, and to ascertain any accidental complication which might occur. If, when the liquor amnii escapes, the dilatation be slightly advanced, and the orifice of the uterus increased only an inch or two in diameter, you may expect more or less delay in the completion of this stage, unless the cervix of the uterus be extremely thin. If it be at all thick, the irritation of the head generally renders it rigid, no matter how dilatable it may have been previously. In this second exploration, therefore, a very cautious opinion should be given as to the duration of the labour. The head, if presenting, may also be felt; but the position cannot be determined until the dilatation is more increased, and the head fully in the brim. The small portion of the head which occupies the slightly dilated os uteri is not sufficient to determine anything about it; and if you pass the finger within the os uteri for the purpose of tracing the sutures and fontanelle, you will only succeed in exciting a great deal of unnecessary irritation in its tissue; neither can you define the characters of the position, through the os uteri, with sufficient accuracy to place any dependence on the evidence they give.

The time has not yet come to decide the position of the head. If this be true, then it is scarcely necessary to point out to you the absurdity of attempting to correct its supposed wrong position at such an early period of the labour. Yet this is one of the objects for which we are instructed that a vaginal examination is necessary, and are even recommended to rupture the membranes for the purpose of altering the position of the head.* The hand

* "If it be discovered early, it is certainly proper to rupture the membranes, and turn the vertex round, which is easily accomplished." Burns' Midwif. p. 394.

may be found presenting with the head under similar circumstances, and the correction of this accident is also assigned as a reason for examining *per vaginam* as soon as possible. We very much question this necessity, or even the propriety of interfering with the presentation so soon. You will recollect, that in order to correct this malposition, the hand and arm must be pressed up above the head, and there maintained until the succeeding pain drives down the head below it. If the os uteri be only slightly dilated, and the head not completely in the brim, you will find it very difficult to accomplish this, and in the attempt you expose yourself to the double risk—1st, of increasing the irritation of the cervix uteri to a much greater degree than might otherwise take place; 2ndly, in your manipulation, you cannot press up the hand without also pressing against the head, which we have assumed has not yet passed into the brim of the pelvis. The effect of this might be, that you may just succeed in pushing it more on the brim than it had been; and the necessary consequence is, that the head not being able to pass into the pelvis, the action of the uterus is directed more upon the shoulder. The side of the face, ear, and neck, might first be directed towards the pelvic cavity, and then glide with the vertex towards the iliac fossa, while the shoulder would occupy the brim. So that, after an attempt of this kind, you would find the arm coming down into the vagina rather more than before, and after a few pains the position completely changed. There cannot be a more awkward accident than to convert a natural into preternatural labour. Therefore we would question altogether the propriety of meddling with the position thus early, because it seems to us far more probable that a little awkward manipulation would produce this derangement, than that it would take place if the position were not interfered with. In fact, nature would be less likely to blunder than you would.

What we have stated to you does not apply to the same position when the os uteri is more dilated, and the head quite within the cavity of the pelvis; we would desire, at present, only to point out to you our reasons for objecting to submit your patient to a very early examination, with a view of unnecessarily endeavouring to make corrections. The only accident which would seem to render an early examination necessary is when the funis is prolapsed. We have stated that so long as the membranes

are entire, and the liquor amnii surrounds the funis, there is little danger that the circulation will be arrested. But if the waters prematurely escape, this alone renders an examination necessary; and therefore the funis may be detected in sufficient time to determine upon the course which it may be requisite to pursue. In ordinary cases of labour, such as we are now speaking of, the membranes seldom give way until the second stage has made some progress, and often remain entire until the head of the child is almost expelled. You seldom, therefore, have occasion to make more than one examination during the first stage; and having obtained by this every information you require, it is better then to commit your patient to the nurse's charge, and to retire to the next apartment until you are again summoned by the urgency of her pains, or by some new symptom presenting itself.

The time which the first stage may occupy, when there is no cause to render it tedious, is very uncertain. It is, of course, much longer with women pregnant for the first time than with those who have had many children. It may last from twelve to sixteen hours; but where this stage is prolonged, the following stages are generally short, and sometimes bear an inverse proportion to each other. You cannot take time as a criterion to judge when the first stage is about to cease and the second to commence; but you may do so by closely observing the pains. We have described to you the character of the grinding pains that accompany the first stage: they frequently merge into the bearing pains so gradually as to require some attention to observe the change. You may remark, when the pain comes on, that the patient is obliged to grasp firmly whatever is within her reach; she retains her breath more than before, and sometimes makes an involuntary effort to force the child down. Her voice also alters, its tone is more subdued, and she seems more patient of her suffering than before. Sometimes the complete dilatation of the uterus is marked by constitutional symptoms; there may be a slight rigor or vomiting, perhaps a strong inclination to go to stool. Whenever any change of this kind is noticed, you have just reason to suspect the commencement of the second stage, and as you cannot tell with what rapidity it may proceed, the patient should no longer be suffered to go about the room. She must now remain in bed, which should

be properly prepared for her reception. The manner of doing so is thus:—A skin of morocco leather, or a broad piece of Indian-rubber cloth, is usually placed next the bed, to protect it from being stained; and a blanket, folded very wide, enclosed in a soiled sheet, is placed underneath the hip of the patient as she lies upon the left side. They should be so fastened together that the whole may be removed at once without difficulty. By this means the discharges are absorbed, and prevented from soiling the sheets of the bed, while these foul clothes may be withdrawn after the delivery of the patient without disturbing her. Trifling as these details may appear to you, we must direct your attention particularly to them, because of a practice which is prevalent among the lower classes here, that seems both inconvenient and dangerous. The parturient woman generally sits at the end of the bedstead without making any change in her dress. The bed and bed-clothes are doubled up in such a manner that she can recline against them something like a sofa, and so she remains during the whole of the second stage, until she is delivered. The intention of this arrangement is, that when labor is quite concluded, the clean bed and bed-clothes may be laid down, the patient's day apparel changed for her night-dress, and everything made "comfortable," as the midwives express it. All this may be, and often is, done without injury to the patient; but recollect the risk that is run by moving the patient so much, at a time when she should be kept perfectly quiet; and if an accident should happen, if hæmorrhage should result from it, imagine your perplexity. Her dress is in your way when you wish to command the uterus. You may not have time to get the bed laid down again, so as to place her in a horizontal position; even moving her for the purpose is dangerous. You are surrounded with a host of unexpected difficulties, and her life might even be the sacrifice of a little want of forethought. You should, therefore, always take care that your patient is loosely attired in her night-dress, and placed on her left side in the bed, guarded in the manner described, when there is any evidence that the first stage is near its completion.

When the os uteri is quite dilated, and the bearing pains become more decided in their character, the patient is anxious to have her body fixed as much as possible during their continuance; she therefore requires to have something within her

reach that she may grasp firmly. Her feet also should be supported while the pain lasts. For this purpose a sheet is generally fastened to the bed-post in such a manner that she can easily hold by it, while the nurse may support her feet by pressing a pillow against them. It is a very common practice to place between the knees a small pillow when the head of the child is beginning to press upon the perinæum, but you will find it both useless and inconvenient. The patient derives very little advantage from it; and it is constantly in the way when you wish to make a vaginal examination, or when the head is passing the vulva. On this point we may be permitted to anticipate the consideration of the second stage towards its conclusion, and observe that when the labor has so far advanced as to render the aid necessary which this is intended to afford, much more efficient assistance may be given by the nurse, if she raise the knee moderately with one hand and support the foot with the other.

When the second stage of labour has commenced, the patient experiences much more fatigue than she did before; as it continues, she feels exhausted by the struggle, becomes heated and thirsty, and often grows dispirited. Hence has arisen the pernicious practice of giving wines, spirits, and other such stimulants, "to help her pains." You must imperatively *forbid heating drinks of all kinds* to be given to the patient. Such a practice would not only increase the sense of exhaustion when the stimulus had passed away, but it might also contribute to increase greatly any tendency to inflammation which might exist in the passages, while the head is passing through the pelvis. Barley water, tea, very thin gruel, or such like drinks only, should be given. *A free ventilation* of the apartment should also be secured; and at the same time care is necessary that your patient be not exposed to any draughts of cold air. For this purpose you must not fall into an opposite extreme, and, as is sometimes done, have the bed-curtains closely drawn to prevent all access of the air. In this manner the parturient woman has been kept in a vapour bath of impure air; an excellent excitant for miliary fever. On the contrary, it is better to have no curtains, or at least to have them only drawn sufficiently to prevent exposure to a draught of air. It is also very essential to have *as few persons in the room as possible*. You

should only permit one person to remain with the nurse, *the nearest married relative* of your patient, with whom she would feel no restraint, and to whom she can communicate freely all her feelings. The officious assistance of friends should be sedulously shunned; it is often extremely disagreeable to the woman herself to have these witnesses of her suffering present; and although she may silently tolerate the inconvenience because she is conscious of the kind motive which actuates them, still it is embarrassing and unpleasant. To the practitioner it is still more inconvenient, because they are often unaccustomed to such scenes, and become alarmed at the agony their friend seems to suffer: their faces express even more distress than the patient's, and like multiplying mirrors, reflect every pang she suffers with tenfold power. If this stage should continue longer than usual, their anxiety is proportionally increased. Anxiety is soon succeeded by a secret if not open doubt of the practitioner's competency, and their ominous expressions of distrust very rapidly produce a corresponding effect upon your patient. She soon becomes dispirited, and fearful for the result; so that at a period when it is most important that the action of the uterus should continue regularly and efficiently, it may be altogether suspended. When years have given you not only experience but station in your profession, you may not be subjected to such misgivings; but be assured, that in the opening of your career, when you are as yet but little known, and have to build up your reputation, you will be subject to these inconveniences, if you are not decided in preventing them.

LECTURE VI.

MANAGEMENT OF NATURAL LABOUR, CONTINUED.

Second Stage—Obstetric Duties of the Practitioner—Vaginal Examination—Its Objects—Support of the Perinæum—Its Intention and Mode of Accomplishment. *Third Stage*—Removal of the Placenta—Support of the Uterus—The Abdominal Bandage—Its Object and Mode of Application—Management of retained Placenta without Hæmorrhage.

IN the preceding lecture, we considered the duties you had to fulfil towards your patient when the os uteri was in the process of its dilatation, and when THE SECOND STAGE was about to commence. We explained to you the general arrangements, both as to the bed, the apartment, and the diet required by your patient during this trying stage. In continuation we would observe that from the moment of its commencement, she must receive your most sedulous attention; she cannot now be left to the nurse; you must sit beside her, attentively observing the progress of the labour, and be prepared to act the moment your assistance is required. You have to witness a struggle of greater or less severity, in the effort of the uterus to force the head through the pelvic cavity. The bearing pains return with a regularity and strength that would seem sufficient to overcome any ordinary obstacle with rapidity, and yet the advance of the head is often slow, and its progress bears no proportion to the effort which seems to be employed. It is your duty to watch carefully the effect produced by the pains, and to sustain your patient through this severe trial with every encouragement. The proportion between the head and the pelvis must be accurately observed, the exact position of the head ascertained, and the progress which it makes through the pelvis carefully noted. More than one vaginal examination is, therefore, necessary; but you must bear in mind, that the passages are now expe-

riencing unusual pressure; there may be, probably, some congestion and increase of temperature, and this might be much increased if the vagina were irritated by frequent examinations. In order to obviate any injurious effects, nature provides a resource in the increased discharge of slimy mucus which now flows abundantly from the vagina. But if examinations be repeated too often, and the passages become irritated, this discharge is diminished; it may be arrested, and the parts become hot and dry, or perhaps it may be succeeded by a thin serous discharge, that rather increases the irritation. This change, therefore, in the character of the discharge, serves as a useful indication that caution is required in this respect. It had not been lost sight of by the older practitioners, who supposed that the frequent introduction of the fingers into the vagina dried up the parts by absorbing the discharge.

The *first object*, then, of a vaginal examination in this stage, *is to determine the proportion between the head and the pelvis.* For this purpose the fingers should be passed carefully between both in the interval of the pains, directing them, in the first instance, between the pubis and head, and moving them round on either side. The ear can be felt if there be sufficient space for the head to pass, but if the head be high up in the pelvis, the finger can only just touch it. If the ear cannot be reached readily, and there seems to be a want of proportion between the head and the pelvis, you have still another means of testing its degree, by examining the presenting part of the head. When it is only slightly compressed, the scalp is simply folded or puckered by the closing of the sutures; as the compression increases, these folds merge gradually into one, which ultimately forms a distinct tumour. This continues to enlarge, so that in cases of impaction of the head, it is sometimes of great magnitude. The manner in which this change takes place, and its degree, is generally a sufficient proof of the amount of the disproportion. If the tumour form very slowly, and never increase to any great size, you may infer that the head will pass safely through the pelvis; but if, on the contrary, it increase rapidly, and attain a great size, the indication must be unfavourable.

The *second object* of a vaginal examination is, to ascertain the exact position of the head. We have already pointed out to you the means of distinguishing the different positions from each

other. We shall, therefore, at present only bring before your notice those positions which we are directed by some authors to alter as soon as they are found out, in order to prevent the head becoming impacted in the pelvis afterwards.

One of these cases is when the head enters the brim in the left fronto-cotyloid position. It is assumed that this cannot pass safely, but will cause great delay and difficulty in the labour: therefore, it is laid down that the correction must be made the moment the position is ascertained. We have already stated to you the experience of Naegele, confirmed by other observers, that nature, if left to herself, will correct this deviation, by rotating the head into the right occipito-cotyloid position. The probability is, therefore, that by meddling too soon you may prevent this, and prematurely force the head into a more unfavourable position than it had been. The moment this position is detected is not, therefore, the time for interference; it is more advisable to wait and observe the course the head will pursue. It may correct itself; it may advance and be delivered in the third position without injury; it may be arrested. The last is the only condition which would justify your aid. The head may then be displaced from its situation, and pressed back in the interval of the pains, and a *very slight rotation* is generally sufficient to make it glide easily in its proper direction when the pains return. The very same observation applies to those instances where the head and hand, or even arm, descend together. This accident is often the result of the pelvis being too wide, and if so, both will be expelled without difficulty; but sometimes the arm comes down a little too much and prevents the head advancing, or the head may be arrested by the hand descending with it. In either case, the hand or arm can be very easily pushed back when the pain ceases, and so maintained until the next pain advances the head, which generally passes down very rapidly as soon as the correction has been made. When the head is in the cavity of the pelvis, there is not the same danger of displacing it as when it is only entering the brim; and consequently, our previous observations on this accident (pp. 98-99) do not apply to the present case. You should not, therefore, when these deviations occur, too hastily assume that the head cannot be delivered. It is more advisable to wait until they become causes of delay.

The *third object* of a vaginal examination is, to note the pro-

gress which the head makes. In natural labour, where no difficulty presents itself, a very few examinations, at proper intervals, will be sufficient for the purpose, because its advance is generally quite obvious; but in difficult labours, where the head makes a very slow progress, and there are other causes of embarrassment present, more care is required; their consideration, however, is beside our immediate subject. Having ascertained the position of the head, and its relation to the pelvis, the next object of attention is its descent upon the perinæum. You must, therefore, *be prepared to give the perinæum support* the moment it suffers any degree of distension. The mode of doing so, which I have been in the habit of adopting, is somewhat different from that directed by the more popular writers on midwifery. Ramsbotham and Rigby both employ the left hand to press against the perinæum, and the right is kept in reserve to make any necessary correction. Churchill and others adopt the same plan. Dr. Rigby admits that "it is awkward at first, because it requires the hand to be considerably twisted, and makes the wrist ache a good deal."*

The left hand is twisted, in order that its fingers may be directed forwards toward the perinæum. Dr. Ramsbotham directs the palm of the left hand to be pressed against the perinæum with the fingers; I presume, the other way. He does not state so; but I draw the inference from his description: and if correct, it is liable to the objection of which Dr. Rigby is fully aware, and endeavours to avoid—that is, that the part of the hand which has the least acute sense of touch is applied to the margin of the perinæum, and you cannot so readily perceive its degree of distension as when the fingers lie close to it. The plan which I have found the most useful and convenient to adopt at this period of labour, is the following:—To sit behind the patient as she lies upon her left side, the back of the chair being towards the head of the bed, and while the head of the child is passing through the pelvic cavity, to press moderately with the left hand over the hip of the patient. Counter-pressure in this way employed is generally grateful to her, and seems to give her some relief; it assists also in keeping the pelvis fixed when the head is passing the perinæum, the most important part of this process. Having the

* Rigby, p. 112.

left hand so employed, the right can be used to support the perinæum, a single fold of a fine napkin should be placed along the edge of the perinæum, and the right hand so applied that the fold of skin between the fore-finger and thumb should correspond to this, the fore-finger and thumb passing on either side of the vulva, and the palm of the hand, resting against a thicker fold of the napkin, applied to the posterior part of the perinæum. By this means you have full power to make any counter-pressure with the palm of the hand which may be necessary, and the fingers being quite close to the edge of the perinæum and vulva, you can easily trace the margin of the perinæum, and feel the head, if necessary. Thus one hand fulfils the office generally assigned to two, and enables you to grasp with the left hand the pelvis, to prevent the patient moving away too suddenly when severer pains come on. If, the head being expelled, this be no longer necessary, you can employ the same hand to support the uterus during its contraction in expelling the body of the child. Besides these advantages, it is certainly less fatiguing. The only inconvenience of this method is, that when the funis is coiled round the neck of the child, so as to make it necessary to remove it, or that the delivery of the shoulders should be assisted, the hands must be changed, that the left may support the perinæum and the right make the required correction. But this is a temporary disadvantage and only arises occasionally. We shall suppose you, therefore, thus prepared to give the perinæum the required support; the only question is, when your assistance is needed. The young practitioner, fully impressed with the importance of preventing laceration, hardly ever commits the mistake of being too late in attending to this point. He very generally errs on the other side: he presses against the perinæum a great deal too soon, and causes unnecessary heat and irritation in consequence, which rather retards its distension. His mistake arises from supposing the perinæum in danger the moment the head touches it. We have explained to you that the head alternately touches and retreats from the perinæum, often for a long time before the perinæum suffers any dangerous distension. You must not, therefore, be too precipitate; it is better to wait until you feel the head protruding, with each pain, through the vulva, because at this time it is getting gradually upon the ischio-pubic ramus, against which it rests, while the anterior

part of the head presses, with considerable force, against the perinæum. Caution is also necessary as to the manner in which the perinæum is supported. The object in view is to obviate the effects of too violent distension. The pains at this time are very unequal, sometimes weak and again very strong; you support the perinæum against the latter by moderate counter-pressure, to prevent accidents; but against the former no such precaution is necessary: you must not, therefore, press with every pain indifferently, but only when the uterus is acting with great force. Again: when the head is nearly protruded through the vulva, anxiety to save the perinæum may be the cause of its rupture. For instance, if you attempt to draw the perinæum back over the head, it will be stretched too suddenly over the bi-parietal measurement, the widest part of the head. If, on the other hand, you push the head too much forwards, pressing with the pains from the sacrum towards the pubis, the same effect will be produced in a different manner; you force the parietal portion of the head too rapidly through the vulva. At this point it is better to continue the same moderate counter-pressure, to make no attempt to hasten the delivery, and to allow the head to pass along the hollow of the hand, in the same manner as it moved along the curve of the sacrum. When the head is passing out of the vulva, you should then direct it forwards toward the pubis; and when it is delivered, examine carefully lest *the funis may be coiled round the neck*. If such be the case, and that it is only a single coil, it will generally be sufficient to draw down a little more of the funis, and loosen it. A single coil seldom retards the delivery of the child, or arrests the fœtal circulation; but two, and even three, coils are sometimes met with, and the child placed in great danger of strangulation. In these cases, as much of the funis as possible should be brought down, and the coils so loosened that one may be drawn over the head. There are cases where this cannot be done, and the only resource left is to tie and divide the funis, and extract the child as soon as possible, in order that respiration may be established. This operation is hazardous to the child's life, and can only be viewed as the lesser of two evils.

If the funis be not found about the neck, the perinæum must still be supported until the next pain, usually a tardy one, expels the shoulders. The same caution must be exercised as before,

lest the arm or hand should lacerate the perinæum as it is coming out of the vulva. This should be particularly attended to in second positions of the head.

Sometimes the shoulders are very wide, and require to be assisted, which may be done by placing the fore-finger of the right hand within the axilla of the child's arm, on the pubic side, and guarding the perinæum carefully with the left hand. As soon as the shoulders and thorax of the child are delivered, it can respire, and is, so far, beyond danger; no haste should, therefore, be used in extracting the body and lower limbs; it is preferable to allow the uterus gradually to expel them, and, while it is doing so, the left hand should be immediately applied over the fundus, in order to maintain a moderate pressure upon the uterus while it is descending towards the pelvis. This should never be neglected, because it insures a uniform contraction of the uterus, and often the expulsion of the placenta into the vagina. When the child is born, such is the anxiety to remove it as soon as possible from the mother, that the tying of the funis is the immediate occupation of the attendant, while the uterus is generally left to itself. The motives assigned by friends for this haste, is their fear lest some accident may happen to the child: it may get cold, etc. Just as often the real cause is a little natural desire to see and exhibit it; you should not, therefore, suffer yourself to be hurried by these solicitations, nor withdraw your hand from the uterus until you have secured it, either by a temporary bandage, applied in the manner we shall presently describe, or by the hand of the nurse, if she is sufficiently intelligent to understand your object. The latter plan is more convenient.

When the uterus is thus prevented from again relaxing, you may attend to the funis. The delay is serviceable to the child, because time is allowed for the transition from the placental to the pulmonic circulation, by which the latter is completely established, before placental life altogether ceases. This is of great importance to the health of the child afterwards. When this is perfectly effected, the circulation of the funis often ceases; but if its pulsation be felt, the funis may be tied if the child cry strongly. The manner of doing so is by applying a strong ligature of housewife thread, bobbin, or narrow tape, about two inches from the umbilicus, and a second about an

inch further. You must be careful to see the part of the funis you are dividing, lest the fingers or any part of the child should be in your way, and also in order to examine the cut surface of the umbilical portion. The blood should be squeezed out of the vessels, and the surface wiped with a napkin, for the purpose of detecting any oozing of hæmorrhage that might take place if the funis were not properly tied. The child may then be removed and the separation of the placenta attended to.

If the bandage have not been previously used, the hand may be again applied to the fundus uteri, which is generally found in a semi-contracted state. With a little attention, you will presently observe it become harder from contraction, although the patient scarcely complains of it. A very moderate pressure on the fundus at this time is often sufficient to expel the placenta completely out of the vagina; but if not, it can be drawn out by the funis quite easily, directing the funis forwards in the axis of the vagina. But if the uterus should not obey the stimulus at first, do not persevere; it is always more advisable to wait for some time than to use too much irritation. Neither should you attempt to remove the placenta by the funis alone. By great violence, it is true the funis may be broken, or the uterus inverted. I do not attribute to you such awkwardness; but by pulling frequently at the funis to ascertain if the placenta be separated, you may excite an irregular contraction of the uterus. Passing the fingers into the vagina is often sufficient to excite the action of the uterus; and drawing the placenta by the funis may excite it still more. If the uterus contract, and the order of its action be not secured by the means already pointed out to you, the great probability is that, being nearly emptied of its contents, the lower fibres will contract first, and retain the placenta. Thus, by pulling too much at the funis, the placenta may be retained. By a little caution, and by moderate pressure on the fundus of the uterus, you will generally secure its favourable separation. This being accomplished, the next and concluding object of your attention is to preserve the uterus in that state of contraction which is so necessary to prevent subsequent hæmorrhage. We have already explained to you the efficiency of the abdominal muscles, when they are strong enough to contract firmly upon the retiring uterus. But when these muscles are rendered inert

from the constant distension they are exposed to, they can give the uterus no support, and there is, consequently, a constant risk that the uterus may again relax and pour out blood, if this want be not supplied by artificial means. Hence the use of the abdominal bandage. The mode of applying it demands your attention; because it may be made useful or mischievous according to the manner it is employed, and many of the objections raised against its use have been founded upon its improper application. Sometimes it is bound so tightly over the uterus, that the patient can hardly breathe, or it may be so applied that the least motion of the patient displaces it, and it becomes twisted round the loins like a rope. All these inconveniences, distressing to the patient and useless for the intended purpose, arise from a mistaken view of the use it is meant to fulfil. The waist is to be compressed into shape, and therefore the patient is bound up so tightly that she can seldom tolerate the pain of the bandage: it is soon loosened, and perhaps altogether discarded. A bandage properly applied may be made to effect two objects; one, to support the pelvis by compressing it as much as possible; another, to support the uterus by *moderate* and *equable* pressure over the whole abdomen. The articulations of the pelvis have undergone a great degree of tension during the passage of the head, and a dull pain sometimes remains, which is much relieved by counter-pressure. The uniform pressure of the intestines is necessary to prevent relaxation of the uterus. The mode of applying the bandage for these purposes is to commence by drawing it evenly over the pelvis, its lower edge, when so placed, being about one inch below the trochanter; this margin should be drawn as tightly as the patient will bear, and pinned securely below the right trochanter. The bandage should be again drawn and pinned in a similar manner across the ilia, so that the pelvis may be embraced by this portion of the bandage, about three inches in width, as tightly as possible. Having accomplished this, the remainder of the bandage should be drawn and pinned with moderate tightness, but equally from the pelvis to the diaphragm, so that the whole of the abdomen be included within it, and not permitted to project over the bandage in the unsightly manner which may sometimes be observed.

When the bandage is properly applied, the patient always

experiences comfort from it, a sufficient evidence of its utility. There is a great variety in the material employed for bandages. Sometimes a piece of calico or a napkin is used; and, again, you will find them more complicated in their mechanism than the most fashionable corset—both are equally inefficient. Calico and diaper are too unyielding, and if pinned tightly, will hurt the patient; nor can they be employed unless they are so loose as to be useless. *The obstetric corsets*, if we might so call them, for drawing in the waist, are liable to all the objections which have been urged against bandages. It is necessary that a bandage should be elastic, that while it supports the abdomen it may yield readily to its action; it should be sufficiently thick or firm not to wrinkle easily, should be soft in its texture, and at the same time strong enough to bear being tightly drawn. A double fold of flannel would answer the purpose, and has the advantage of being easily pinned; but if you remember the intention the bandage is to fulfil, your own judgment will best direct you to the kind of material which will suit your object. From what we have stated, you will perceive that a bandage may be made useful or injurious, according to the manner in which it is applied. You should not, therefore, entrust this simple but important part of your duties to another. It is sometimes a practice to commit to the nurse its application: it would be imprudent for you to do so in the first instance, or so long as there is any risk of hæmorrhage, from relaxation of the uterus; it will be sufficient time to leave the management to her when your patient is secured from danger. If ordinary discretion is used, it may be applied without offending the feelings of the most sensitive person, and therefore no motive of false delicacy should prevent the practitioner fulfilling this essential duty. When the bandage is applied, the folded sheet, etc. that had been placed under your patient during labour should be removed, and replaced by others dry and warm, in order that she may be induced to sleep, and that she may not afterwards be disturbed. It is the more necessary to attend to this, because it too frequently has happened that hæmorrhage has been induced by imprudence on the part of the nurse, who, when the attendant has left his patient, immediately sets about making her “dry and comfortable,” and in doing so, causes so much excitement in the process of dressing her, and changing the bed-clothes, that

hæmorrhage is the result. The patient, after her delivery, always experiences a nervous shock, often very slight, but still sufficiently obvious. Although happy in her relief from suffering, and in the birth of her offspring, she still feels depressed; and this period, beyond all others, is that in which perfect repose is absolutely necessary. Too much caution cannot, therefore, be exercised to prevent her being disturbed. Having secured to your patient perfect quietness and freedom from interruption, your immediate duties are completed; but still caution is necessary, and although you should retire from the apartment, it would not be advisable to leave the house for at least an hour after her delivery, or until she falls into a sound sleep.

It sometimes happens that the placenta is retained after delivery, without any hæmorrhage taking place; and although we shall have again to direct your attention to these retentions, when accompanied by flooding, a few words may not be out of place here, in reference to these very frequent retentions, where no hæmorrhage arises. The causes generally assigned for retained placenta are, either *inertia* of the uterus, *hour-glass contraction*, or *adhesion*; but one quite as frequent, if not more so, is *suspended action* of the uterus. The former causes are generally attended by hæmorrhage, but with the latter it is very seldom the case. The placenta is retained merely because the uterus is deprived of the necessary irritation to cause its efficient contraction. In such instances, the first contractions of the uterus not being supported, the organ becomes, as it were, accustomed to the presence of the placenta; and it remains imperfectly contracted about it, without any further effort at expulsion. In this way, the placenta may remain two—four—six hours in the uterus without being expelled. If the rule which we have laid down be observed, and a steady but moderate pressure be maintained upon the fundus of the uterus during its contraction, this will seldom happen; but if the placenta be not separated then, it is better to wait for some time—from half an hour to an hour—and again to excite the uterus to contraction. For this purpose, the fundus should be brought, as nearly as possible, towards the centre of the pelvis, and grasped firmly with both hands; as soon as it becomes hard, strong pressure upon it is generally sufficient to cause the expulsion of the placenta; if it should not, do not use any violence; rather let the

nurse, under your direction, maintain the fundus in the same position, while you pass the fingers, and, if necessary, the hand, into the vagina, in order to stimulate the uterus to contraction. For this purpose, the funis should be held firmly in the left hand, and the fingers of the right hand passed along it within the vagina. Sometimes this alone excites contraction; but if not, all the fingers, in a conical form, may be introduced within it as far as the os uteri: in doing so you will often find a large portion of the placenta lying at the upper part of the vagina; you may even feel the insertion of the funis, but do not attempt to withdraw it, pass the hand still toward the os uteri, and by irritating it, the portion of the placenta that lies within the cervix is often detached, so that the whole placenta may be removed. If this be not sufficient, withdrawing the hand along the vagina for a short distance will excite contraction; but if both means fail, the fingers must be introduced in the same manner within the os uteri, to dilate it, when the upper part of the placenta may be grasped, and the whole removed. The assistant should press firmly on the fundus uteri while the hand is being withdrawn. In many instances the placenta is found in the upper part of the vagina alone, and can be very easily removed; but no attempt of this kind should be made until the hand has passed above it, so as to have it completely within its grasp. When efforts are made to draw the placenta away by the lower portion, there is always a risk that it may be broken in the attempt, especially if it be caught by the cervix uteri. In one instance which came under our notice, a small portion of the placenta was adherent to the neck of the uterus, and the remainder being dragged away in this manner gave rise to hæmorrhage that terminated fatally.

Another cause of retention of the placenta without hæmorrhage, is irregular contraction of the uterus. This is excited, as we have stated, by drawing the funis frequently, for the purpose of ascertaining whether the placenta is separated. Sometimes, also, it is produced by frictions over the anterior wall of the uterus, or rather, over the corresponding portion of the abdomen. By this manipulation, the uterus is often pushed over to the iliac fossa, where it remains in a semi-contracted state; or the anterior portion of the uterus is contracted, while the posterior, where the placenta is commonly attached, is

relaxed. To remove this irregularity it is necessary, not only to grasp the fundus firmly, as before mentioned, but to pass the fingers over the posterior wall, as low as the abdominal parieties will admit, when, if the irritation excite the relaxed portion, the order of uterine contraction is instantly restored, and the placenta will be immediately expelled. In these cases of retention, it is seldom necessary to wait longer than an hour to have it removed; and if the uterus be carefully attended to during its contraction, and firm pressure afterwards used, if necessary, you will very seldom have any occasion to wait so long, or to pass the hand into the uterus to withdraw the placenta.

With these observations, we shall conclude the management of natural labour, and consider the further treatment of the parturient woman after her delivery in a future part of the course, when we enter upon the subject of lactation.

LECTURE VII.

DIFFICULT LABOURS.

Difficult Labours divided into Tedious and Laborious Labours—Causes of Tedious Labours—Over-Distension of the Uterus—Extreme Obliquity—Gradual Escape of Liquor Amnii—Hysterical Excitement—Mental Despondency—Rigidity of Os Uteri from Inflammation—Toughness of Os Uteri—Extreme Rigidity, etc.

THE first exception to the definition which Denman has given of natural labour is that in which labour exceeds twenty-four hours. It then becomes *difficult labour*, and to this subject we would now direct your attention. According to this definition, when the vertex presents, and no accident occurs, time alone would seem to be the criterion which is to determine the class to which labour is to be referred: but this is not the case, and you must understand this definition to be only of general, not universal application; that while, on the one side, there are cases in which labour is so severe, and the obstruction so great, as to render it difficult, nay, even dangerous, before twenty-four hours expire, there are also instances, on the opposite side, in which labour may be prolonged far beyond the prescribed period, and yet present no difficulty to the practitioner, but the difficulty of sustaining his patience. There are many circumstances, therefore, besides *time*, which should be considered, in reference to difficult labour, and you must only accept the definition we have given you as one which will embrace a larger number of such cases than any other that we might propose.

The causes which render labours difficult vary exceedingly. The delay arises sometimes in the first, sometimes in the second stage. In one instance, the constitution of the patient or the rigidity of the passages, may retard the delivery; in another, the disproportion between the head of the child and the pelvis may

impede the progress of labour. It is necessary, therefore, to classify these causes, and for this purpose we would include, under the head of "*difficult labours*," two subdivisions—1st, That in which labour is merely prolonged beyond the average period, without being, at any time, unusually severe,—it is then called "*tedious labour*,"—2ndly, That in which, without reference to time, there is a powerful struggle carried on by the uterus to overcome some unusual resistance. This may be called by the expressive term "*laborious labour*." The causes which produce the former are most frequently met with in the first stage of labour; those that give rise to the latter, generally occur in the second. These divisions embrace a great variety of causes, which may be classed under several heads. Tedious labour may depend either upon inefficient action of the uterus, or rigidity of the passages; and as their consideration will form the subject of the present lecture, we place these causes before you in a tabular form in the order in which we shall consider them.

Inefficient action of the uterus from—

1. Over-distension of the uterus.
2. Extreme obliquity of the uterus.
3. Gradual escape of the liquor amnii.
4. Hysterical excitement.
5. Mental despondency.

Rigidity of the passages—

1. Rigid os and cervix uteri.
2. Contracted vagina.
3. Rigid perinæum.

Such are the usual causes of tediousness in labour. We might add others which are less frequently met with; but it is unnecessary to propose to you a complicated classification. The first of these conditions, then, is:—

1. *Over-distension of the uterus.*—This cause of delay is not very commonly met with; but when it occurs and prolongs labour, the uterus is so immensely distended by the liquor amnii, that, like the bladder in retention of urine, it is for a time paralyzed. The accumulation of fluid arises from a diseased condition of the amnion, which is often thickened and marked with broad patches of a white colour, as if lymph had been effused between it and the chorion. This thickening of the membranes and dropsy of the amnion generally accompany each other; but the delay in labour principally depends upon the latter cause.

When the uterus is thus over-distended, the grinding pains, which last only for a moment, return with longer intervals be-

tween them, and sometimes they cease altogether ; so that if the cause be unknown, the patient may remain for an indefinite time teased by these inefficient spasms of the uterus, or labour may be quite suspended. Very little attention, however, is sufficient to detect this condition of the membranes. In such cases, the child is generally small, and the evidence, through the abdomen, of its presence in the uterus, is not so manifest. The foetal heart often cannot be heard, the placental murmur is indistinct, or may be absent ; at the same time, the abdomen is greatly distended, and the sense of fluctuation over its surface general. When the vagina is examined, no presentation can be felt, the membranes, tensely distended by the liquor amnii, alone protrude through the os uteri. Such evidences are sufficient to determine the cause of delay, and as the dilatation of the os uteri has generally made some progress with these tardy and feeble pains before the case can be considered tedious, the liquor amnii may be discharged without much risk of the os uteri becoming irritated ; nevertheless, caution is necessary in this simple operation, because if the liquor amnii be discharged suddenly, the violent gush of the fluid may derange the position of the child, or bring down the funis into the vagina. The safer plan would be to puncture the membranes within the os uteri as high as you can reach, so that the liquor amnii may escape gradually. When the uterus is thus relieved, the pains increase in strength and frequency, so that labour generally proceeds rapidly to its conclusion.

2. *Extreme obliquity of the uterus* is the next cause of delay. This obliquity may be either lateral or anterior. When the obliquity is lateral, the uterus is generally inclined to the left side of the abdomen, in place of taking its more usual direction to the right side ; it rests very much upon the iliac fossa, and the pains, which may be strong, have little effect upon the os uteri, although quite dilatable ; this circumstance will excite a suspicion of the cause, but an examination of the uterus through the abdomen will at once determine it.

Mere change of position is sometimes sufficient to remove this difficulty. Those pains which had been short and irregular, and consequently ineffectual, while the patient lay on her left side, become steady and efficient when the patient changes to the right side. The dilatation of the mouth of the uterus proceeds rapidly to its completion, and the labour, that had been lingering for

hours before, is often terminated by a few strong pains. If this mode be not sufficient for the purpose, a broad bandage may be applied round the abdomen, and drawn towards the right side as firmly as the patient can conveniently bear.

When the uterus is inclined very much forwards, it may arise either from the pelvis being so shaped that the axis of the brim is nearly horizontal, or from weakness of the abdominal parietes; possibly from both combined. In these instances, the uterus is sometimes so completely displaced, that the direction of the fundus is reversed, and the uterus hangs down over the pubis so as to rest upon the thighs. Labour is, of course, impeded; but if the bandage be so applied as to draw the uterus gradually upwards, or even if it be supported by the hand during its contractions, labour will often advance rapidly to a favourable termination.

It is advisable to allow the patient to lie more upon her back than her side, but as there is a great difference in the position which different women find most favourable to their pains under these circumstances, you will sometimes meet with cases where they lie, not only on their side, but with the body completely across the bed. This difference may depend upon the cause of the obliquity: if it arise from weakness of the parietes of the abdomen, to lie on the back would be the most likely position to correct this deviation; if it depend upon a horizontal aspect of the pelvis, the body would be inclined forwards, to lessen its distance from the axis of the uterus. It would be unsafe, however, to theorize too much on such a subject: and happily it is the less necessary, because nature generally prompts the woman to adopt the position which is the most favourable for her. You should not insist, therefore, too much on any given position, but rather let your patient try different postures, and accommodate yourself to that which she feels to be the easiest. The mouth of the uterus is often greatly displaced in these cases: it is directed very much towards the promontory of the sacrum; and hence, in order to correct the obliquity, some have advised that the forefinger be passed within the opening, and the os uteri drawn towards the centre of the pelvis. How is it possible to alter the pendulous fundus by such means? But if, in order to correct the position of the fundus, it is also necessary that it be raised by the opposite hand, the introduction of the finger is not required, because then

the mouth of the uterus will correct itself. Such means, therefore, should be avoided, because they are calculated to excite irritation. Some serious mistakes, however, may be made, as to the cause of delay, when the os uteri is absent from its usual situation. For instance—when the anterior lip has become so thin as to resemble the membranes, it may be mistaken for them, and this extreme tenuity sometimes takes place when the anterior lip becomes the most dependent part of the uterine tumour. The delay might easily be attributed to rigidity of the membranes, and the supposed membranes—that is, the uterus—accordingly punctured. Another error is, the supposition that the os uteri is either closed by a cicatrix, or is imperforate. A mistake of this kind has led to the sagacious operation of dividing the cervix to make an artificial os uteri: such an instance has been mentioned by Dewees.* You should, therefore, always make the most careful vaginal examination, and you will seldom fail in finding the os uteri (often not larger than a sixpence) directed upwards towards the promontory of the sacrum.

3. *The gradual escape of the liquor amnii* also gives rise to tediousness. If this take place when the os uteri is slightly dilated, in other words, when the latter is so long exposed to the pressure of the head of the child as to become irritated by it, the result is rigidity of the os uteri, and its consideration falls naturally under that head. But if the os uteri be dilated to any extent, and not easily excited by irritation, this accident may have an opposite effect, and hasten rather than retard delivery.

We would now direct your attention to much more important causes of delay—viz., those which depend upon the temperament of the patient: such as are produced by high mental excitement, or by its opposite, great depression of mind, acting through the constitution on the uterus.

4. *Hysterical excitement* is the first of these causes to which we shall allude. Hysteria is one of the most frequent affections of the female constitution. In the course of your practice, you will meet with many instances of that highly-excitible temperament which is associated with this disease. When these hysterical females become pregnant, and the period of parturition arrives, the practitioner has often to undertake the management of a case

* Dewees' Midwifery, p. 90.

far more embarrassing than those in which mechanical impediments obstruct delivery.

As the time of her trial approaches, the patient becomes anxious and excited; she has perpetually before her mind this long-looked-for event; her attention is constantly alive to the most trifling circumstance that has reference to the suffering she has to undergo. Consequently, when at length the moment arrives, her mental excitement is strained to the highest pitch; labour has scarcely commenced when it is suspended by her anxiety, and is again renewed only to be again interrupted in a similar manner. Proceeding in this irregular course, the first stage of labour may be prolonged considerably beyond the time it would otherwise occupy; and when this is the case, the patient (who from the beginning was not very tolerant of her suffering) being taken as it were by surprise, and never anticipating such delay, becomes still more anxious. She may possibly be alarmed for her safety; and then her mental inquietude bursts through all restraints; she begs urgently to be released from her agony, and if you hesitate to comply with a request to which you dare not accede, she loses all confidence in you; her impatience knows no bounds; every pain is interrupted by the most violent exclamations, and sometimes terminates in a fit of hysterical convulsions. Such a case, therefore, requires the most careful attention, not only in its medical treatment, but in the circumspection that is necessary in its management. Conversation should not be permitted within the hearing of the patient; none but her immediate friend and the nurse should be allowed to remain in the room; and while you use every exertion to give her encouragement, and, if possible, to lessen her excitement, you must not commit yourself by hasty promises. With regard to medical treatment, your chief attention should be directed to the state of the bowels. If it be in your power to enter upon a course of treatment before labour has commenced, your chances of success will be greatly increased. In many of these cases there is great constipation previous to parturition; but in all the evacuations are of an unhealthy character, dark, viscid, scanty and offensive. A preliminary course of alteratives, combined with stimulating aperients, will correct this condition, and lessen the hysteric influence; but if you are not given the opportunity for such treatment, and are for the first time called upon to take charge of the case when labour has actually com-

menced, it would be advisable to have an assafœtida enema given before labour has made any progress. Scybalæ lodge frequently in the rectum and large intestines, causing great irritation, which contributes to impede the action of the uterus. When the bowels are unloaded, narcotics, combined with the diffusible stimulants, will more efficiently control the irritability of the patient. Women of this temperament are often greatly fatigued by their own useless and impatient efforts to hasten their delivery, as well as by the length to which labour may be prolonged. In such instances, a full anodyne frequently procures some hours' refreshing sleep; after which the pains become more regular and active. If you should find that, notwithstanding such treatment, the action of the uterus is still weak and inefficient, ergot of rye may be given in moderate doses to excite it; but its use should follow the treatment previously mentioned, and only when the uterus continues to act feebly, although the nervous irritability of the patient has been tranquillized. A judicious application of these adjuvants will aid you in exciting regular and dilating pains, and will bring this stage to a favourable conclusion. If the first stage be not so much prolonged as to exhaust the patient, the second will generally terminate without much difficulty: these hysterical symptoms often disappear when the expulsive efforts of the uterus absorb the attention of the patient. Cases of this description are embarrassing to the practitioner from their tediousness, but seldom terminate unfavourably, if they are properly treated. The same remark does not apply to the next cause of delay.

5. *Mental Despondency*.—This source of difficulty is but briefly alluded to by the majority of obstetric authors. "Depressing passions of the mind" are enumerated among the causes that retard labour, but are not dwelt upon in proportion to their importance. Fortunately, extreme cases of this kind are rarely met with; but instances might be quoted in which death was the result of such a cause. Cases are occasionally recorded of *unaccountable sudden death* after labour, which might perhaps be explained in this way, if all the circumstances of the case were understood; at least, the few instances that have fallen under my notice, seemed to admit of such an interpretation. In one case death would undoubtedly have taken place, had not the cause of depression been so obvious.

A poor, emaciated woman entered the Dublin Lying-in Hos-

pital, January, 1834, to be delivered of her eighth child. "Sharp misery had worn her to the bones;" her pulse was feeble, the action of the uterus weak; notwithstanding this, she was delivered in an hour after admission; no hæmorrhage took place, and the placenta was separated without any difficulty; but her delivery was followed by the most alarming depression, which required the utmost care and attention to prevent her sinking altogether. Fortunately, strong beef-tea, and other nutritious diet, had been given to her from the time of admission, so that, with the addition of stimulants, and maintaining the temperature of the surface, she gradually recovered. This was a case where poverty and starvation produced their usual effects, and consequently one more under the control of treatment than those melancholy instances in which some cause operating on the mind alone, produces some extreme nervous shock which we cannot relieve, because we cannot "minister unto a mind diseased." An instance of this kind occurred in the same institution the following year, January, 1835. A young woman was admitted in labour of her first child. She was evidently above the class of persons usually admitted into that establishment. She seemed rather to shun observation; and there were no symptoms attending labour that required interference. It proceeded to its conclusion without any interruption, and terminated within ten hours from its commencement. The pains were feeble, but they were sufficiently strong for the purpose; the patient herself appeared also weak. She was delivered of a girl; and in about half an hour after the placenta was expelled; but the pulse instantly sunk, syncope followed, and every means that could be used failed to prevent dissolution, although the discharge from the uterus was not increased, nor was there the least evidence of hæmorrhage, either externally or internally.

An inspection was made twelve hours after death, and no cause could be discovered to explain an event so unlooked for: her history, however, may do so. She had been one of a respectable family, delicately reared, and educated in the strictest moral principles. She had been seduced, betrayed and deserted; and, to complete her miseries, had to endure her hour of trial in the reception-ward of the Dublin Lying-in Hospital. I shall only mention another instance of this kind, which will, perhaps, more distinctly illustrate the effect of extreme nervous shock.

In the beginning of the year 1834 a poor woman had walked

some distance to the Dublin Lying-in Hospital, and when near it was suddenly seized with the pains of labour. She was delivered in the street, and with much difficulty brought into the house before the placenta separated. It came away, however, without difficulty; and the trifling hæmorrhage that followed was easily arrested. Her alarm was very great, but after some time it subsided; she slept, and nothing further occurred out of the usual course until the following day. On that morning a patient was brought into the same ward to be delivered who was extremely boisterous; she occupied the next bed to this woman, who lay so quietly that she seemed to pay little attention to the disturbance. In the course of the day, however, she complained of being overcome by her cries. She felt faint, as if she were sinking; she had slight pains in the epigastrium, some sickness of stomach, pulse rather rapid, compressible and soft. The woman who caused this was fortunately delivered, and thus all further annoyance was removed; but this patient did not recover from the effect that it seemed to produce on her. Stimulants were given to her, the extremities and surface kept warm, and the most perfect quietness observed in the ward—but all to no purpose. In the evening she was seized with syncope, so alarming as to excite the greatest apprehension for her safety; the extremities became cold, her motions passed involuntarily, and she died in about three hours. The uterus was perfectly contracted; there was not the slightest appearance of hæmorrhage from the vagina, nor any symptom present to explain the cause of dissolution.

A very careful inspection was made after death: all the viscera of the abdomen were quite healthy, the uterus firm, and contracted to its usual size. There were some old adhesions in the lungs; the heart was small, and contained very little blood on the right side; the vessels were all sound; and the only alteration in the brain was an increased quantity of serum in the ventricles and at the base. No other explanation, therefore, was left, but the probable one, that she sunk in consequence of extreme nervous shock. Her own sudden delivery produced a strong impression on her mind in the first instance. This was again excited and increased by the violence of the patient alluded to, and hence the effect. It is probable that she would have recovered from the first shock, had it not been again renewed by this accident.

These instances will illustrate the influence of the mind on

the constitution at this critical period ; they are fortunately rare, but those cases where the same cause operates in retarding, and sometimes in suspending, the action of the uterus, are more frequently met with. The sympathy (to use a popular term) that exists between the brain and the uterus is matter of daily observation, the change of feelings and temper that frequently result from pregnancy, the hallucinations that occur after delivery, from the slightest temporary aberration to long-continued mania, all prove the influence of the uterus on the mind. So, on the other hand, a disturbed mind suspends the action of the uterus, just in the same manner as it interferes with the healthy action of the digestive organs. As in the latter class of cases you find the appetite gone, the digestion imperfect, the liver disordered, and the bowels constipated, so in the former parturition may be greatly prolonged, and the patient recovered with difficulty from the effect of a labour that otherwise would have been happily concluded within the average period. Such cases may come under your notice ; it is therefore necessary to recollect their characters.

They differ altogether from the former class of patients ; there is no restless excitement about them ; on the contrary they submit to their suffering with a quiet resignation, which might be called fortitude, only that the feeble pulse and listless expression pronounce it the indifference of despair. The patient meets her trial without hope, with the settled conviction that she will not escape. The sufferings that attend the birth of her offspring are only lighter pangs added to the accumulation of sorrows that have already overwhelmed her ; she therefore makes no complaint, no inquiry ; but the pains are feeble, the dilatation of the os uteri is consequently slow, and the labour protracted. The uterus is evidently unequal to the required effort.

You have no clue to unravel the intricacy of these symptoms, as you are, of course, never informed of their cause. Nevertheless, the quick and feeble pulse may excite your suspicions ; the tendency to chill and the coldness of the extremities will increase them, and the constant, although passive, watchfulness of your patient, will confirm your apprehensions.

The moment you perceive the nature of the case no time can be lost. A treatment is required the opposite to that generally employed. Stimulants may be given moderately, carefully

observing their effect, the temperature of the surface and extremities attended to, and the bowels (which are always constipated) relieved by warm and stimulating enemata. Ergot of rye, in moderate doses, to excite the specific action of the uterus, is useful.

You may thus succeed in securing a favourable dilatation of the uterus, before the patient becomes exhausted; so that, in the second stage, the uterus may retain sufficient power to complete the delivery in a short time; but if this be not the case, artificial assistance becomes necessary, in order to abbreviate its duration as much as possible. When the child is partly born, you must be careful, also, not to withdraw it too rapidly from the vagina and uterus, because the danger that attends the case is not confined to the effect of tediousness in labour; you have still to guard against the syncope that may follow the complete contraction of the uterus. It should, therefore, be permitted to expel the remainder of the child very gradually, while an equable pressure is maintained by a broad bandage over the abdomen.

The SECOND DIVISION of causes that render labour tedious, is, *rigidity of the passages*. This may exist either in the cervix uteri, vagina, or perinæum. At present we shall confine our attention to rigidity in the cervix uteri. In a former lecture, we took occasion to point out to you, that under the term "rigidity" were included different states of the os and cervix uteri, presenting different degrees of resistance. We shall first speak of those cases where *the rigidity has been induced* by some accidental cause, and secondly, consider *idiopathic rigidity*.

If the os uteri be much exposed to irritation, it is rendered rigid; the lips become swollen, hot, and tender: when these signs of commencing inflammation present themselves, the cervix is less disposed to yield to the action of the uterus, and become rigid. One of the most frequent causes of this kind of rigidity is the *gradual escape of the liquor amnii*, by which the head of the child descends upon and irritates the cervix. If this irritation be long continued, you have to contend, not only against the effect of inflammation, but also against a spasmodic contraction of the circular fibres of the body round the child. When this happens, an additional resistance is offered to the

action of the fundus. In the treatment of it, therefore, promptitude is necessary. If the patient be strong, plethoric, and disposed to make violent straining efforts, a free depletion from the arm will be of much use; it diminishes the tendency to inflammation, and produces a feeling of exhaustion in the patient, which induces her to bear her pains more patiently. In order to insure such an effect, depletion may be followed by tartarized antimony, in small doses, so as to excite nausea. Women who may have been previously very violent and intolerant of their pains, are soon subdued when the sense of exhaustion that attends sickness is excited. If, on the contrary, your patient be of an opposite temperament, this treatment cannot be employed; local depletion is preferable; a dozen leeches may be easily applied to the cervix uteri; warm emollient enemata may be given, and if the woman be much fatigued, or if the pains become feeble and irregular, an anodyne is often very beneficial; some sleep is procured, the irritation of the cervix is diminished, the spasmodic contraction of the fibres disappears, and the pains return with more regularity and strength.

Another cause that renders the cervix uteri rigid, from a tendency to inflammation, is its accidental compression by the head against the pelvis. Sometimes the mouth of the uterus, partially dilated, is driven down with the head into the pelvic cavity, where it is sometimes tightly wedged between the head and the pelvis. Cases are recorded in which the whole cervix uteri has been completely separated by a circular slough, and expelled with the head. In order to avoid this, it is necessary to use every means to combat inflammation, and to prevent, as far as possible, the effects of extreme pressure. Local depletion either by leeches or by scarification, may be employed with advantage, and when the pains cease, the head should be pressed back towards the brim, to relieve the constriction of the cervix uteri. This plan of treatment, with warm emollient enemata, which act as fomentations, will generally succeed; but if the impaction be so great as to render it impracticable, instrumental aid becomes necessary. The consideration of this we must defer to another opportunity.

Again: the cervix and os uteri may suffer only a partial constriction, and thus become inflamed, and retard delivery. The head of the child may rest on the pubic side of the pelvis in

such a manner as to compress the anterior lip of the uterus, and prevent its dilatation. A band is thus formed before the head, which when long pressed upon, is swollen, tender, and rigid. The treatment of the cervix when in this state has become a kind of *vexata questio* in obstetric practice. Some practitioners of station and experience have advised that the anterior lip of the os uteri be pushed up by the fingers, above the head, in the interval of the pain, and there maintained until the returning contraction of the uterus drive the head below it; while others of equal reputation deem such practice to be objectionable, and calculated rather to increase than to diminish the difficulty, by exciting more inflammation. I may mention the late Dr. Hamilton, of Edinburgh, as holding the former opinion; Dr. Collins, of Dublin, the latter; as a proof how men of very extensive experience often arrive at opposite conclusions on what would seem to be a simple practical point. It is my duty neither to draw you into controversy nor to give you too dogmatic an opinion on a question so nicely balanced by authority. I must assume, on the evidence of Drs. Hamilton, Burns, and Breen, that this kind of artificial dilatation may be accomplished in some instances with safety and success. My own experience, however, confirms that of Dr. Collins, and is opposed to this practice: the opportunities I have had of putting it to the test have taught me that success is by no means so easy as it is described to be; that the anterior lip may be pressed back again and again, and yet return to the same situation as before; that it is difficult to get the head to pass the introduced finger; and that these attempts, when unsuccessful, only increase the swelling and inflammation of the soft parts. I have also met with instances, in which, after a failure of this kind, the anterior lip was retracted without any assistance, when the contractions of the uterus succeeded in at length altering the direction of the head. It appears to me, therefore, that this kind of manipulation may be employed, and would be serviceable if the fingers were placed against the head of the child, in order to relieve the constriction of the anterior lip, and to direct the head more towards the pelvic cavity. I am still, however, disposed to object to the practice of artificial dilatation of the mouth of the uterus for the former purpose. While on this part of our subject, I must apprize you of a deception which

may be produced by certain states of the os uteri. The anterior lip is sometimes hypertrophied, and projects so much before the head that it seems to be the cause of difficulty. It is scarcely necessary to say, that you will employ your time very fruitlessly, if you attempt to push back this hypertrophied portion of the os uteri.

That resisting condition of the cervix uteri, to which we have applied the term "*toughness*" (an expression borrowed from Dr. Hamilton), is often met with. The os uteri forms a thick gristly ring, rather dry, and without tenderness. It yields very slowly to the repeated efforts of the uterus, and hence this stage is generally very much prolonged; but so long as inflammation is not excited, no injury is caused by the time occupied in dilatation; nay, more, these cases often recover more rapidly than where the labour is short, although severe. The treatment is altogether of a negative character; rather to watch its progress attentively than officiously to attempt to hasten it. Every care should be taken to prevent inflammation taking place; and vaginal examinations should be made as seldom as possible. The patient should not be suffered to fatigue herself by endeavouring to assist the pains, and by fruitless efforts to shorten her sufferings. The room should be kept perfectly cool; and, as there is generally very great thirst, heating drinks should be strictly forbidden: warm emollient enemata may be given with advantage; and if a stimulus be required to excite the action of the uterus, the addition of common salt in a moderate dose will generally answer the purpose. In such a case as this the first stage may occupy thirty, forty, fifty hours without injury, but if attempts be made to hurry it to a completion, either by mechanical dilatation or by ergot of rye, they will only cause what you are, or should be, always anxious to avoid—inflammation. The former is a direct, the latter, an indirect irritant, because, when the uterus is much excited to contract, by the specific action of the medicine, the membranes are broken prematurely and the head driven forcibly down upon the undilated os uteri: thus your difficulties are rather increased.

The patient may become fatigued and dispirited by the continuance of this stage, and the pains feeble and inefficient. When such is the case, a full dose of opium is often very ser-

viceable; and the nearer the time of natural rest at which it is given, the more likely it is to succeed. A woman who has been fatigued by constant pain during the day, will often sleep the greater part of the night after an anodyne, and awake quite refreshed by her sleep. The dilatation of the os uteri then proceeds more rapidly to a favourable conclusion. Time, in fact, is the only remedy for this condition of the passages.

The last kind of rigidity to which I shall have to allude is that in which *the os uteri is like cartilage*, and will not yield to the most powerful and constant action of the uterus; the membranes are usually broken and the waters discharged early in this stage, and therefore the uterine action is increased to its full extent. Inflammation is the almost certain consequence of the struggle that ensues; you have, therefore, complications of the worst description to contend against. The issue of a case of this kind is the spasmodic and irregular contraction of the uterus about the body of the child, and frequently its death, before the uterus is opened to any extent. It therefore becomes a case for delivery by perforation. But there are some instances in which the dilatation is brought to a successful termination by extreme care in the management of the case. As a preliminary treatment, the frequent use of the warm bath, although an exception to ordinary rules, is found to be very beneficial. When labour begins, warm emollient enemata should be given from time to time, and the patient placed at once under the influence of tartarized antimony. General depletion may be employed if the patient be robust; if otherwise, local depletion is preferable, and it is indicated as soon as the least tenderness of the os uteri is observed.

If these means fail, it becomes a question whether we should wait for the death of the child, in order to remove it by the crotchet, or incise the unyielding cervix. The former practice involves a sacrifice of life, but generally secures the mother from the injurious effects which may follow. The latter may be the means of preserving the child, but if the incision lead to a laceration of the uterus, the mother is at once placed in the most imminent danger of her life. The fear of such a consequence, it appears to me, has prevented any attempt being made thus to cut through this Gordian knot of difficult labour in its first stage, but whether this, like other operations, is only

surrounded by chimeras of the imagination, which some bold spirit will dissipate, remains yet to be proved. Incision has been performed without accident; the same may happen again, and I confess, in a case such as I have described to you, I should be more disposed to adopt the shorter course, in the hope of saving the child, than to wait until its death enabled me to remove it. This, however, is but an individual opinion, and needs support.

LECTURE VIII.

LABORIOUS LABOURS.

Causes of delay in the Second Stage of Labour: Head of the Child misplaced—Too large, and too much Ossified—Hydrocephalic—Pelvis like the Male Pelvis—Deformed—The Ovate Pelvis—The Cordiform Pelvis—Head of the Child above the Brim of the Pelvis—Fixed in the Brim—Arrested in the Cavity—Characters of the Arrest and Impaction of the Head.

THE causes of delay in the first stage of labour have been detailed to you in the preceding lecture; those which occur in the second stage we have now to consider. The former gives rise to labours that are called "*tedious*;" the latter, to those that are termed "*laborious*," in consequence of the extreme severity of the struggle that ensues in the effort to force the head through the pelvis. Difficulties in the second stage of labour are produced either by disproportion between the head and the pelvis, or by some mechanical impediment obstructing the head in a pelvis that otherwise would permit it to pass. In cases of disproportion, the cause may exist either in the head of the child or in the pelvis, or in both combined.

We shall therefore direct your attention to each separately—1st. To the head of the child as a cause of delay; 2nd. To disproportion in the pelvis; and having considered the management of, and rules of practice applicable to, cases of this description, we shall then briefly conclude with some observations on those mechanical obstructions that are sometimes met with.

1st. *The head of a child* may be a cause of difficulty from its *irregular position*. It may be *too large*, or *too much ossified*, or it may be *hydrocephalic*.

We have already sufficiently dwelt upon the positions of the

head; very few observations upon them as causes of delay in labour will now be necessary. Some of these positions, as face and face-to-pubis presentations, have been assumed as, rather than proved to be, causes of difficulty. We have shown you that the majority of these cases terminate within the ordinary period. Nevertheless, there are some exceptions in which labour is prolonged, and in which danger may arise if a judicious treatment be not adopted.

When the head is arrested with *the face towards the pubis*, you will have no difficulty in detecting the position by the rule we have laid down; neither, when it ceases to advance, is it hard to correct. Sometimes the correction may be made with fingers, but it is preferable to introduce the vectis on the pubic side of the pelvis, to press the head back in the intervals of the the pains, and to rotate it gradually towards the sacral side. When I say "gradually," I mean that you should rather trust to frequently repeated efforts to give the head its proper direction, than to make a violent attempt to change the position in the first instance. The adoption of such a course would only end in disappointment, and complicate your difficulties. It is obvious that in these cases the head should not be suffered to remain too long arrested, because it may be impossible then to alter it, and you will be compelled to extract the child with the forceps—an operation which, in such a case, would be one of no little difficulty—dangerous to the soft parts of the mother, especially to the perinæum.

If *the face-presentation* be in a similar manner arrested, as few vaginal examinations as possible should be made, merely such as are sufficient to determine the arrest; when this is ascertained, recourse must be had to the forceps. But it is very improbable that you will meet with a case where such an operation is necessary. That accidental displacement by which the antero-posterior measurement of the head is fixed across the cavity of the pelvis, has been already alluded to. This is the most easy to correct of any of these deviations, but if it be left too long arrested, you may not be able to press up the forehead with the fingers; the vectis may therefore be applied for the purpose of altering the direction. The last irregularity that we shall speak of is when *the forehead becomes the presenting part*. In these cases, the forceps must be employed to extract the

head. You will find it to be a very difficult operation, and one which requires great caution in the performance. These are the principal varieties of position which may cause a delay in the second stage of labour.

The cause of difficulty we have stated to be that in which *the head of the child is too large and too much ossified*. This will require a more attentive consideration, because it appears to me that this increased development of bone occurs most frequently in male children; and through the elaborate researches of Professor Simpson, of Edinburgh, we now have statistical proof—1st. “That the dangers and difficulties of parturition are *greater to the mother* in male than in female births;” and 2nd. “The dangers and accidents from parturition and its results are *greater to the child* in male than in female births.” Professor Simpson correctly attributes these effects to the greater size of the head at birth in male infants, a fact already noticed by the late Dr. Joseph Clarke, of Dublin. I should be disposed to add, that not only is the size greater, but the ossification of the bones is more advanced, and therefore they are less disposed to yield. The head of the healthy male child is rounder, and the fontanelles smaller. That of the female, more oblong, and the fontanelles more distinct. These characters, along with the increased size, contribute to produce greater difficulty in its passage through the pelvis. The head being too much ossified, is therefore a very frequent, and too often a very embarrassing cause of delay in the second stage of labour. When a vaginal examination is made, the posterior fontanelle is not distinctly felt; it seems to be only a central point to which the lambdoid and sagittal sutures converge; the sagittal suture is indistinct, the head presenting a round, firm, equal surface. When the head is so formed, it will generally pass safely through a pelvis of the average proportions, although slowly and with difficulty. It is when the pelvis deviates from the standard, and is diminished in its proportions, that these difficulties increase to a dangerous extent.

As far as I have had the opportunity of judging, this kind of head is very often met with when the pelvis is in a similar condition—too much ossified. I shall have again to bring before your notice the varieties of the pelvis which have been described to you in the commencement of the course, and I shall beg your

attention especially to that variety which possesses many of the characters of the male pelvis, and to take it in connection with the large, round, ossified head of the male child. This is of essential importance when we have to compare it with other varieties in the head and in the pelvis, and to derive from these facts a conclusion as to rules of practice. At present I shall only direct your attention to the propositions I have quoted as established by Professor Simpson. The valuable report by Dr. Collins, of the practice of the Dublin Lying-in-Hospital, has been the basis of his calculations. I am happy to have it in my power to confirm Prof. Simpson's statements by the results of my own experience in the same hospital. Having noted all those cases in which the duration of labour extended to or exceeded twenty-four hours, the following results were obtained in reference to this question :—

There were in the total number of patients delivered (5699), 213 such cases, of which five were twin births ; 126 of these were boys, and only 92 girls, or in the proportion of three to two. Forty-six of the boys died, thirty-five of the girls, being nearly one-third of the whole number. About eighty of these 213 cases occurred when that hospital was under the superintendence of Dr. Collins, and are therefore included in 16,654 cases reported by him ; the remainder were observed during the two years subsequent to that report. From these facts, then, you may conclude, that the majority of cases where labour is difficult are those with male children ; and as the greater size of the head is proved to be the cause, the difficulty must be in the second stage, and they belong, therefore, to the class of laborious labours which we are now considering. We have stated that in these instances also the head is more advanced in its degrees of ossification. Hence you will perceive that this cause of difficulty, dependent upon the child, embraces a very large and important class of cases.

The next cause is of a very opposite character. *The hydrocephalic head*, it is true, is enormously increased in its size, but its ossification is retarded ; and it might rather be compared to a bag of fluid than to the solid tumour which the head generally presents. Hence some of these cases are attended with no difficulty ; and instances have occurred to me, in which the head, though morbidly enlarged, passed easily through the pelvis, but,


on the other hand, some of the very worst cases in obstetric practice have depended upon this cause. As a proof, I place before you the very opposite results of the practice of Dr. Collins and Dr. Lee on this point. Dr. Collins gives six cases in which the child was still-born, from hydrocephalus: In three of these cases labour lasted only one hour; in a fourth only two hours; the fifth was twenty-six hours in labour, not stated to be severe; the sixth was brought into the hospital, after being thirty-two hours in severe labour. In none of these cases did the mother suffer. Dr. Lee gives five cases:—one, twenty hours in labour; one, sixty hours; one, seventy-two hours; the fourth stated as being “very long;” the fifth, “too long” in labour. In three of these cases, the mother died; two of them in consequence of the uterus being ruptured. These last two cases are detailed at length by Dr. Lee in his lectures.*

If it were necessary to bring forward additional proofs to convince you of the importance of educating your sense of touch so as to distinguish the different positions of the head, such instances as these would afford abundant evidence. It is obvious to you, that if the cause of delay were known to be hydrocephalus, no patient should be suffered to remain “very long in labour.” The disease is fatal to the child; and if the head be arrested, there is no other operation than to perforate the head and discharge the fluid. The practice is self-evident when the cause of difficulty is understood. But when a patient is allowed to remain “too long in labour,” under such circumstances, it can only arise from an imperfect education in the sense of touch, in consequence of which the true position of the head is unnoticed. Those who are quite satisfied that the head presents, without caring much how it is situated, will be liable to those mistakes; but when, by a constant and careful attention, you acquire facility in recognizing its different positions, you will at once detect the remarkable change which hydrocephalus produces in the characters you have been accustomed to notice. The increased size of the posterior fontanelles, the mobility and separation of

* It is right to state, that these cases, recorded by Dr. Lee in his “Clinical Midwifery,” (pp. 54–60,) were cases in which Dr. Lee’s assistance was required, not those attended by him from the commencement of labour.

the sagittal suture, the great overlapping of the parietal bones during a pain, if the head have entered the pelvic cavity, and the general looseness of these bones, would sufficiently point out the nature of the case and determine the practice.

Such are the principal causes of difficulty depending upon the head of the child, in the second stage of labour. The most important of these is certainly that, in which the head is too large and too much ossified—the head of the male child. Let us now turn to the difficulties presented by the pelvis; and, to prevent confusion, we shall assume that the head presents in the first position.



Masculine Pelvis.

It will not be necessary again to enter upon a detailed description of the varieties in the irregularly formed pelvis; it will be sufficient to recapitulate briefly those leading points of difference that are of practical importance; and with this view we would beg of you to consider attentively that form of pelvis which has been described as *resembling the male pelvis*, to compare it with *the pelvis deformed by disease*, and to observe the points of contrast between them. The first to which we would direct your attention, is the difference in the degree of ossification. The pelvis resembling the male is more osseous than the standard pelvis; the diseased pelvis less so. The former we shall call (for brevity) *the masculine pelvis*, and proceed to consider its especial characters. *Its weight* is greater than the ordinary pelvis, in consequence of the increased deposition of bone; but, what is more

important, this deposit occurs in most unfavourable situations. The symphysis pubis is very narrow, and a bony ridge is sometimes formed behind it. The spine of the ischium is longer, and drawn inwards. The tubera ischii are larger and rougher, from the same cause. There is much less mobility in the sacro-coccygeal articulation, and the sacro-iliac synchondrosis is perfectly unyielding.

Its *shape* is unfavourable to the passage of the head. The brim is rather lessened in its transverse and oblique measurements; but the difficulty which this pelvis presents is not generally at the brim, it is rather at the pelvic cavity that an arrest or impaction takes place. The cavity is much deeper than the standard pelvis, and becomes narrower towards the outlet. The head often passes down almost to the outlet, and then becomes *impacted*, because the arch of the pubis is too narrow to allow the occiput to escape from the cavity; it is therefore driven down still further, and becomes fixed between, or above, the tubers of the ischia. Its advance is also opposed by the resistance of the coccyx, and perhaps the spines of the ischia. The head may be arrested before it reaches the outlet, in consequence of the difficulty of its rotation towards the conjugate axis of the pelvis.

If, in addition to these impediments depending upon the shape of the pelvis and its increased ossification, you should be obliged to contend with obstacles arising from the large and ossified head of the male child, a very embarrassing combination of difficulties would present itself, requiring the most skilful treatment to bring to a successful issue. They are, however of a character perfectly the opposite to those more generally described—the difficulties of the deformed or diseased pelvis—the only cause of embarrassment depending on the pelvis, which is pointed out by the majority of obstetric writers. Let us, then, contrast the one with the other.

The softened pelvis has a *less deposit of bone* than the standard pelvis. The symphysis pubis is often wide and perfectly smooth; the tubera ischii small. The articulations are less resisting, and I think I might add, on the authority of some recorded cases, the pelvis itself is capable of some degree of expansion. The shape of the diseased pelvis is never uniform. In that which is generally attributed to *rickets*, the brim is oval, and, in the extreme deformity, is of an hour-glass shape; the cavity is shal-

low ; the outlet very wide ; the coccyx abruptly curved.* The principal difficulty in such a pelvis is at the brim, the antero-posterior measurement being too narrow ; but if this be overcome, no other impediment presents itself but the coccyx, which although abruptly curved, is generally sufficiently flexible to oppose but a slight resistance to the head of the child. In that deformity which is ascribed to *mollities ossium*, in the *cordiform pelvis*, its irregularities impede the passage of the head, both in the brim, cavity, and outlet.† If the head pass through the cordiform brim, it is still opposed in the cavity by the planes of the ischium, which are pressed inwards : the pubic arch is also too narrow to allow the head to escape. The coccyx is abruptly curved in the same manner as in the former pelvis, from which it differs in the close approximation of the tubera ischii. The extreme of such a deformity would render the passage of the child impossible ; it is only, therefore, to cases of slighter deviation into this irregularity that our present remarks must apply ; we would call your attention to those instances alone where the head of the child may pass through the pelvis slowly, or may be drawn through it with difficulty. Contrasting a pelvis of this kind with the masculine pelvis, and assuming that the pelvic cavity in both presents the same apparent contraction, still the difference in the degree of ossification must cause a corresponding difference in the degree of opposition to the passage of the head. The articulations of the softened pelvis, although not actually moveable (even this is asserted by some), are yet capable of yielding, to a certain extent ; the bones themselves may admit of slight expansion, and if we might admit the probability that the head of the child is also less ossified, you will perceive a strong contrast to the difficulties which the masculine pelvis presents under apparently similar circumstances.

It is necessary to request your attentive consideration of these differences, in order to render intelligible to you some very difficult questions connected with the rule of practice in such cases. The contrast is not confined to the pelvis alone ; but, as you might naturally anticipate, there is a similar difference in the constitutional strength and temperament of the parturient woman.

* Fig. 1, p. 42.

† Fig. 2, p. 42.

We have already briefly alluded to the characteristics of those strong, muscular, masculine women with whom the pelvis is unusually ossified, who have often great rigidity of the passages, and with whom we sometimes find that almost undilatable rigidity of the os uteri. Let us now point out to your notice those feeble habits whose unhealthy constitutions are indicated by these evidences of disease in the pelvis. They are generally of a leucophlegmatic temperament; the subjects, perhaps, of thatameleon disease, hysteria, therefore nervous and excitable, of fair complexion, soft skin, bones fine, but swollen at the joints, the flesh often flabby, and the tissues relaxed. In such habits the os uteri is seldom rigid, the vagina is smooth, and very yielding, and if leucorrhœa have existed, may be even flaccid; the perinæum is also quite dilatable. Inflammation is not so readily excited in the passages as in the former case, and the whole difficulty of the case in this second stage, is, as it were, centred in the pelvic deformity.

It should be also noticed, as a constitutional difference, that in these feeble habits the uterus does not possess the same energy; it is sooner fatigued, and exhaustion would be more readily induced if its action were greatly prolonged, than in the former class of patients. You perceive, therefore, that two cases which present precisely the same apparent amount of resistance to the passage of the head may be perfectly opposed in every other respect, and consequently, that the same treatment, if applied to both, might be mischievous to either. It is essential to point this out to you, because rules of practice are too often laid down by authors as being applicable indifferently to all cases which present the same amount of disproportion, and hence a great deal of useless controversy has sprung up as to those rules, in consequence of the opposite experience of the disputants; for example, if the question be as to the rule when instrumental aid is called for. The practitioner who meets with the diseased pelvis as his chief cause of difficulty, and living perhaps in a manufacturing town, or other unhealthy district, who finds the majority of his patients to be those we have just described to you, such a practitioner would find that he could not suffer his patient to continue in labour beyond a certain period, without great hazard to her life and to his own reputation. He would also have it in his power to deliver his patient with the forceps successfully

where the disproportion was such, that had it been the masculine pelvis he would have done irreparable mischief. It is not surprising, therefore, that his opponent, accustomed to a healthy district, and meeting with cases where the chief difficulty arises from a pelvis of the latter kind, should wonder and protest against such temerity. The judgment of each, as to the rule, may be correct, but the experience upon which it is founded being different, the opinions must of necessity be opposed when they attempt to argue *a particulare ad universale* (as logicians have it), without exercising a just discrimination on those points of difference that I have attempted to lay before you.

Having made these general observations on the irregularities of the pelvis, as causes of delay in the second stage of labour, permit me now to direct your attention to the manner in which the head of the child is retarded by the different causes mentioned.

The head may not be able to enter the brim of the pelvis ; it may become fixed within the brim ; it may pass through the brim, and remain arrested in the cavity ; and, lastly, it may meet no impediment until it is almost expelled, when its advance is opposed just at the outlet. Let us consider each in detail.

The head may not be able to enter the brim of the pelvis.—This may happen when the head is hydrocephalic, or the brim of the pelvis much deformed. In the former case, the moment it is ascertained, as we have already stated to you, the head must be perforated. In the latter, it is of importance to determine the degree of disproportion in the brim, in order to decide on the practicability of delivering the child. In the extreme deformity of either the ovate or cordiform brim, when the antero-posterior measurement is, perhaps, only an inch, or an inch and a half, it would be impossible to do so *per vias naturales* ; * and therefore the Cæsarian section, or what might be called delivery *per vias praternaturales*, becomes a question for your consideration. But the deformity may not be extreme, and yet the head be prevented entering the brim. It is such cases as these that cause the practitioner the most embarrassment, and which present to him the

* In Elizabeth Sherwood's case, delivery was effected by the crotchet, when the antero-posterior measurement was less than an inch. But this exception is so remarkable as rather to prove the rule.

greatest difficulty as to the course he should pursue. It is for the purpose of determining the rule in these cases that our professional mechanicians have contrived an endless variety of pelvi-meters. It is for this object that the profession have made many fruitless efforts, and even entered upon no little controversy, to determine the least possible space through which the child might pass. I shall not occupy your time uselessly with their discussions, but shall only point out to you the errors you must avoid. Recollect that the child must be delivered by destructive instruments, that its sacrifice is involved in the question you have to consider; and this remark applies equally to those who have the temerity to apply the forceps when the head is above the contracted brim of the pelvis. Whether the forceps or the crotchet be employed, the result to the child is the same; the only difference being, that in the former case the hazard to the mother's life is greatly increased. You must not, therefore, place implicit confidence in the accuracy of the measurement that you make of the pelvis, and at once proceed to operate, because it is, or you think it is, within the space through which the head may pass; you might be altogether deceived; and every experienced practitioner knows how often he is deceived in the estimate he forms of the space in the pelvis, although the utmost care may be taken to determine it. Do not, therefore, trust to pelvi-meters, however ingeniously contrived; rather let time and a close attention to the symptoms which present themselves decide your practice. When the character of the labour is changed, when inflammation begins in the passages, when the premonitory symptoms of exhaustion appear, when, after a reasonable time has elapsed (say six hours) without the head making any progress, although the os uteri is quite dilated and the pains have continued regular and powerful—when any of these conditions are met with, and you have reason to think, from an examination of the pelvis, that the brim is too contracted, then, but not until then, are you justified in interfering. The simplest, and we might add, the most efficient pelvi-meter is the hand of the practitioner. If it be the educated hand of the experienced obstetrician, the evidence is certain; but even the attentive student may acquire much accuracy by observing a few simple rules. In ordinary vaginal examinations, when one finger only is introduced, the superior part of the sacrum is quite out of reach. When the point of the finger,

therefore, touches the upper part of the sacrum, it is evidence that the sacrum must be pressed too much forwards; but if it touch the promontory, or just below it, the contraction at the brim must be such as to render it very doubtful if the head can pass. The length of the forefinger is four inches; but from the resistance of the soft parts, as well as of the remaining fingers which are without the vulva, not more than three inches of the forefinger can pass within the pelvis. If, therefore, the distance between the arch of the pubis (taken below the sub-public ligament and soft parts adjoining) and the promontory of the sacrum be only three inches, that between the superior surface of the symphysis and the same point must be less than three inches. You may then take it as a rule, that if your forefinger touch near the promontory of the sacrum, the head will not pass. In more doubtful cases, there are other modes of examination to confirm your suspicions. In some cases, when the cavity and outlet are very wide, the whole hand may be introduced, and the degree to which the fingers are pressed upon, when passed towards the brim of the pelvis, will be your guide. In the standard pelvis, the fingers may be slightly separated from each other. In the pelvis which is contracted at the brim, but through which the head might pass (at least to a certain distance) the fingers are pressed close, but not so as to overlap each other. In greater degrees of disproportion, the fingers cannot be passed forwards without overlapping, and the degree to which this takes place will indicate the amount of contraction in the brim of the pelvis. In extreme cases, three, two, sometimes even one finger will hardly pass between the sacrum and the pubis. In other cases, where this cannot be done, in consequence of the outlet being also distorted, you have, in this circumstance, additional evidence of the character of the pelvis. In some instances, the ischio-pubic rami are very close, and the coccyx abruptly bent; in others, having the same alteration in the coccyx, the symphysis pubis is pressed down towards it. In these varieties it is difficult to make an examination. Only one finger, generally, can be introduced, and if, with this distortion of the outlet, you find the sacrum also pressed forwards, you can hardly hope that the head will pass the brim. These modes of determining the disproportions of the pelvis are far more convenient, and (as it appears to us) are quite as efficient as the more complicated contrivances to which we have referred. But,

with the exception of very extreme cases, we would again caution you not to assume too hastily, even with every evidence in this way obtained, that the head cannot pass the brim, until time and a close observation of the labour removes every doubt.

The head may become fixed within the brim.—This might occur with any of the irregularities of the pelvis that have been described to you. The degree of disproportion may be just sufficient to allow the head to enter the brim, and no more; but as, in the majority of these deviations, the cause retarding its advance exists in the pelvic cavity, such instances may be most conveniently included under the next division. At present, I would direct your attention to one, and only one variety of the deformed pelvis where the head is often fixed in the brim. In the ovate pelvis (that which is produced by rickets) the conjugate measurement of the brim is diminished, while the transverse and oblique are increased, the cavity is shallow, the outlet wide. When the head, therefore, has entered the brim, and is there arrested, the conjugate measurement is the only cause of obstruction; but if this difficulty can be overcome, the head would rapidly advance. Such seems to be precisely the case where the long forceps may be applied with the most success. The shape of the brim is very favourable to its application, and if the head can be drawn through the brim safely, no further difficulty remains. (Vide p. 198.) We would desire to impress this fact the more upon your minds, because it is necessary to discriminate between the application of the



Head fixed in the brim of the pelvis.—(*Smellie*.)

long forceps in this especial case, and its use when the head is in the same situation in other irregularities of the pelvis, and also because this kind of arrest is much more frequently described, and even figured (as you see in the above plate of Smellie's), than those which we believe to be more commonly met with. When the head is thus fixed in the brim, its long axis corresponds exactly to the transverse measurement. The bi-parietal space is compressed between the sacrum and pubis, hence it would be impossible to feel the ear of the child; the sagittal suture is overlapped, but there is always sufficient space between the head and the sides of the pelvis to introduce the blades of the long forceps.

The head may pass through the brim, and remain obstructed in the cavity.—This division embraces the largest number of, and the most important, cases you have to consider. It is to this that the observations apply which we have previously made in reference to the large and ossified head of the male child, the masculine pelvis, the cordiform variety of the deformed pelvis, and the constitutional differences to be noticed in these cases. It is here that the rules of practice are the most contradictory; and however difficult it may be to do so, it is, in such cases, of the utmost importance that they should be, as far as possible, determined. We shall, therefore, first lay before you the characters of this obstruction, both in its slighter form, when it is called *arrest*, and in its increased degree, when it is named *impaction*, and then endeavour to deduce from the facts within our reach a safe conclusion as to the rule of practice.

The term "*arrest*" is applied to those cases where, although the head ceases to advance, the cause either does not depend upon disproportion in the pelvis, or when disproportion exists, it is not so great as to render the delivery of a living child impossible. For instance, when the uterine pains are feeble and inefficient, if the position of the head be unfavourable, or if the arm descend with the head, its progress may be *arrested* without any irregularity in the pelvis. The term "*impaction*" is employed when the head not only ceases to advance, but when there is every evidence that its further progress is beyond the power of the uterus. The use of this term is therefore confined to those cases in which there is great deformity in the pelvis, or to those in which a very large and ossified head is wedged in the deep narrow cavity of the masculine pelvis.

When the head is *arrested* in the pelvic cavity, it may be readily distinguished from the *impacted* head. In the former case, if the head be slightly pushed back, the finger can be passed with facility between the head and the pelvis, the ear may be touched, the parietal bones do not overlap each other strongly, the scalp is only puckered, or, if a tumour be formed on the presenting part, it is diffused, increases slowly, and seldom attains any magnitude. In the latter instance, when the head is *impacted*, it cannot be so easily displaced; it is impossible, without force, to pass the finger between it and the symphysis pubis; the ear cannot be felt, and the urethra is compressed. The parietal bones are strongly overlapped, and cause the sensation to the finger that is expressed by the homely simile of "the sow's back;" a tumour grows very rapidly, and to a great size, often completely obscuring the character of the presentation; the vagina is also swollen and congested. If, however, the death of the child take place, it becomes less in size, softer, crepitant, and œdematous, while the serrated edge of the suture may be felt still more distinctly.

Having given this explanation of the terms which are commonly employed to signify the different degrees of disproportion, and having pointed out to you the characters by which they may be known, we shall proceed, in the next lecture, to consider the management of those cases where the head is delayed in the pelvic cavity.

LECTURE IX.

LABORIOUS LABOURS.

Symptoms of Inflammation—Of Exhaustion—Treatment—Different Degrees of Disproportion in the Pelvic Cavity—Case in which the Head moves very slowly through the Pelvis—Opposite Opinions as to its Management—Discussion of the Question—Attempt to determine it by Statistical Results.

IN the preceding lecture we pointed out to you the manner in which the head may be impeded in the pelvic cavity, and explained the terms used to designate the different degrees of disproportion. We shall now bring before you the effects that result therefrom, those symptoms that indicate danger in labours of this class, and the general treatment required to secure the patient from injury. We shall then endeavour to determine, as far as is possible, the circumstances that justify artificial assistance, and the nature of the aid that is called for. You will recollect, that in speaking of the second stage of natural labour, we directed your attention to the striking change in the action of the uterus when the waters were discharged and the os uteri fully dilated.* The action then becomes much more powerful, and the effort to force the child forward, is assisted by every muscular aid that can be called into action by the patient. Hence the danger that may arise if the advance of the child be opposed by any unusual resistance from the pelvis. If the head be arrested, but still more if it become impacted, the most serious consequences might follow, if not prevented by a judicious treatment. It is therefore necessary, in all such cases, that you watch very closely any alteration in the symptoms that accompany the struggle, in order to foresee the approach of

danger, and to enable you to act with that promptitude and decision which the circumstances require.

The unfavourable results that follow these labours, and, indeed, the majority of all difficult labours, depend generally upon one of two causes—inflammation or exhaustion. Inflammation of the passages may arise, and, if it proceed to any extent, may terminate either in slough of *some part* of the vagina (more especially the vesico-vaginal septum, because most exposed to pressure), or in slough of the *whole vagina*, if the inflammation assume any erysipelatous character, or perhaps in *slough and laceration* of the portion of the vagina connected with the neck of the uterus, thus leading to *rupture of the uterus* itself. Each of these consequences, although differing in degree, is equally to be avoided. It is therefore important to observe the earliest symptoms of inflammation.

Another cause of danger is *exhaustion* of the uterus. When the uterus becomes powerless, artificial assistance is absolutely required to deliver the child. The danger, however, does not consist in this necessity, but rather in the cause that led to it. Atony of the uterus is not easily produced; and when it occurs, the shock of the constitution is very great. Sudden death has been the consequence of it; and the patient is always placed in imminent danger, because of the symptoms of constitutional disturbance which present themselves. Beside this, the uterus having lost to a certain extent its powers of contraction, hæmorrhage may be the result; and this increases the exhaustion of the patient, and, consequently, the danger of the case.

Hence, one of the most serious effects that can follow prolonged labour is, that the patient becomes, in the *true meaning* of the term, exhausted. It is one which should be guarded against by every precaution, and one which justifies the most prompt interference the moment any indication is given of such an unlooked-for alteration in the constitution of the patient. But do not confound exhaustion with mere fatigue. Fatigue is the ordinary effect of long-continued exertion, and, therefore, of long and severe labour; exhaustion, a very rare result of it; nevertheless, the term exhaustion is applied indifferently to express both effects. The patient will tell you that “she is quite *exhausted*.” Her friends will exclaim, that “she will sink from *exhaustion* if not relieved.” This language is well understood

and appreciated by the intelligent practitioner in cases where there is no exhaustion whatever; but we fear we must add that it is an appeal very readily listened to by some who are not quite so prudent, and who feel quite as much inconvenience from fatigue as the patient. You must not be betrayed into impatience from such a cause, but rather steadily observe the character of the labour, and be prepared to recognise the earliest symptom of exhaustion the moment it presents itself. Thus you will be able to interfere with promptitude when your assistance is required, and, on the other hand, will not be hurried by these urgent solicitations into an unnecessary, and perhaps an injudicious, attempt to terminate the labour.

When the head of the child moves very slowly, or ceases to advance through the pelvic cavity, the second stage of labour is often greatly lengthened. In such instances, the uterus continues to act for a certain time with the same power as before; but if no effect be produced, its action is suspended in the same manner as in the first stage, only for a shorter time. When the pains are renewed, they are not so strong as before; the uterus seems, as it were, conscious of the difficulty; the pains are shorter, although very often severe. At this time the woman is disposed to use every effort to force the child forward, and, impatient of the delay, will strain with all her strength: failing in her object, she becomes alarmed and dispirited, and her mental anxiety still further interrupts the action of the uterus. She should therefore be dissuaded from fruitless attempts of this kind, because nothing is so well calculated to excite inflammation, if not to produce exhaustion. It is the more necessary to point out this to you, because the patient is too frequently recommended to adopt this very practice. She is often desired "to bear down with her pains," and "to assist herself" at a time when she can give herself no assistance whatever.

After this comparatively feeble action of the uterus has continued for some time, the pains return in their former strength, especially if the patient has had even momentary sleep. If the difficulty be still insuperable, they may either again cease, or continuing, may produce inflammation of the passages. When this is in the least degree indicated, the pains are very short, extremely severe, and in their intervals the patient still complains of pain, and a feeling of soreness. If the uterus be ex-

amined through the abdomen, you will observe a very perceptible difference in the sensation it communicates. It feels almost as hard and contracted in the interval as during the pain; the patient cannot bear to have the abdomen touched. Beside this alteration in the character of the pains, we have other symptoms, both local and general, to guide us. Febrile irritation, which had been previously absent, shows itself; the pulse becomes quick, frequent, and not easily compressed; the tongue is dry, and the patient has great thirst; the countenance is anxious, and the features slightly collapsed; there may be a distinct rigor, but more usually there is but a slight chill. Locally, the vagina becomes hot and tender, the mucous secretion ceases; it is therefore dry, or, if the inflammation increase, a serous discharge may be substituted. When inflammation thus commencing is neglected, and it extends to the uterus, there are further local changes depending upon the effects produced on the contents of the uterus. The membranes undergo decomposition, and then a thick, yellow, oily, and very offensive discharge flows from the vagina. This is increased if, the death of the child having taken place, its tissues undergo a similar change. You should bear in mind, however, that there are cases where the death of the child may have taken place either in the beginning of labour, or before it has commenced. In such instances, putrescency may take place rapidly, and similar appearances to those just mentioned might present themselves without any inflammation having existed; you must not, therefore, confound the dark and fœtid discharges from the vagina, that depend upon such a cause, with those that occur in connexion with inflammation of the vagina and uterus.

When the head remains in the cavity of the pelvis, and the least tendency to inflammation is manifested, the most prompt and decided measures are necessary. In strong and plethoric habits, free general depletion, followed by nauseating doses of tartar emetic, emollient enemata, and local fomentations, will check the advance of inflammatory symptoms, and give time to determine the important question as to the delivery of the child. In those feeble constitutions where you have reason to dread lest exhaustion should ultimately take place, you cannot employ general depletion, nor any means that would reduce the strength of the patient to any extent. It is preferable to

deplete locally, to use fomentations, enemata, and, after the bowels are evacuated, to give a *moderate* dose of liquor opii sed., combined with some diaphoretic. Inflammation does not so readily arise in these cases, and when it occurs it is more easily arrested; but exhaustion would be much more likely to follow prolonged labour than in those more robust constitutions we have alluded to. Both cases demand from you an equal attention; but you perceive it must be directed to a different object. We shall therefore briefly detail to you those premonitory symptoms that indicate the approach of exhaustion. When the uterus is becoming weak, and its strength is, in the strictest sense of the term, almost exhausted, the pains become short and inefficient, the intervals between them longer, and sometimes they are quite suspended; but the nervous system being excited, the patient derives no advantage from such suspension. She does not sleep, but continues watchful, anxious, and restless; the pulse is increased in frequency, is very easily compressed, and its pulsations are disturbed and rendered irregular by the most trifling causes. The countenance of the patient is peculiar, to say that it is collapsed would not convey its exact expression. Watchfulness and anxiety are portrayed on features that equally indicate languor and listlessness. An observable change takes place, but to explain how is not easy. If these monitors be neglected, further symptoms of constitutional disturbance will soon present themselves. Constant watching, increased restlessness, hurried respiration, irregular chills, and slight delirium, all rapidly succeed each other, and hasten to the most unfavourable conclusion. When there is the least reason to apprehend exhaustion, promptitude in delivery is imperative; but before interference is thus called for, much may be done to prevent its necessity. Rest is very important in these cases, because it is an evidence that the over-excited nervous system is tranquillized; therefore, when the pains are becoming weak, or return only at long intervals, a moderate dose of opium is often of great service: if the patient sleep, even for a short time, the uterine action is renewed with much more power. I have expressly stated a *moderate* dose of opium, because the case now under our consideration is that in which the same medicine has produced the most opposite effects, according to the dose administered, and, consequently, it has sometimes

been discarded, because it has been misapplied. The object which it is intended to accomplish by opium, is to subdue the nervous irritation which precedes exhaustion, and to restore, by rest, the energy of the uterus. The effect produced by too large a dose may be to paralyze all nervous power, and thus at once to cause uterine exhaustion. You must therefore exercise proper caution in the use of it; twenty to twenty-five minims is generally sufficient for the purpose. After a temporary rest has been thus produced, if the uterus still continues to act feebly, ergot of rye may be given in an equally cautious manner, carefully attending to its influence on the pulse, and especially on the circulation of the fœtus. If in either case, after giving this medicine, the rate of the pulsations be diminished, you must not persevere in its employment, otherwise the death of the child may be the result.* It is also necessary to be

* Ergot of rye should be given with great caution. Its effect has been closely observed by Dr. Beatty, and he is led to the following conclusions:—

“That the administration of ergot to a woman in labour is attended with danger to the child, whenever a time sufficient for the absorption and transmission of its noxious properties elapses before the child is born. The degree of effect produced, differs with the time that elapses between the exhibition of the dose and the birth of the child. Hence, the ergot should never be given in any case where there is a likelihood of the labour lasting more than *two hours* after its administration; except when it may be employed to save the mother's life; and secondly, if delivery is delayed to two hours, we should resort to artificial assistance to save the life of the child.”—*Dublin Journal*.

Dr. Hardy has since made its effects the subject of his observations, and has drawn up a series of Tables to determine—

- 1st. The period after administration that its action commences.
- 2d. Its effects on the maternal pulse (when any), and how soon evident.
- 3d. Its effects upon the fœtal heart (when any), and how soon produced.
- 4th. The state of the uterus and lochial discharge during convalescence, in cases in which it has been exhibited.

1. It appears that, in some cases, ergot of rye acts on the uterus so soon as seven minutes after its exhibition, whilst in others a much longer period of time is required; from about ten to fifteen minutes may be stated as the average. In those cases where the children have been expelled alive, I have always observed the action of the uterus to commence within twenty-five minutes. On the other hand, when a longer period than this elapses before the uterus takes on action, the use of instruments has been necessary to perfect the delivery, or the children have been dead-born.

2. In nineteen cases there was a marked diminution in the frequency of the mother's pulse, following the administration of ergot. This effect generally began to take place from about fifteen minutes to half an hour. In all these instances,

careful to avoid the use of *secale cornutum*, if the delay in this stage arises from great disproportion between the head and the pelvis. It must be obvious to you, that in a case like this it would be very dangerous to use a means of exciting the action of the uterus, over which you can have no control. A preparation which exerts a specific influence on the uterus, which often causes the most violent action, and that not returning at intervals, as ordinary pains do, but which excites a *continuous* effort of the uterus to expel the child, is not the safest to employ

when the depression of the pulse occurred, the fœtal heart underwent a similar change. In several cases where the circulation of the patient underwent this depression from the action of the ergot, the effect continued several days.

3. In the majority of cases, a diminution of the fœtal heart's pulsations followed the exhibition of ergot. The period at which this effect begins to be produced, varies from fifteen minutes to half an hour. The most common effect is a diminution in the frequency of the pulsations: this is succeeded after some time by an irregularity in its beats, which irregularity continues, more or less, until the sounds intermit, and at length after a variable period become quite inaudible. In those cases where the number of pulsations have been steadily reduced below 110, and *at the same time with intermissions*, the child will rarely if ever be saved, although its delivery should be effected with the greatest possible speed.

4. The volume of the uterus is often found much greater than after ordinary labours. In addition to this enlarged state of the uterus, it has sometimes a firmly contracted feel, which generally continues for several days. In a few instances the lochial was rather pale and scanty, although nothing unfavourable occurred during convalescence to account for this circumstance. With some few exceptions, the women had generally good recoveries. The children that were born alive, all, with one exception, did well. In this case, delivery was effected by the forceps, as the fœtal heart had fallen so low as 100 from the exhibition of the ergot. It died in three hours after delivery.

General results of Dr. Hardy's Tables.

Cases in which the uterus expelled the child alive	7
Cases in which the children were born alive by the application of the forceps or vectis (after ergot was given)	7
	—14
Cases where the uterus expelled the child dead-born (after ergot)	15
Cases in which the forceps or vectis were applied (after ergot), but the children were dead-born	6
The children dead (from ergot), delivery effected by the crotchet	13
	—24
	—

Dublin Journal, vol. xxvii. p. 224.

43

From the above summary of cases given by Dr. Hardy, it appears that in forty-eight cases where ergot of rye had been given, thirty-four children were still-born,—nearly three-fourths!

when there is much resistance opposed to this action. The remedy, when cautiously administered, is useful, however, in those cases where the delay chiefly arises from want of power in the uterus, which may be exhausted if not thus artificially stimulated to action.

Having given you this brief outline of the general treatment necessary to these protracted cases, we shall now enter upon the more difficult question of their management, to secure the delivery of the child, assuming that instrumental aid is not rendered imperative by the presence of inflammation or exhaustion. It is hardly necessary to state to you that there is every variety in the degree of disproportion between the head and the pelvis. In some instances it is so slight that the child may be safely delivered without any assistance; only it will occupy a longer time in passing through the pelvis. In others, the amount of difficulty may be so much increased as to render it doubtful whether the head can pass without assistance; and it is in these cases that the rules which are given for your guidance are the most contradictory. Again: you may have a still greater disproportion, in which there is no doubt about the improbability that the head can be expelled by the natural efforts of the uterus, although there is very great doubt, and no little dispute, as to the means by which the head must be extracted. Lastly: you have occasional instances in which the narrowness of the pelvis is such, or the magnitude of its distortion is so great, that the safe delivery of the child is hopeless: the head must be lessened; it must be destroyed before it can be brought into the world. In extreme cases of this kind, even this cannot be done; but recourse must be had to the difficult and dangerous operation of removing the child from the uterus by laying it open, in order to save the mother from the dreadful alternative of dying undelivered.

In those cases of slight deviation from the standard pelvis, where there is every evidence of space sufficient for the head ultimately to pass through the pelvis, if nature be allowed time for the purpose, you would not, of course, interfere with her; although I believe instances might be quoted where *very adroit operators* have, even in such cases, relieved the tedium of a long attendance by a ready application of the forceps. It is sufficient to say, that the united testimony of the profession, given in

every standard work of midwifery, is opposed to such a practice; and if any accident should arise from this mischievous meddling, the operator is fully responsible for all the consequences that follow from it. But in those more doubtful cases, in which there seems hardly sufficient space for the head to pass safely through the pelvis, the practice is not so clear, nor is the evidence of the profession so unanimous on the subject. When, in such instances, the head is actually arrested, and so remains for some hours in the same position—a sufficient length of time to satisfy you that the uterus cannot advance it—if the ear can be felt, or the finger be passed easily between the head and the pubis, you may use the forceps to deliver the child, and I think the weight of authority will support your practice. But when the head is not so arrested, but, at the same time, advances so extremely slowly that it seems to be arrested, you have here the discordance of authorities at once confounding you. Burns devotes the greater part of a chapter* to prove the impropriety of delay under such circumstances, and advocates the application of the forceps in cases of arrest, or rather of slow progress of the head, because “in such cases then we may experience much evil from trusting too long to nature, but add little to the sufferings of the patient, and nothing to her hazard, by instrumental aid.† He applies the same principle to cases of impaction, which we shall presently consider. The evil he dreads is uterine exhaustion, if this second stage be much prolonged. In this view he has the support of the late Professor Hamilton, who equally dreaded delay. Dr. Campbell also gives a similar opinion, but more guardedly expressed. “It may, however, be repeated, that while the delivery is advancing, and the patient continues free from unfavourable symptoms, the use of the forceps is to be abstained from altogether. *But whenever the progress is slow and imperceptible, and the subordinate means already recommended have failed to accelerate the transit of the fœtus, the case should be watched, and this instrument applied with very little delay after the passages are prepared.*”‡ Other names might be added to this list of advocates for interference in the case

* Burns, chap. vi. p. 428.

† Ibid., p. 434.

‡ Campbell's Midwifery, p. 281.

supposed. But let us turn to the other side, and you will find the eminent names of William Hunter, Osborne, and Denman, opposed to this practice. Dr. Osborne would wait until exhaustion had actually taken place—a maxim for which he has been very severely, and I admit very justly, criticized. Dr. Denman's fifth aphorism states, "It is meant, when the forceps are used, to supply with them the insufficiency or want of labour pains; but *so long as the pains continue*, we have reason to hope they will produce their effect, and *shall be justified in waiting*."* When the pains cause the head to advance, although very slowly, they are producing their effect; and the case therefore comes within the limits of the aphorism. Dr. Collins observes, "Let it be carefully recollected at the same time, that so long as the head advances EVER SO SLOWLY, the patient's pulse continues good, abdomen free from pain on pressure, and no obstruction to the removal of urine, interference should not be attempted, *unless the child be dead*."† Dr. F. Ramsbotham's third rule on this point is, "If the head advance at all, and be not impacted, provided the strength and spirits are good, there is seldom need to interfere."‡ Dr. R. Lee's name might also, I think, be added, as being favourable to this rule of practice.

I shall not, gentlemen, so far trespass on your patience as to ask you to unravel with me this tangled web of contradictory experience. It is sufficient if I convince you of the difficulty of the subject, and if it induce you to give a patient attention to the only mode that I can adopt to draw a legitimate conclusion—that is, to derive it as nearly as possible from facts, without reference to opinions. I think this may be done. Bearing in mind that the great and leading principle to be observed in these difficult cases is, to preserve both mother and child, if possible, from injury, I think it is in our power to compare the results of cases where the forceps has been applied with those where it has been withheld, and thus determine the practice which presents the greatest success. We shall first direct your attention to the following tables of operative midwifery, derived from reports given by British and foreign practitioners;

* Denman's Aphorisms, p. 12.

† Collins's Report, p. 18.

‡ Ramsbotham's Obstetric Medicine and Surgery, p. 309.

you will find in them the total number of cases given in each, the number of forceps operations, and the results to mother and child when they are given.

COMPARATIVE VIEW OF FORCEPS OPERATIONS AND PERFORATIONS.

BRITISH REPORTS.

Date.	Place.	Name.	Total Cases	For- ceps	Deaths.		Perforations.	D'ths: Moths	Total Opera- tions.
					Chil'n	Moths			
1781	Lond'n	Dr. R. Bland . .	1,897	4	—	—	8	—	12
—	Ditto	Dr. Merriman . .	2,947	21	6	—	9	—	30
1828 to 1843	Ditto	Dr. F. Ramsbotham	35,745	49	11	3	38	6	87
1787 to 1793	Dublin	Dr. J. Clarke . .	10,387	14	—	2	49	16	63
1826 to 1833	Ditto	Dr. Collins . . .	16,414	24	8	4	79	15	103
1835 to 1837	Ditto	Dr. Beatty . . .	1,182	9	4	—	3	—	12
1835 to 1840	Ditto	Dr. Churchill . .	1,640	3	1	—	12	1	15
1832 to 1835	Ditto	Dr. Murphy . . .	5,699	14	5	1	29	6	43
			75,911	138	35	10	227	44	365
	Lond'n	Dr. R. Lee . . .	—	55	38	9	127	23	182

FOREIGN REPORTS.

FRENCH.

{Date.	Place.	Name.	Total Cases.	For- ceps	Deaths.		Perforations.	D'ths: Moths.	Total Operations.
					Chil'n.	Moths.			
1797 to 1811	Paris . .	Bolvin . .	20,357	96	23	—	16	—	112
1812 to 1820	Ditto . .	La Chapelle .	22,243	77	18	—	12	—	89
			42,600	173	41	—	28	—	201

GERMAN.

1821 to 1825	Wurtem- burg .	Riecke .	221,923	2,740	636	127	98	35	2,838
1801 to 1821	Vienna .	Boer . .	26,965	100	—	—	43	—	143
1797 to 1827	Ghent .	Jansen .	13,365	341	—	—	5	—	346
1811 to 1827	Prague .	Moschner	12,329	120	—	—	4	1	124
1825 to 1827	Bonn . .	Kilian . .	9,392	120	—	—	4	—	124
1814 to 1827	Dresden	Carus . .	2,549	184	—	—	9	—	193
1817 to 1828	Berlin .	E. Siebold	2,093	300	—	—	1	—	301
1823 to 1827	Ditto . .	Klugè . .	1,111	68	14	—	8	3	76
	Heidel- berg .	Naegele .	1,711	55	—	—	5	—	60
			291,438	4,028	650	127	177	39	4,205

You will perceive that in these tables the number of forceps

operations in British practice is 138; in thirty-five of which the child was still-born, being in the proportion of one in every fourth case. In order to prevent error in this proportion, we have separated Dr. Lee's forceps cases; the total number of which given by him is fifty-five; the mortality of children, thirty-eight; which would be quite out of proportion (being more than one-half) if these cases were not carefully examined. I have endeavoured to do so, and to make the necessary corrections. In nineteen of these fifty-five cases, the forceps failed: they therefore became cases for perforation; of the remaining thirty-six cases, one-half the children, eighteen, were lost, but twelve of these eighteen were destroyed by other causes than the forceps. Deducting, therefore, all such cases from the whole number, the remainder will be twenty-four forceps cases, in which eighteen children were saved and six lost, being in the same proportion, one in four.

In the French reports, forty-one children were lost in 173 forceps operations, being one in four, nearly.

In the German reports of Riecke and Klugè, which state the mortality of the children, the number of the forceps operations united is 2,808; the deaths of children, 650; being also one in four, nearly. Thus, then, we may conclude, that one-fourth of the children delivered by the forceps are lost. What is the result when these protracted cases are left to themselves? Is the mortality increased? I do not think such will be found to be the case. In order to determine this question, I must refer you to Dr. Collins's valuable report—the only report which, from its extreme accuracy and minuteness, affords the elements upon which to form a calculation. Dr. Collins has given tables to show the duration of labour in all the cases he reports; he has also given separate tables to show the duration of labour in forceps cases, and in those which are preternatural. We may also assume, that perforation being only had recourse to “when, after the most patient trial, the impracticability of labour being terminated in safety by any other means was clearly proved,”* that all these cases exceeded twenty-four hours. From these data, then, we shall endeavour to draw a fair conclusion.

* Collins, p. 23.

*Cases of labour protracted to 24 hours and upwards from Dr.
Collins's Report,*

TOTAL CASES, 430.		STILL-BORN CHILDREN, 150.		MOTHERS DEAD, 40.	
Delivered by forceps	12	Still-born	4	Mothers Dead . .	0
by perforating	79	Ditto	79	Ditto	15
preternaturally	15	Ditto	6	Ditto	0
naturally	324	Ditto	61	Ditto	25
	430		150		40

From this table you perceive, that of 430 cases in which labour lasted twenty-four hours or exceeded it, 324 of them were natural cases, delivered without assistance, and that of these 324 the children were lost in sixty-one instances, which would be about one in five cases. The result of my own inquiries on this subject is nearly similar, and has been obtained from the same source, the Dublin Lying-in Hospital.

Report of 218 cases of labour protracted to or beyond 24 hours.

Delivered.	Cases	Boys.			Girls.			Mothers Dead.	Causes of Mother's Death.
		Liv- ing.	Dead	Putrid.	Liv- ing.	Dead	Putrid.		
By forceps	14	4	1	—	5	4	—	4	{ 1 Puerperal fever. 1 Rupture of uterus. 1 Puerperal fever. 1 Rupture of uterus.
By perforations	29	—	20	1	—	7	1	6	
Naturally	175	76	19	5	52	22	1	8	
	218	80	40	6	57	33	2	18	3 Puerperal fever.

In 5,699 cases, 218 were protracted to this degree; and of these, 175 were delivered naturally, and forty-one children not putrid were still-born, being one in four, nearly. Thus, then, you perceive that, taking the widest, and, we would say, the fairest view of this question, the proportion of still-born children in these difficult and protracted cases is nearly the same, whether the forceps be employed or otherwise; that the difference, if any exist, in favour of Dr. Collins's practice of leaving these cases to nature. But this is only one view of the question.

It may be said, and has been said, in the energetic language of Dr. Burns, that the mother must be considered. "From the strength of the recommendations of the partizans of nature, we should suppose that whenever the child could actually be born without aid, no hazard occurred; and, on the other hand that instruments must of necessity prove not only very painful in their

application, but dangerous in their effects. Now, the first supposition is notoriously wrong, for *innumerable instances* are met with, where the mother does bear her child without artificial aid, and much, doubtless, to the temporary exultations of the practitioner, but, nevertheless, death takes place, or at the best, a tedious recovery is the consequence.”* Is such the case? It is totally opposed to my personal experience; on the contrary, I have been surprised at *the rapid* recovery of patients who have suffered this protraction, when I had erroneously anticipated, from that very circumstance, all the unpleasant consequences here detailed. But I would again ask you to put aside, for the present, individual experience, and examine the facts. In doing so, our data are more limited than those which assisted us in the former question, because, in the French reports, there is a most ominous silence regarding the mortality of the mothers—they say nothing about it. In the German reports, we are limited to that of Dr. Riecke, who gives 127 deaths in 2,740 cases, being one in twenty-one, nearly. But take Dr. Churchill’s more extensive researches on this question, from whose valuable work on operative midwifery these tables of foreign practice are partly formed. He states, that “amongst the French and Germans, in 479 cases, thirty-five mothers were lost, or about one in thirteen.† Dr. Churchill gives the proportionate mortality in British practice as one in twenty-one; but you perceive that, in the comparative view we have placed before you, there were ten deaths in 138 cases, which is about one in thirteen. Compare this with the result where the cases have been left to the natural efforts. In Dr. Collins’s report, there were twenty-five deaths in 324 cases, or one in thirteen, precisely the same as where the forceps had been used. Among those cases which I have observed, there were eight deaths in 175 cases, or one in twenty-two—a proportion in which I can place the more confidence, because it is derived from personal observation.

With regard, then, to the second question, the mortality of the mother, take the estimate in any way you please, and you must arrive at the same conclusion—viz., that the mortality is certainly not increased when these cases are not interfered with,

* Burns, p. 434.

† Churchill’s Operative Midwifery, p. 134.

and all the dreaded consequences which Dr. Burns anticipates from such practice have no foundation in fact. But we might even go farther: we might say, that so far from such evils following our Fabian practice, the evidence seems to point the other way, and to prove that the actual mortality is diminished. The twenty-five deaths reported by Dr. Collins include cases of puerperal fever, and other causes of death which might be called accidental, because he gives, under a distinct head, the number of deaths, the "effects of tedious and difficult labours."* These are just eleven cases, or one in thirty cases, nearly. The eight deaths which took place under my own observation, include three deaths from puerperal fever, leaving only five deaths from the severity and protraction of labour, which would be in the proportion of one to thirty-seven cases. Caution, however, is necessary, when we would derive a just conclusion from statistics. It is therefore possible, that if the reports of these forceps operations were more fully given, so as to separate the deaths from accidental causes from those resulting from the operation, the proportion of mortality would be diminished in the same ratio.† We do not wish you to assume more than what we think has been proved—viz., that the mortality of the mothers is not increased by leaving these cases to nature. The safety of the mother or child cannot, therefore, be advanced as a reason for instrumental delivery, when the head is making a very slow but a certain progress.

One argument, however, has been much used by the advocates for interference, which is very clearly expressed by Dr. Burns: "Granting (he observes) the recovery to be excellent, is it no consideration that the patient has been subject to twelve, perhaps twenty-four, hours of suffering of body and anxiety of mind, which might have been spared?"‡ You must perceive that if this argument be worth anything, it will admit of a much more extended application than Dr. Burns would give to it. It might be employed to justify the use of the forceps in every case where the head was within reach, and labour at all severe. Because,

* Collins, p. 365.

† It is probable that the reports include deaths from puerperal fever, and those following delivery of the *impacted* head by the forceps.

‡ Burns, p. 434.

why should your patient be exposed to *any* bodily suffering or anxiety of mind, if it were in your power to relieve her from her miseries? On this principle, the forceps might be used (as indeed they have been used) in every tenth case, and the practitioner relieved from the most anxious portion of his duties. But the design of nature will not thus be thwarted; and we might reply to such an argument in the language of Naegele—"If we admit that proportionate difficulties, according to the constitution of each individual, and an effort of strength (requisite in child-birth), are inseparable from the nature of this process, we must conclude *that an abbreviation of this process, though performed by an able hand, before the salutary change, on which the preservation of health depends, has taken place in the organization of the mother, that a premature and sudden removal of these difficulties cannot be a matter of indifference; that such a violent interference with the functions of nature must incur the risk of destroying the health, though this should not ensue for some time after.*"* A more immediate injury, however, sometimes follows the application of the forceps in the case we are supposing, as well as in cases of impaction, which we shall have again to consider. The pressure of the instrument may cause slough of the neck of the bladder or the urethra, and thus establish a fistulous opening into the vagina; and the incontinence of urine that follows renders the patient's life miserable afterwards. It is difficult, in all instances, to trace this accident to the use of the forceps. When a forceps operation is described to us, we are seldom told that any mischief is the consequence. The splendour of success is very dazzling, and while we admire the operation, we are too often left in the dark as to the effects. Nevertheless, I have been able to trace this accident clearly to the use of the forceps in several instances. The usual account given by the patient is, "that she had been delivered by instruments, and the child's life saved." Dr. R. Lee, in his "Clinical Reports," gives a candid and clear statement of the results in the forceps cases he details. "Four died from the rash and inconsiderate use of the forceps; seven had the perinæum more or less injured; one had the recto-vaginal septum torn; five were left with cicatrices of the vagina, after sloughing; and one with

* Naegele's Mechanism of Parturition, by Rigby, p. 88.

incurable vesico-vaginal fistula.”* Dr. Collins records only *one* case of vesico-vaginal fistula in the whole of his report of 16,654 cases—that was a case of perforation—consequently this accident never was found among those cases which were delivered naturally. The only case of fistula which occurred in the 5,699 cases to which I have so often referred, was one in which I employed the forceps to deliver a child that presented the forehead. The principal cause of difficulty in Dr. Collins’s cases, was the large head of the male child forcing its way through a very osseous pelvis; the pressure on the soft parts must be very great, and if fistula could be produced by great protraction of labour in cases that ultimately were delivered without assistance, it must have been an accident of frequent occurrence in these cases, when the soft parts were so much compressed; but such did not happen, and therefore they afford a very favourable contrast to the cases delivered by the forceps in nearly similar circumstances. The intelligent practitioner would therefore hesitate to expose his patient to the risk of vesico-vaginal fistula, for the mere gratification of shortening the severities of labour.

We have been reluctantly compelled to dwell longer upon the management of this degree of disproportion than we desired. The difficulty of the question it involves, and the contradictions among the most experienced writers, must be our apology. In the case that we are considering, that in which the second stage of labour is protracted, and the head of the child advancing *very slowly*, we have shown you that there is no increased danger to the mother or child by leaving the case to nature in place of delivering by the forceps; that if there be any difference in the ratios of mortality, it is in favour of non-interference, and rather against the forceps. We have pointed out, as far as imperfectly detailed facts would enable us, that the post-partum accidents of labour follow operations with the forceps more frequently than cases which are left to themselves, and, consequently, the conclusion at which we must arrive, is hostile to the use of that instrument, under the circumstances stated. But recollect, that there is no general rule without an exception, and you will sometimes meet with cases so feeble in their habits that they will not endure a protracted labour without great risk of exhaus-

* Clinical Midwifery, p. 32.

tion; you may be called to patients where you dare not temporise, whom you must deliver although the head is making a tardy progress. We only ask you to consider these as the exceptions, not often met with, but still necessary to be studied and understood. It is for this reason we have brought before your notice the symptoms of exhaustion, and those which precede it; the same desire to direct your attention to the study of individual cases which may be exceptions to the general principle, we would wish to govern you, leads me to bring before you the varieties, not only in the formation, but in the resistance of the pelvis, so that you may know where an operation might be undertaken and where it cannot be attempted. If we have placed this subject before you with sufficient clearness, we shall conclude by directing your attention to that opposition in the practice of experienced authorities which has rendered its discussion so necessary.

Name.	Total Cases.	Forceps.	Proportion, 1 in	
Ramsbotham . .	35,745	49	729½	London
Clarke	10,387	14	742, nearly	Dublin
Collins	16,414	24	684, nearly	Dublin
Boivin }	42,600	173	246	Paris
Lachapelle . . }				
Boër	26,965	100	269½	Vienna
Kilian	9,392	120	78	
Carus	2,549	184	14	Dresden
Siebold	2,693	300	9	Berlin

Ramsbotham, one forceps operation in 700; Siebold, one in nine cases!

LECTURE X.

LABORIOUS LABOURS.

Management of Cases when the Head of the Child is impacted—Opposite Opinions as to the Application of the Forceps—Examination of the Question—Forceps as an Instrument of Compression—Baudeloque's Experiments—Perforation—Advantages of Auscultation—Case when the Head is retained at the Outlet—Accidental Obstructions—Ovarian Tumours—Polypus—Fibrous Tumours—Osteo-sarcoma.

THE object of our attention in the preceding lecture was to determine the rule of practice in those cases of laborious labour where the head advanced *very slowly* and with much difficulty, through the pelvic cavity. The conclusion at which we arrived was opposed to interference under such circumstances. That the *general rule* should be to leave the case to nature, but at the same time to recollect that there are *some exceptions* to this rule, depending upon constitutional or accidental causes, where instrumental aid becomes necessary. We shall now examine the next degree of disproportion, that by which *the head is impacted in the cavity of the pelvis*; a case where the ear cannot be felt, where the soft parts suffer very great pressure, and where it is hopeless to expect the delivery of the child by the unassisted efforts of the uterus.

The management of cases where the head of the child becomes impacted has been, I regret to say, almost as much a question for controversy as that which we have just discussed. It is admitted that the child must be delivered by the resources of art, but how these resources are to be applied is the matter in dispute. Some consider that even in these cases the forceps, skilfully employed, may effect the object in view; the woman may be thus delivered, and possibly the child preserved. Others dread such application of the instrument, because of the

injury that may be done to the passages, and consequently they esteem the probable danger to the mother to be a risk too great to encounter for the very slight chance of saving the child. Hence the question lies between perforation of the head of the child and its forcible extraction by the forceps.

It would be most desirable to determine the rule of practice in these very difficult cases, by an application of the same principle that was proposed to you in the last lecture. If we could compare such cases as have been delivered by the forceps when the head was impacted, with those in which recourse was had to perforation—if we could contrast the results—we might be able to arrive at a conclusion that would satisfactorily resolve our doubts upon the subject; but, unfortunately, that is impossible. We have no statistical knowledge of the effect of the forceps in these special cases; and the mortality that is reported under the head of perforation seems to be disproportionately increased by the circumstances under which the operation has been generally performed.

From the earliest period, the profession have been accustomed to look upon craniotomy with dread—I might almost say, with horror. A natural reluctance to destroy human life—no matter under what necessity—has been greatly increased in some countries by religious prejudices; and the anathema of the doctors of the Sorbonne still exerts an influence that paralyzes the judgment of the practitioner. Hence we read of cases allowed to remain several days in labour, until not only the death, but the putrefaction, of the child, gave evidence that the perforator might be employed without any stings of conscience. The result of such practice was, as might be supposed, inflammation of the passages, advanced to such an extent that the mother was sacrificed to this procrastination; and hence in the tables of mortality, we find that *one mother in every five*, and sometimes *one in every four*, died after the operation. We cannot, therefore, determine the rule of practice by statistical returns. We must only hope to do so by a fair examination of the question itself, by collecting the general experience of the profession, and by submitting to you the ground upon which we have formed the opinion which would govern us as to the course to pursue. We do not wish you to adopt this opinion

unless you are satisfied of its correctness; we but ask you to examine the subject dispassionately, and to discard from your minds the damnatory language that too frequently is employed by some obstetric authors. When you find an operation spoken of as "murderous," you are not disposed to become the murderers; you doubt and hesitate, and perhaps ultimately commit a *double homicide*. An appeal to harsh expressions is generally esteemed an evidence of weakness in argument; therefore, when you find these hard words, you can appreciate their value, and pass them by for more conclusive reasoning.

In order to compare the forceps and perforator in the case supposed, you must view the forceps as something more than a substitute for the power of the uterus. In order to extract the head, it must also lessen its dimensions; it must be employed for the purpose of *compression* as well as for *extraction*. When we come to examine the different instruments used, you will find, in the varieties of the forceps, that some are shaped especially for this purpose, which is sedulously avoided in the construction of others—a sufficient proof of want of unanimity on this important subject. Let us, then, examine the forceps as an instrument for compressing the head of the child, so as to adapt it to the diminished space in the pelvis.

We have already evidence before us to prove that the power of the forceps for this purpose is extremely limited. The experiments of Baudeloque are referred to in almost every popular work on midwifery, and notwithstanding the critical objections raised against them, they are sufficiently important briefly to state them to you. Being desirous to determine the extent to which the forceps could compress the head, Baudeloque performed nine experiments on the heads of still-born children with Levret's forceps, an instrument of the strongest kind, and especially adapted for compression. The utmost force was exerted to reduce the head, a force so great as to bend one forceps, although highly tempered; the head was not lessened more than two lines, unless where the bones were unusually soft and loose, and then only to four lines. These experiments satisfied Baudeloque that the diminution could not be, in any case, so much as accoucheurs had stated, and that the degree of reduction should never be measured by the distance between the handles,

when pressed together, nor from the amount of force employed to approximate them.*

In these experiments more force was used than you could ven-

* 1627. Ces expériences ont été répétées de suite sur neuf enfans morts à l'instant de leur naissance, ou peu d'heures après, qui étoient d'une grosseur différente, quoique tous parfaitement à terme. Pour les rendre plus concluantes, nous fîmes en sorte de restituer à la tête de ces enfans, en la plongeant dans l'eau chaude et en la pétrissant un peu avec les mains, la souplesse que présente au toucher celle des enfans vivans, et nous nous servîmes du forceps allongé dont il est parlé au § 1614. Nous nous en procurâmes trois semblables, *de la meilleure construction et de la meilleure trempe*. Nous appliquâmes cet instrument d'abord selon l'épaisseur transversale de la tête, comme nous le recommandons ailleurs; et ensuite suivant la longueur du crâne, c'est-à-dire une branche sur le milieu du front en descendant de la fontanelle à la racine du nez et l'autre sur l'occiput, pour connaître la réduction qu'on pouvoit opérer dans ces deux directions, et ce que la tête acqueroit dans un sens, en perdant selon l'autre. Quelque soit le degré d'écartement que laissèrent entre elles les branches du forceps à l'extrémité qui se termine en crochet, toutes les fois qu'elles furent placées sur les côtés de la tête nous les rapprochâmes exactement et nous les fixâmes dans cet état de contact au moyen d'un ruban, pour que la réduction de la tête ne variât point pendant que nous mesurerions de nouveau ces dimensions, et que nous les comparions à celle qu'elle avoit avant l'expérience. *Baudeloque*. tom. ii. p. 17.

1635. On peut conclure d'après ces expériences. 1°. Que la réduction qu'éprouve la tête de l'enfant entre les serres du forceps est différente à quelques égards, selon que les os du crâne présentent *plus ou moins de solidité au terme de la naissance* et que les sutures, ainsi que les fontanelles, sont plus ou moins serrées. 2°. Que cette réduction ne sauroit être en aucun cas aussi grande que des accoucheurs l'ont annoncée, et qu'elle ira *difficilement et bien rarement, au-delà de quatre à cinq lignes*, lorsqu'il instrument agira sur les côtés de la tête. 3°. Qu'on ne doit jamais évaluer son étendue d'après l'écartement des branches de l'instrument, à l'extrémité opposée à celle des serres, et le degré de rapprochement qu'on leur fait éprouver avant d'extraire la tête, ni d'après les forces qu'on emploie pour les rapprocher ainsi. 4°. Enfin, que les diamètres qui croisent celui suivant lequel on comprime la tête, loin de s'augmenter dans les mêmes proportions que celui-ci diminue, ne s'augmentent pas même pour l'ordinaire d'un quart de ligne: et en deviennent quelquefois plus petits (Op. Cit. p. 20, 21).

The following are the results of eight experiments, briefly stated:—

No.	Reduction in Bi-Parietal M.	
1	3 lines	* At this degree of reduction, the suture was torn, and the brain escaped.
2	2 lines	* Instrument bent.
3	2 lines	
4	4 lines	Bones very soft, sutures and fontanelles loose.
5	4 lines	Equally soft.
6	4½ lines	The same.
7	3 lines	
8	2 lines	The ninth experiment is not stated by Baudeloque.

ture to exert if the child were living, and yet the space gained was scarcely sufficient to admit the blades of the instrument to be introduced within the pelvis. They seem to me, therefore, conclusive as to the limited power of the forceps when used as a compressing instrument. Nor can I agree in the opinion of Dr. Rigby, that "the slow and gradual pressure of the forceps thus exerted (by tying the handles tightly together, and tightening them after every successive effort,) upon the head of a living fœtus, will have a very different result to that of the experiments of Baudeloque, and others, in attempting to compress the head of a dead fœtus by the application of a sudden and powerful force." * It is impossible to grasp the forceps for the purpose of moving the impacted head, without applying to it a sudden and powerful force; and if this force be maintained, in the interval of the pains, by ligature, such powerful, constant, and, at the same time, unequal pressure, acting on the head of the child, would appear to me much more hazardous than even the compression of the contracted pelvis, which is known to be a frequent cause of the child's death.

The possibility, therefore, of reducing by the forceps the impacted head to that degree that will enable you to draw it safely through the pelvis, seems to me extremely doubtful. If it were the large head of the male child, advanced in its ossification, and wedged in the deep narrow cavity of the masculine pelvis, I would say it is impossible. The only case where it might, perhaps, be successfully attempted, is in the diseased pelvis, which may be capable of some degree of expansion, and where the head of the child, being less ossified, is softer, and more compressible. †

Let me now direct your attention from the child to the mother; and admitting it is possible, and only possible, to save the former, let us inquire into the risk to which the latter is exposed in the attempt to accomplish this object. The very nature of the case implies an unusual degree of pressure on the soft parts between the

* Rigby's Midwifery, p. 142.

† On this point, I am much more disposed to agree with Dr. Meigs, who remarks—"One of the most dangerous errors relative to the forceps that a student could take up, would be the opinion that the forceps is a compressive instrument by its very design: it is not so—it is an extractor—it is a *tire tête*; and I think it ought to be established as a principle in obstetrics, that where there is not space enough for the descent of the head without the forceps, there cannot be produced a due proportion by merely squeezing the head down to the required dimensions by such an instrument."—Meigs' Philadelphia Practice of Midwifery, p. 295.

head and the pelvis ; congestion must be the result ; and if inflammation have not already taken place, the passages are in such a state that inflammation could be most easily excited. The blades of the best contrived forceps cannot be applied to the head when it is tightly impacted in the pelvis, without bruising the soft parts to a certain extent. This contusion becomes a centre around which inflammation takes place, may increase to any extent, and terminate either in a local slough of the compressed part, or a general gangrene of the vagina, if the inflammation assume an erysipelatous type. In the former case, the separation of the slough may be the formation of vesico-vaginal fistula. In the latter, death may be the result. You will admit that such consequences are of too grave a nature to hazard for the *slight chance* of saving the child ; and therefore, when you are placed in the unpleasant alternative, either to save the child at the risk of the mother's life, or to sacrifice the child in order to preserve her, you must adopt the maxim which governs British midwifery, and consider the safety of the mother to be your first object. But you will seldom be placed in such a dilemma, if you observe closely a case of this description. If you are satisfied that the forceps cannot be safely introduced ; if you think that you cannot compress the head sufficiently to extract it without exposing your patient to a tremendous hazard ; it does not follow that you must destroy the child in order to deliver her. In the great majority of such cases, Nature provides against the difficulty of the case by doing so herself. When the head is thus wedged, the liquor amnii discharged, and the uterus strongly contracted about the body of the child, it is seldom saved from the effect of this extreme pressure ; its death is the result ; and if the case be left altogether to itself, the child becomes putrid, the bones of the head looser and more compressible, and thus it is possible that it might be expelled by the uterus. Formerly, it was customary to wait for "these signs of the death of the child," before perforating ; but being those of putrescency, the patient was exposed to all the consequences that would follow decomposition of its tissues in the uterus, and hence, the death of the mother was too often the result. But now we have it in our power to ascertain its death by another means, which is available long before putrescency takes place. The stethoscope has been found to be a valuable aid to the obstetrician ; it sometimes enables him to determine the exist-

ence of pregnancy when all other means fail ; but I know of no case where it is of more important service than in that which is before us, nor is there any in which its evidence is more certain. In pregnancy, when the child is small, the liquor amnii abundant, or the muscles of the abdomen strong, the foetal heart may not be heard ; but in parturition, when the liquor amnii is discharged, the child full grown, and perhaps large, the muscles of the abdomen stretched to their fullest extent, its pulsations are perfectly audible ; and if once heard, there can be no change in the situation of the sound, because the child is fixed in its position. A close attention, therefore, to the foetal pulsations is necessary in such a case ; and when they rapidly increase in frequency, then intermit, again return more feebly, and ultimately cease, you can have no doubt the death of the child has taken place. To prove to you the value of the evidence in this way obtained, I shall quote the very important experience of Dr. Collins, as to these kinds of labours. He says : “ I have no difficulty in stating, and *that after the most anxious and minute attention to this point*, that where the patient has been properly treated from the commencement of her labour, where strict attention has been paid to keep her cool, and her mind easy, where stimulants of all kinds have been prohibited, and the necessary attention paid to the state of her bowels and bladder, that, under such management, *the death of the child takes place, in laborious and difficult labour, before the symptoms become so alarming as to cause any experienced physician to lessen the head*. This is a fact I have ascertained, beyond all doubt, by the stethoscope, the use of which has exhibited to me the great errors I committed before I was acquainted with its application to midwifery, viz.—*in delaying delivery often, I have no doubt, so as to render the result precarious in the extreme, and in some cases even fatal.*” * This observation of Dr. Collins would apply to many cases of perforation that are recorded, and which have been followed by such frightful consequences, that it is not surprising that they should excite the disgust of the profession. They were cases where the operation was useless, because performed too late. By means of the stethoscope it is in your power to prevent this, and to deliver the child in sufficient time to save the mother from injury. No one is justified in destroying a living child, unless there

* Collins's Practical Treatise, p. 16.

is clear evidence, from the symptoms, that the mother is in danger. According to the old rule of practice, therefore, you were placed in the dilemma, either to wait for such symptoms, or for the signs of putrefaction in the child—alternatives equally dangerous to her; but if the death of the child can be known the moment it takes place, and if it be true that its death precedes those dangerous symptoms, it is obvious that its removal by the crotchet is no longer objectionable, and perforation is deprived of all its horrors. So far as the safety of the mother and the preservation of the passages from injury are concerned, there is no comparison between perforation and the forceps. In this respect perforation is a far safer operation, if ordinary caution be exercised; the objection—the sole objection—that condemns it, is the fact that the child must be destroyed, either by the uterus, or by the instrument. We freely admit the cogency of the argument; but when it is weighed against the still greater objection, that in the attempt to save the child the soft parts of the mother may be injured to a most dangerous extent, while the preservation of the child is extremely doubtful—when we find, in the imperfect history of these operations, such as they are given to us, that the child is very generally lost, or, if there be an exception in which the child is with difficulty saved, the case is recorded with that triumphant acclamation that proves the success to be unexpected,—when the risk to the mother is so great, and the prospective advantage so doubtful, you will admit that the balance is in favour of an operation by which, if properly performed, and with sufficient promptitude, the safety of the mother is at least secured.

We are not generally favoured with a faithful history of cases that illustrate the mischievous effects produced by the forceps. On the contrary, while the post-partum accidents of a skilful operation are deeply concealed in the shadows of the back ground of the picture, the surprising, the almost miraculous, power of the instrument is put prominently forward, with all the vividness of a most glowing and high-coloured description. Thus the truth is concealed from you, and so would remain, until exposed by your own dear-bought experience, except that you find scattered through the works of men whose skill is acknowledged, ominous hints and anxious warnings against the improper application of these instruments. Many evidences might be quoted to this effect: we shall direct your attention to a few of them. Your

late respected professor, Dr. Davis, paid a great deal of attention to the subject of instrumental labours, and was disposed to advocate a much bolder use of the forceps than what I should recommend; nevertheless, he candidly admits, that “of all the instruments used in the practice of midwifery, those of the present class [the forceps] are unquestionably *the most dangerous to the mother*, inasmuch as in all cases where the forceps are used, the maternal tissues are more or less liable to contusion. All the fangs and framework of the instrument are made of tempered steel, and let them be ever so well covered and defended, they will still retain a great degree of hardness, calculated to bruise and to fret the soft and living texture which might be interposed between their covered surfaces and the solid walls of the pelvis.”*

The same impression of mischief leads Dr. F. Ramsbotham to warn the practitioner that “cautiously and tenderly must this iron instrument be used! We must recollect that no sensation can be imparted to the operator’s hand of any injury that may be done to the woman; and we must remember that one injudicious thrust, one forcible attempt at introduction, one violent effort at extraction, may bruise, may lacerate, may destroy!”† Dr. Blundell addresses his pupils thus—“When, however, you lay your hand upon the tractor, or forceps, remember, that the accoucheur who is meddlesome may be guilty of occasioning laceration of the perinæum, rupture of the vagina, compression and death of the child, inflammation of the abdomen of the mother, and many other fatal consequences, *which I myself have had occasion to see*—a list of offences surely sufficient to alarm the prudent.”‡

But let us come to more direct evidence. Riecke, in his report of the practice of the Kingdom of Wurtemberg, gives the results of a very large number of cases, and amongst them those in which the attempt was made unsuccessfully to remove the impacted head by the forceps. He observes—“Almost always, perforation was preceded by attempts to apply the forceps, and to the great injury of the mothers, because perforations, not

* Davis’s Obstetric Medicine, p. 786, 8vo. edition.

† Ramsbotham’s Obstetric Medicine and Surgery, p. 299.

‡ Blundell, by Castle, p. 526.

preceded by such attempts, presented much more favourable results. . . . The trials at extractions with the forceps—which many accoucheurs continue to the extinction of the infant's life (although foreseeing the necessity for perforation)—exhaust the mother to that degree, that she necessarily sinks under the effect of these violent efforts.”* In allusion to similar inquiries, Dr. Collins remarks—“It is from being thoroughly convinced of these facts by long and extensive observation, that I consider the forceps quite inapplicable when the head becomes fixed in the pelvis, and the ear cannot be reached by the finger except by violence, in consequence of disproportion existing between the head and the pelvis. . . . The results I have witnessed from such practice [delivery by the forceps] were most distressing: in some, the neck of the bladder or urethra either lacerated or the injury by pressure from the forceps so great as to produce sloughing and consequent incontinence of urine; in others, the recto-vaginal septum destroyed, either of which renders the sufferer miserable for life; and in two cases where the mouth of the womb was imperfectly dilated, so much injury inflicted on this part as to terminate in death.”† Dr. R. Lee, in his lectures, quotes the paragraph at full length from which these passages are extracted,

* Riecke, in his report, gives 84 cases of perforation, in 31 of which the mother died, being in a proportion rather more than one in three. He explains it thus:—

“Presque toujours la perforation du crâne avait été précédée de tentatives pour appliquer la forceps, et cela au détriment des mères: car les perforations non précédées de ces tentatives offrent des résultats *beaucoup plus favorables que les autres*. La répugnance des accoucheurs à pratiquer la perforation du crâne lorsque l'enfant est encore en vie d'une influence très fâcheuse pour les mères. . . . Les tentatives d'extraction avec le forceps que beaucoup d'accoucheurs continuent jusqu'à l'extinction de la vie d'enfant tout en prévoyant la nécessité de la perforation du crâne épuisent les mères au point qu'elles succombent presque nécessairement à la suite de ces efforts violents. Les suites ne sont pas moins fâcheuses lorsque l'accoucheur se décide à attendre patiemment que l'enfant soit mort.* La perforation du crâne exécutée avec les précautions convenables n'est pas en elle même une opération bien dangereuse. Parmi les femmes soumises à cette opération il s'en est trouvé une que l'a subie dans onze accouchemens consécutifs et jamais elle n'en a éprouvé de suites fâcheuses.”—*Archives Médicales*. tom. xxii. p. 375.

† Collins's Practical Treatise, p. 12-13.

* The means of determining the death of the child by the stethoscope is not alluded to by Riecke.

and adds—"The accuracy of these remarks is fully confirmed, by all the forceps cases which have come under my observation, which exceed sixty in number."* It would occupy too much time to accumulate further testimony to the same effect.† I trust sufficient has been placed before you to authorize the conclusions at which I have arrived, and which are now submitted to you—viz., that when the head is impacted in the pelvic cavity, it cannot be delivered by the forceps without such injury to the passages as might endanger the mother's life; that the probability of preserving the child's life is not sufficiently certain to justify an attempt which might be so hazardous; that in the great majority of these cases the death of the child takes place naturally, and it may be removed before symptoms dangerous to the mother present themselves; and lastly, that if it should happen that the reverse occurs, and danger to the mother—whether from exhaustion or extending inflammation—is indicated before the death of the child, that then perforation is called for, rather than render the risk to the mother a certainty, by the dangers that result from a forcible extraction by the forceps.

We have been obliged to dwell at some length on these disputed questions connected with the practice of midwifery, when the head is fixed in the cavity of the pelvis; we shall therefore only allude very briefly to the last stage of delay, when—

The head is retarded at the outlet.—When the head is in this position, it may arise from the perinæum being rigid, or from the arch of the pubis being too narrow, so that the head cannot pass out between the ischio-pubic rami. In either case, there is a long-continued pressure on the perinæum, which must excite inflammation, and increase its rigidity, if the delivery of the head be not assisted. The strictest attention is necessary to subdue any tendency to inflammation in the perinæum; fomentations must be sedulously employed, and, if necessary, depletion

* Lee's Lectures, p. 305.

† "I wish that my present subject permitted me also to state what I have found on dissecting the parts after the use of the crotchet, and, *in particular, where the forceps had been used*, as I must presume, in a case improper for them. The injury which the seemingly harmless instrument, the forceps, is capable of doing, might then be proved, and a wholesome admonition given to young surgeons."—Sir Charles Bell on the Muscularity of the Uterus; *Medico-Chirurgical Transactions*, vol. iv. p. 339.


by leeches. By this means a rigid perinæum will in many instances gradually yield to the head, and allow it to pass; but sometimes the vectis may be passed on the pubic side of the head to assist its advance. Great caution is required not to bring the head down too suddenly on the perinæum, which must also be protected by the counter-pressure of the hand. Where the difficulty arises from narrowness of the pubic arch, the forceps is preferable, both because you have more power to overcome the resistance of the tubera ischii, and there is less danger of the pubic blade injuring the adjacent soft parts than if the vectis were employed, and much force used in the extraction.

ACCIDENTAL OBSTRUCTIONS are occasionally causes of difficulty in the second stage of labour, and when such are presented to your notice they always demand the most serious attention. First, because the majority of them depend upon some organic disease, the existence of which renders labour, if at all severe, extremely dangerous to the patient. Secondly, because the nature of the obstruction is often very obscure; and if a tumour impede the progress of the head, it may be doubtful whether it is ovarian, polypoid, or malignant. The extent of its attachments, and the possibility of removing it, may also be a difficulty. You have many reasons, therefore, for extreme caution and a very guarded prognosis under such circumstances. Tumours of this kind are, to a certain extent, moveable, and may be either soft and fluctuating, or firm and elastic. Sometimes the sacrum becomes the seat of osteo-sarcoma, and then a hard and perfectly unyielding tumour obstructs the head.

Beside these tumours, another description of obstacle is occasionally met with, the consequence of previously existing inflammation. Cases have been observed *where the os uteri has been agglutinated* in such a manner as altogether to prevent its dilatation. One case is recorded by Dr. Ashwell, in which the os uteri was absent; nor could the first stage proceed, until an artificial opening was made to allow the head to pass. These are, however, very rare cases; a more common effect is the alteration produced in the vagina. Sometimes a *band is found crossing the vagina* and preventing the advance of the head. *The walls of the vagina are occasionally united* for a certain distance, and, again, *a stricture may be formed in the vagina*, just as in the urethra. It is very necessary to make the most careful examina-

tion when a case of this kind presents itself, because the extent of the injury done is not always confined to the part which obstructs the head; partial sloughs of the vagina are just as likely to be the result of the previous inflammation, and when these have separated, the portions that have been thus partially destroyed must have been left extremely thin, and badly calculated to resist any extreme pressure. Thus the vagina may be very easily lacerated if labour is suffered to continue for any great length of time.* When a band is formed in this manner, it may easily be divided by a guarded bistoury, and the obstruction removed; but when a stricture is the cause of difficulty, more care is necessary. The head should be pressed back in the interval of the pains, and the portion of the vagina above the stricture accurately examined; if it be of the usual strength and thickness, or if its density be increased, it is better to allow labour to proceed, in order that the head may gradually dilate the stricture; but if, on the contrary, the vagina be felt unusually thin, it would be extremely dangerous to do so, because the head may force its way through the attenuated portion, and perhaps convert the rectum and vagina into one common cloaca. It would be preferable to divide the stricture in different places, so that it may yield to the head more readily. When the walls of the vagina cohere, the case becomes very embarrassing, because they must be separated; and yet there is a danger that the vagina may be cut through, if great caution be not exercised. You have also the additional difficulty, that it is almost impossible to ascertain the condition of the vagina behind the portion that is united. Some risk must therefore be encountered. It is better to allow the labour to proceed sufficiently far to determine the extent to which the head may separate the parts adherent, employing every necessary means to counteract any inflammation that may arise. The head may overcome the resistance to a certain extent, so as to render the division of the remaining portion much safer and more easy; but if this cannot be accomplished, the walls of the vagina must be dilated, so as to expose perfectly the adhesion which is necessary to divide by cautious and frequently repeated incisions with the knife.

* In a very interesting paper on this subject in the Dublin Journal, vol. xxiii., Dr. Doherty states some facts worthy of attention. He mentions one instance of a



Ovarian Tumour.—Merriman.

Ovarian tumours sometimes descend into the pelvic cavity, and obstruct the head of the child. If the tumour consist of several cysts, the smallest may pass down between the vagina and rec-

“slough of the entire inner surface of the vagina. From the vulva to the os uteri not a point escaped; and to prove how care and attention can prevent ill-consequences, even in such a case as this, I would remark that the patient who suffered this loss of substance, left the hospital with the passage almost as capacious as before delivery,” (p. 67). In another case, where there was occlusion, “There was no *direct* communication between the vagina and uterus; an orifice *below* the occlusion admitted the finger into the cavity of the bladder, into which the os uteri could be felt to project, owing to the destruction of the vesico-vaginal septum also *above* the adhesion.” This patient was not pregnant.

The advantages of nauseating doses of tartarised antimony in relaxing a rigid vagina, Dr. D. illustrates by the case of a middle-aged woman, who was admitted into the Dublin Lying-in Hospital in labor of her first child. “On examination, the finger could be passed up the vagina only with the greatest difficulty, and when introduced, felt as if surrounded by a cylinder of iron, so thick and hard were the walls of the canal and perinæum. They seemed in truth not to possess the slightest elasticity; so much so, indeed, that I was some time before I could persuade myself that it was not one of those hard, encysted tumours surrounding the rectum that my finger came in contact with. And yet all this rigidity disappeared in less than six hours, under the use of tartar emetic, and the child was expelled without any difficulty” (p. 69.)

“While I deprecate the rashness of operating without a well-marked necessity, I

tum; cases are also recorded where very large tumours are found in the same situation. One of these cases is given by Dr. Merriman, along with a very accurate drawing of the tumour. Their contents vary so very much in their consistence and density, that they are not always easily recognised; but if there be any sense of fluctuation, or even if the tumour be very elastic, the probability is, that it is an ovarian cyst, containing fluid more or less deeply seated. Unless the size be great, it is possible that the head may press the cyst against the sides of the pelvis, and pass below it; a small tumour also may be pushed back towards the brim of the pelvis, when the pains are absent, and perhaps be prevented from again descending when the action of the uterus returns; if by neither of these modes the removal of the obstacle can be accomplished, the only resource left is to puncture the tumour and allow the fluid to escape. This may be done, although there be no distinct sense of fluctuation, because the fluid is often thick, like honey, and may be deep-seated, which will communicate to the fingers an elastic feel rather than that of fluctuation. Besides, if a mistake be made, and you should puncture a polypus or a fibrous tumour, no great injury is done. It is when the sense of fluctuation is distinct that caution is required, especially if the tumour should appear towards the pubic side of the pelvis. The bladder has sometimes prolapsed before the head of the child, and presented a fluctuating tumour. It is not necessary to tell you that this should not be punctured. But the danger of these cases does not generally arise from the delivery being obstructed, but from

would at the same time beg to dwell on the fatal consequences which may ensue from postponing the use of the knife beyond the moment at which it becomes necessary. Every additional instant then adds to the risk of laceration, and of course, increases the chance of a lamentable termination to both mother and child; a fact which was forcibly impressed upon me by a case wherein the attendant, while sitting at the bed-side with the scalpel in his hand, delayed employing it, in order to afford another practitioner who was present, an opportunity of examining the part; *meanwhile a strong pain suddenly came on, and the vagina was extensively torn* (p. 72.) In a case of face presentation, a thick band surrounded the upper portion of the vagina. This band was nicked in one or two places, the pains continuing of natural strength and frequency; the face gradually descended, and as it progressed the remainder of the band appeared to relax, and every hope was entertained of a fortunate issue, till about two o'clock, P. M., when rupture took place (p. 73).

the effect which labour produces on the disease ; the tumour is necessarily exposed to a great deal of irritation ; the patient is weakened if not exhausted by the struggle which takes place ; and, when labour is concluded, she is quite unequal to combat the effects of that irritation. Dr. Merriman has collected the history of eighteen cases of ovarian tumours obstructing parturition. One half the mothers died, three recovered very imperfectly, and six only may be said to have escaped ; sixteen of the children were still-born, and four were born alive. I shall quote from Dr. Merriman's work the results of these cases as he has given them, from which it appears that the greatest success attended those cases where the tumour was opened :—

“From the enumeration of the cases already referred to, it appears that

“Twice, the labour was effected by the pains, unassisted by the art of the accoucheur ; but one of these women lost her life, and one of the children was still-born.

“Five times, the perforator was used after a longer or shorter duration of labour. Three of these women died, another recovered very imperfectly, and one got well.


“Five times, the labour was terminated by turning the child ; all the children were lost, and only one mother recovered.

“Three times, the tumours having been opened, the labour was afterwards trusted to nature. Two of these women recovered, but the other remained for a long time in an ill state of health. Two only of the children were preserved. In three cases, the tumours being opened, it was still found necessary to have recourse to the perforator. One of these women died, one remained in an ill state of health for eighteen months, and then sunk under her sufferings ; the third recovered.”

Dr. Merriman remarks, “Upon the whole, the evidence we at present possess is more in favour of opening tumours when they contain a fluid, than of any other mode of procedure ; for of the nine women who recovered more or less perfectly, *five appear to owe their safety to this operation*, and of the children born alive, two were preserved by the same means.”*

Polypus has been found sometimes to interfere with delivery.

* Merriman on Difficult Parturition.



Polypus.—Ramsbotham.

An interesting case of this kind is given in Dr. F. Ramsbotham's "Midwifery,"* in which the polypus nearly filled the pelvic cavity. However, the mouth of the womb dilated rapidly, the membranes burst speedily, and in less than an hour after my [Dr. Ramsbotham's] arrival, the head, under the action of powerful throes, forced the principal bulk of the tumour external to the vulva (which still, nevertheless, retained its attachment to the uterus by the stem), and itself instantly followed."† Thus it is possible that a moveable tumour of this kind, although very large, may be driven down before the head. If it be small, and detected early in labour, it might also be in your power to prevent the tumour descending. It might be pressed back when the pain ceases, and so retained until the head passes beyond it. But if neither can be accomplished, if the tumour remain an impassable barrier, it should be removed, not by ligature, but by excision; the polypus should be drawn down as much as possible by a forceps proper for the purpose, a temporary ligature applied, and the stem cut through.


We shall not here enter into a discussion of the comparative merits of the treatment of polypus by ligature and by excision; this must be reserved for another opportunity. But in reference to the present case, we would only observe that the risk of dangerous hæmorrhage after excision is not so great as to justify the

* Ramsbotham's Principles and Practice, etc., p. 237.

† Op. cit. p. 237-S.

adoption of the only alternative—destroying the child. You will have little difficulty in recognizing polypus when it descends so low into the vagina as to interfere with labour. Beside the firm, fleshy feel of the tumour, it is extremely moveable, and when the head is pressed back in the interval of the pains, its pyriform shape and long narrow stem will be more obvious. It is not likely that the ovum could be brought to maturity if a large polypus occupied the cavity of the uterus; it is, therefore, fair to assume, that when a polypus is found to impede parturition, it must be attached to the mouth of the uterus, and therefore it can be the more easily traced to its origin, so that you have every facility to assist your diagnosis.

Another cause of obstruction may arise from a fibrous tumour of the uterus. A remarkable and very interesting case of this kind is recorded by Dr. Beatty,* in which the tumour was so large and apparently so attached, as it was thought would render the Cæsarian section necessary. It was agreed, however, to wait, and to observe the action of the uterus, as long as it might be done with safety. After some time, and to the surprise of those in attendance, the tumour appeared to retreat from its situation, while the child began to occupy its place, and to present its foot; this was seized, and the delivery with great difficulty completed. The child was still-born, but the mother recovered.



Osteo-sarcoma.

* Dublin Medical Journal, vol. xvii. p. 411.

OSTEO-SARCOMA sometimes grows from the sacrum. The bony tumour may be so large as to render delivery *per vias naturales* impossible, and therefore the Cæsarian section must be had recourse to. But it may be small enough to prevent this necessity, although it may be difficult to save the child. A case of this kind came under my own observation, where a tumour, about the size of an orange, was connected to the middle of the sacrum; it was perfectly immovable, and of bony hardness; the head of the child could not pass it, nor was there the least hope that it could be drawn by the forceps through the narrow space left in the pelvic cavity; the head was therefore perforated, and the child removed: the mother perfectly recovered.

Beside these more usual causes of obstruction to delivery, Dr. Drew had detailed, many years ago,* two very remarkable cases of tumours in the pelvis. The first patient, who was not pregnant, died in consequence of it. An examination was made after death: there seemed to grow from the left sacro-sciatic ligament a tumour, which “was perfectly round, about sixteen inches in circumference, of a fat, gristly substance, without any appearance of circulation in it.” The root seemed to be its principal attachment, because when that was cut through it came away quite easily. The result of this inspection satisfied Dr. Drew of the practicability of removing it by operation; and, although a rare variety of tumour, it so happened that very soon after (as is often the case) a similar tumour came under his notice when the patient was in labour.

Dr. Drew states, “It was exactly the same. The tumour grew out of the right side, and occupied the whole cavity of the pelvis so completely as to admit of passing only one finger between it and the pubis, by which I could scarcely reach the head of the child.” Dr. Drew proposed to remove it, which was assented to. An incision was made through the perinæum, at the right side; the tumour was exposed, the finger passed before and behind its root, which was easily divided with a knife, and brought away. The wound being dressed, labour proceeded, and in six hours, the head being within reach of the forceps, was delivered safely. The patient recovered rapidly. These rare cases may present themselves: if such should happen,

* Edinburgh Medical and Surgical Journal, vol. i. 1805, p. 20.

you have sufficient encouragement not to despair altogether of giving relief.

We have now concluded the consideration of difficult labours, so far as their causes, symptoms, and general treatment, are concerned. We have directed your attention to cases where the aid of the vectis, the forceps, or the perforator, is called for. It remains to us, therefore, to consider the mode of applying these instruments.

LECTURE XI.

OBSTETRIC OPERATIONS—THE FORCEPS.

Instruments employed for preserving the Lives of the Mother and the Child—The Vectis—Of limited Application—Mr. Gaitskill's manner of using it—Proposed mode of Operation—Objection to use it as a Lever—The Forceps—Operation with the Short Forceps when the Head is resting on the Perinæum—Operation when the Head is in the Pelvic Cavity—Operation when the Head is fixed in the Brim of the Pelvis.

IN the three preceding lectures, we endeavoured to point out to you the situations at which the head may be impeded in its progress through the pelvis, and the varieties in the degree to which it may be compressed; we were also desirous to place before you the evidence upon which the conclusions therein stated were founded, in reference to disputed questions as to the rule of practice in certain cases of difficult labour. Turning from those controversial subjects, and, leaving the questions when instrumental aid is called for, and what kind of instruments should be employed, your attention must now be directed to an equally important subject—obstetric operations.

The instruments employed in operative midwifery may be arranged into three classes. 1st. Those calculated to preserve the lives both of mother and child, as *the vectis, the forceps. The fillet* was formerly used for the same purpose, but is now discarded from practice. 2nd. When the preservation of both lives is impossible, those intended to preserve the life of the mother by sacrificing the child. These include, *the perforator and crotchet, the craniotomy forceps, the osteotomist, the cephalotribe.* 3rd. When the delivery of the child cannot be effected even by such means, and the safety of the mother is more than doubtful, there still remains the operation of opening the uterus

through the abdomen, and thus removing the child, with some chance, at least, that it may survive.

The rules, therefore, which govern the application of instruments, are founded upon these three principles :—1st, *to preserve the lives of the mother and the child* ; if this be doubtful, 2nd, *to preserve the life of the mother without reference to the child* ; and when this cannot be done, or, at least, seems so from the circumstances, 3rd, *to save the child, if possible*.

The *vectis* and *forceps* belong to the first class. The *vectis* consists of a single blade, shaped like a blade of the forceps, only more abruptly curved, and when used in the manner we have recommended, it is intended to act as an extractor, to assist the feeble action of the uterus, to correct malpositions of the head, or to overcome any unusual resistance of the perinæum. It is not, therefore, an instrument of much power, and its use is limited to the removal of slight impediments to the passage of the head. The advocates for this instrument do not, however, confine themselves to such a restricted application. They employ it as a substitute for the forceps, and even claim for it a superiority over the long forceps, in those cases in which the head is arrested in the brim of the pelvis. We must dissent from such a view of the utility of the *vectis* : in order to give to it the same power which the forceps acquires by the counter-pressure of the blades, an amount of force must be employed which might be very dangerous to the patient. Imagine the head fixed in the brim of the pelvis, the *vectis* applied to the occiput, the practitioner using all his strength to extract, and at the same time to keep the *vectis* in its position, and the instrument slipping from its situation into the vagina ; by supposing such a case, you can readily understand our objection. Nevertheless, as the operation of delivery by the *vectis* at the brim of the pelvis has received the support of Dr. Blundell, who recommends the instrument improved by Mr. Gaitskill, and his mode of using it, we shall briefly quote Mr. Gaitskill's rules for applying it in such cases.

After giving directions as to placing the patient, etc., he proceeds, "The preliminaries being settled, the next thing is, the safe introduction of the instrument. To do this with facility and safety, the accoucheur should kneel on a pillow by the side of the bed, and introduce all the fingers into the vagina as far as

the brim of the pelvis, at the side of the sacral promontory (either right or left, according to the situation of the occiput); as he passes up the instrument the fingers should be gradually withdrawn. The instrument is to be pressed up into the cavity of the uterus, being careful that it is in the inside and not on the outside, gliding it over the parietal bone till the screw part of the handle presses on the fourchette of the os externum. This attained, the handle should now be held firmly in the right hand, while the index and middle finger of the left, fixed about two inches from the screw part within the vagina, become a fulcrum. On this fulcrum, or point of support, the instrument is made to move from the sacro-iliac symphysis towards the hollow of the ilium, by the action of the right hand on the handle. In this way it describes the section of a circle, and glides on the occiput. Should the occiput point to the right ilium, the left hand must be employed; if to the left ilium, the right hand must be used. When a labour pain takes place, the accoucheur should gently aid it by drawing down in the line of the axis of the pelvis—i. e., an imaginary line, directed from the umbilicus through the centre of the axis of the pelvis. In this way the occiput is depressed, while the chin approaches the child's breast, and its head is reduced to the smallest compass, and is thus enabled to pass through the cavity of the pelvis. As soon as the occiput is brought so low as to press on the perinæum, the instrument should be withdrawn, and re-introduced with the usual precautions. The object now in view is, to place the instrument over the face of the child. To effect this, the hand must be passed up, as at first directed, to the right or left sacro-iliac symphysis, according to the situation of the face. When the instrument gets above the brim of the pelvis, a finger or two must be inserted by the side of the instrument, and pressed on till it (the instrument) passes over the forehead on to the face, so as to embrace the chin. An imaginary line drawn through the centre of the child's mouth, ear, and occiput, is the present situation of the instrument, and quite the reverse of what it was before. The practitioner has now nothing to do but to draw down during the time of pain, increasing the power according to the degree of resistance.”*


* London Medical Repository, 1823, p. 379—381.

Such is the mode in which Mr. Gaitskill applied it when the head was high up within the cavity, or in the brim of the pelvis ; but we confess our fears to recommend to you such a manner of employing the vectis. The cases in which it may be used with most advantage, are those in which the head is arrested at the outlet, in consequence of the uterus being unable to overcome the resistance of the perinæum. So long as the pains continue with any regularity and strength, you should not interfere, except for the purpose of preventing inflammation ; but when the pains become feeble, suspended, or return at long and irregular intervals, then the vectis may be applied with even more advantage than the forceps, because there is less risk of injuring the perinæum. We shall proceed to describe to you the manner of performing such an operation.

You must first observe those preliminary measures necessary in all obstetric operations. The urine should be withdrawn from the bladder by an elastic gum catheter, of rather a large size (No. 10), and without a stilette. It is always safer to use a catheter of this kind because there is less risk of injuring the urethra, if it should be compressed, than if the unyielding silver catheter were employed. An enema should also be administered, to relieve the large intestines : and when these points are secured, the patient, lying on her left side, should be drawn as near to the edge of the bedstead as possible. The pelvis must be raised more than usual, and if the patient has been lying on a bed, and not on a mattress, it would be advisable to place a hair cushion under the hips. Adopting Mr. Gaitskill's position, you may kneel with one knee on a pillow, and in the interval of the pains introduce the first two fingers of the right hand between the head and the symphysis pubis : passing them on either side of the symphysis, the tip of the ear will be felt without difficulty : the finger must remain applied to it while the vectis is being introduced : the instrument should be held about the middle, between the two forefingers and thumb of the left hand, and the handle directed obliquely downwards and backwards towards the coccyx, in order that the blade may lie flat upon the head when the instrument is passing between it and the fingers of the right hand. This is difficult to do when the vectis is very abruptly curved, and therefore the curvature of the blade makes a very essential

difference in the value of the instrument. If too much curved, it cannot easily be applied; if too gradually curved, like the forceps, it will slip from its position if any force be used in extraction. You must, therefore, attend to this point particularly, in the selection of the instrument you use. When the blade is so applied, press it gently forwards with a slightly oscillating motion, until the edge reaches the ear, which is now placed exactly between the finger and the vectis. The handle must, therefore, be depressed still more, in order that the edge may pass over the ear; when this is safely accomplished, the finger may be withdrawn, and the vectis passed forwards to its proper position. The direction of the handle is now completely altered, and looks downwards and forwards, its junction with the blade corresponding nearly to the sub-pubic ligament. When the instrument is thus applied, "then (in the language of Dr. Denman), grasping the handle of the instrument firmly in the right hand, wait for the accession of a pain," which, although absent before, almost always returns when this new irritation is applied to the uterus. While you assist the pain at this stage of the operation, great caution is required. It is here that the mischievous principle of the action of the instrument, as a lever, may do so much injury. If the pubis, or ischio-pubic ramus be made the fulcrum, the soft parts must be contused, and a slough may be the result. If, to avoid this, the fingers of the left hand press the blade strongly against the head, and thus guard the soft parts from pressure, making, as it is said, a fulcrum of the fingers, the lever is only altered from one of the first order to one of the second. In the first, the pubis is the fulcrum. In the second, the head and face of the child. Serious injury may be done in either case. You should, therefore, carefully avoid using the vectis as a lever; and in order to do this the more certainly, it is better to pass two fingers of the left hand between the head and the perinæum, and to grasp the shank of the instrument with the remaining fingers; counter-pressure is thus made similar to the forceps, and the vectis may be used solely as a tractor. Again, in the language of Denman, we would say, "When the pain ceases, let the instrument rest, and on its return, repeat the same kind of action, alternately resting and acting, in imitation of the manner of the pains." Proceeding thus cautiously, the head will soon advance and press strongly on the perinæum.

The introduced fingers may then be withdrawn, and the vectis maintained in its position, rather for the purpose of acting with it, if the pains should again become feeble, than to extract the head by its means, if the uterus be sufficient to expel it; thus the perinæum will be better secured from injury.




Extraction with Vectis.*

When *the vectis is used to correct mal-positions of the head*, it is better not to use one too much curved; one blade of the forceps will often answer in these cases; the head is higher in the pelvic cavity, and does not generally press on the perinæum; too great a curvature would interfere with its introduction; while, on the other hand, there is seldom occasion to use it as an extractor, because when once the correction is made, the head will readily descend without assistance. Independently of its limited power, the vectis is liable to some disadvantages from which the forceps is free, and which should be guarded against. It is necessary to grasp the instrument very firmly, and sometimes to exert your strength to keep it in its place. Sometimes the vectis will not retain the head so securely that the instrument may not slip, and though it is easily replaced, still a good deal of force is necessary to keep it in its position. If, in such a case, the handle be smooth and round, there is also a risk that it may turn in the hand without your knowledge, and therefore do much mischief. The handle should always be made rough, and with one side, at least, flat.

The forceps is more general'y used in the practice of mid-

* In these sketches the perinæum etc., is exposed, to show the position of the head, which is faintly outlined.


wifery, and is an instrument of much more extensive application. It may be employed when the head is at the outlet, in the cavity, or in the brim, of the pelvis. Hence you will find, in obstetric authors, two kinds of forceps spoken of—the long and the short forceps. It is necessary to bear in mind this distinction, because the mode of operating with the latter is not the same as with the former instrument. This is the more important, because there seems to be some degree of confusion in the description given by authors of operations with the long forceps. For instance, when the head is arrested high in the cavity of the pelvis, a longer forceps is required than when it is at the outlet. The former operation is therefore sometimes mentioned as a delivery by the long forceps. In speaking of these operations, we would wish to be understood to mean, not only a different kind of instrument, but a different mode of applying it from that adopted when the short forceps is used. In order to avoid this confusion, we shall describe the operations required in three different cases: first, *when the head is resting on the perinæum*, the operation with the short forceps; secondly, *when it is arrested in the pelvic cavity*, which might be considered an intermediate operation. And lastly, the operation with the long forceps, *when the head is fixed in the brim of the pelvis*.



Passing of the Pubic Blade.

The operation, when the head is resting on the perinæum, may be undertaken in cases similar to those in which the vectis is employed, and is preferable, if there be any diminution in the transverse measurement of the outlet. The preliminary steps of this operation are the same as for the vectis; but it must be re-

membered, that the temperature of these, as well as of all obstetric instruments, should be raised to that of the vagina, and they should be greased before being introduced into the passages. Having made these previous arrangements, the pubic blade of the forceps, with the lock looking upwards, must be passed over the head in a similar manner to the vectis, and when so placed, the handle may be raised towards the pubis, and there maintained by an assistant *in its exact position*. The handle should not be moved to the right or left side, because it is of great importance to observe the precise direction of the pubic blade when the sacral blade is being introduced. Taking, then, the lock of the former as your guide, as soon as the pain ceases, pass two fingers of the left hand between the head and the perinæum, and holding the sacral blade lightly by the handle with the right hand, endeavour to guide it so along the introduced fingers that the edge of the sacral may pass along the lock of the pubic blade. As the sacral blade passes forwards, and the locks approach each other, the handle of the pubic blade should be taken in the left hand, and drawn slowly towards the perinæum. In this manner the locks will glide together, and the instrument be applied without much difficulty.



Introduction of Sacral Blade.

When this is done, and the pains return, the handle of the forceps should be held firmly, and, the perinæum being supported at the same time by an assistant, traction made—at first, very moderately, carefully observing the action of the uterus; and as you perceive that the pains are inefficient, the force may be increased. With each effort the handle may be drawn, first,

with a slightly waving motion to either side, and then upward, towards the pubis, in order that the head may pass in the axis of the vagina. When the head advances, and you are satisfied that the difficulty is overcome, it is better to leave the rest to the uterus so long as it acts, because there is less risk of injury to the perinæum. For the same reason, it would be advisable to unlock the forceps, and to withdraw the sacral blade, leaving the pubic to be made use of as a vectis. When the perinæum is tightly stretched over the blade of the forceps, passing out with the head of the child, it is very easily lacerated. The time that the operation occupies is of no importance. The object you should have in view, is to assist the action of the uterus, not to anticipate the pains, or to hurry the delivery. Thus a considerable time may elapse before the operation is concluded. Be careful, therefore, not to make unguarded promises of prompt relief.

The forceps used in this operation is altogether shorter than that employed in either of the other operations. It is about nine or ten inches long; the distance between the extremities of the blades is about one inch and a half; that between the centre, about three inches. The intention is to prevent the head of the child being much compressed in the effort to extract it. There is some difference, however, in the construction of these instruments, which will be understood by the examples placed before you. The short forceps of Dr. Denman* had the extremities of the blades closer, but the handles were very short, with the same object, that of preventing much compression. The short forceps of Dr. Conquest† has the fenestræ wide, in order that parietal prominences may pass through them. Dr. Aitken's and Dr. Collins's‡ are very similar in shape, and correspond with the description we have just given, only that Dr. Aitken's has a small moveable roller between the handles, to prevent compression.

The operation, when the head is arrested in the pelvic cavity, is one which requires a much more attentive consideration, because it is here that the difference in the practice of the most experienced accoucheurs is so remarkable. We have already submitted to you our reasons for the rules proposed for your adoption. They are, 1st, Not to interfere, or to apply the forceps, if

* (Vide fig. 3, p. 245.) † (Vide fig. 5, p. 245.) ‡ (Vide fig. 4, 6, p. 245.)

the head be slowly advancing, unless it should happen that dangerous constitutional symptoms are approaching. 2ndly, Not to apply the forceps *when the head is impacted*. The view of the operation which we wished you to take would confine it to cases of arrest. It is of importance, therefore, to make a very careful examination, *per vaginam*, before the delivery is determined upon. *First, to ascertain that the head is arrested*. Sometimes it ceases to advance, while the tumour on the presenting part increasing, is mistaken for its further descent. Both fingers, therefore, should be introduced, and passed high up, between the head and pelvis, in the interval of the pains, to determine the arrest. *Secondly, you should decide on the degree of disproportion*. In cases of arrest, the ear can generally be felt, which cannot be done without great difficulty in cases of impaction. Hence, *as a general rule*, to feel the ear is a diagnostic mark of this distinction; but you should not confine your attention to this point alone, because it sometimes happens that when the head is arrested, it is so placed that the ear cannot be felt, especially if the head be lengthened, and a tumour be formed upon it. The object of the rule is to determine the amount of space there is for the introduction of the instrument. Therefore, if the ear be out of reach, while the fingers can be passed with facility between the head and the pelvis—if the catheter can be passed easily—if you can press the head back without difficulty—and if the vagina be not swollen from the extreme pressure—then the forceps may be applied. *Thirdly, you must decide upon the time of its application*. It appears to me that *four hours* would be quite sufficient to allow the head to remain in the same position, to authorise your interference. But if there be the least indication of pain, swelling, or heat in the passages, you should not delay one moment from the time that these symptoms present themselves, when you are satisfied that the forceps may be applied. Promptitude is the secret of success, and in nothing is it more evident than in the case we are supposing. It is possible the pains may be strong and frequent, and it is generally a safe recommendation not to interfere so long as the uterus seems to have sufficient power, but rather to wait until the pains become feeble, or the action of the uterus is suspended. Nevertheless, in the case before us, you cannot act upon such a rule. If the head be arrested—if the


pains be strong but inefficient—if inflammation set in—to hesitate to deliver must be considered the most mischievous vacillation. Every hour spent in these useless efforts of the uterus, only increases your difficulty—only renders the operation more hazardous—and diminishes your chances of success, because the application of the forceps to parts already inflamed must contuse them to a certain extent, and if so, the contusion will terminate in slough. Although agreeing with Dr. F. Ramsbotham in many of the principles of his practice, I find myself opposed to him on this question. The summary of symptoms which he gives to authorise the use of the forceps, when labour does not continue twenty-four hours, seems to me to be founded upon a principle very hazardous to the safety of the mother. He states, “If, then, the pains are subsiding gradually, or have entirely disappeared; if the strength is failing, the spirits sinking, the countenance becoming anxious; if the pulse be one hundred and twenty, one hundred and thirty, one hundred and forty, in the minute, the tongue coated with white slime, or dry, brown, and raspy; if there have been two or three rigors; if, on pressing the abdomen, there is great tenderness of the uterus; if there be green discharge; if there be preternatural soreness of the vulva, with heat and tumefaction of the vagina; if the head have been *locked* for four hours, and made no progress for six or eight hours; if the patient be vomiting a dark, coffee-ground-like matter; if there be hurried breathing, delirium, or coldness of the extremities, *then we are warranted in having recourse to the forceps, although the labour have not lasted the limited period of twenty-four hours, or even twelve*; and we should be acting injudiciously to allow the case to proceed until the last four symptoms appear, without relief being offered.”* Dr. F. Ramsbotham enumerates these symptoms, to authorise the delivery of the locked or impacted head—a case in which I have already stated that I do not think the forceps can be at all safely employed; but to apply the instrument when inflammation has advanced to such an extent as to engage the constitution in an irritative fever, and only to deliver before the last four symptoms of exhaustion appear, and that, too, when the head is impacted, seems to me to be dangerous in the

* Ramsbotham's Ob. Med., p. 313 and 314.

extreme. I am more anxious, therefore, to impress upon you the importance of not waiting, or withholding your assistance, the moment such symptoms commence; but if, unfortunately, it should happen that they have advanced to the degree so well described by Dr. Ramsbotham, then the safety of the mother must be your first consideration, and you should select the operation that will best secure it. With this view, I again repeat, perforation is your only resource, although I admit it to be a very painful alternative. When you have determined upon the necessity for delivery by the forceps, and the time for performing the operation, the same preliminary arrangement should be made as in the former instance, using still greater caution in your antiphlogistic measures. Hence, if the vagina be swollen and hot, the urine retained, the pulse quick, depletion, some time before operating, would be advisable, the urine being, of course, removed. If the ear be felt, the pubic blade may be passed in the same manner as in the preceding operation, but if not, the presentation must be carefully examined. You can usually trace the lambdoid suture passing upwards from the posterior fontanelle; direct the pubic blade along this, and it will guide it to the ear. You may also take the rule, with regard to the pelvis, laid down by Dr. Rigby, and introduce the first blade behind the trochanter, still bearing in mind its relation to the lambdoid suture: thus the first step of the operation can be generally taken successfully. The passage of the sacral blade is rather more difficult. It may be introduced in the same manner as in the former instance, but its advance is frequently checked as it approaches the brim of the pelvis. If such should happen, be very careful not to use force in pressing it forwards. It is better to act with the pubic blade, for a short time, as a vectis, and if the head advance even slightly, the opposite blade will frequently glide into its place.

When the forceps is applied, it is well to dislodge the head from its situation in the first instance, because it constantly happens that in these cases of arrest some accidental displacement of the head is the cause of difficulty, which the uterus cannot alter; but when the head is relieved, it will glide into the correct position, and may be delivered without difficulty. If, however, you find that with the following pain the head is still arrested, the forceps must be seized firmly, and, in order to

secure your hold, a coarse napkin might be placed loosely round the handles. A steady and powerful traction should be maintained so long as the pain continues, and when it ceases, the grasp of the instrument must be at once released, and remain so until the succeeding pain, when the same steady traction may be renewed. Thus you will generally succeed in bringing the head through the opposing part of the pelvis, and as it advances more easily and approaches the perinæum, again recollect to leave it as much as possible to the efforts of the uterus. While the head is thus drawn through the pelvic cavity, you should bear in mind the direction in which it must pass; that when the forceps is in the axis of the pelvic cavity, the shank of the handles would lie between the tubera ischii, but when the head is in the hollow of the sacrum, the handles would then be directed forwards towards the pubis. You should therefore first draw, with a waving motion, directly towards you, and as the head advances, direct the handles forwards. It is necessary also to observe the rotation of the head in its lateral direction from the oblique towards the antero-posterior measurement of the pelvis. At the same time, it is



Application of Dr. Radford's forceps when the head is fixed in the brim, etc.

advisable rather to follow than to guide the direction of the head in its progress, because, as it descends, it will *naturally* change its position, which might be prevented by the operator's awkwardness in holding the forceps, and attempting too hastily to turn it.

The operation when the head is fixed in the brim of the pelvis differs from either of the preceding operations. The blades are applied over the occiput and face of the child, and not over the ears. This may easily be done in the case to which we have confined this application of the forceps; but it would appear to us extremely difficult and dangerous to do so in other deformities of the brim of the pelvis. Two fingers, and as much as possible of the right hand, should be passed behind the trochanter, towards the centre of the ilium, on the superior side of the pelvis, and if the anterior fontanelle be felt distinctly, the longer blade of the forceps (if they are unequal) should be passed over the fontanelle to the face of the child; the shorter blade may then be passed in the opposite direction over the occiput, guiding it by the lock of the introduced blade. When properly applied, the handles look downwards and backwards towards the perinæum, and in the axis of the brim; traction must be made in this direction, and when the pain commences, the handles of the instrument should be held, as in the former case, firmly, and the force gradually increased, according to the resistance. Two or three steady trials will generally succeed in extricating the head from the brim, when it will rapidly advance without assistance; the forceps might be removed, but it is safer not to do so, lest any impediment might delay its further progress.

It is necessary to remember the change in the direction of the head when it is passing into the hollow of the sacrum. The blades of the forceps might also be changed, so as to direct them over the ears of the child. Great caution is required in extracting the head from the brim of the pelvis, because it is impossible to use any force without compressing the handles strongly together. There is, therefore, danger lest the face of the child be bruised, or, what is quite as likely, the frontal bone bent in. In some cases, the edge of the blade of the forceps has been buried in the frontal bone, even when the instrument was intended to pass over both ears. Such a fatal accident would be much more

likely to occur in this operation, if sufficient care were not taken to avoid it.

If the *posterior fontanelle* be felt on the superior side of the pelvis, the longer or facial blade of the forceps should be passed along the opposite side to the ilium, and then the occipital blade behind the trochanter, as in the former operation. The head of the child always lies in the transverse measurement of the brim, with the occipito-frontal axis corresponding to it. There are only two positions: one with the face to the superior, and the second with the face to the inferior, side of the pelvis. The operation in the second case is, therefore, just the reverse of that in the first position.

In the construction of instruments, some forceps are made especially for this operation. It is considered objectionable (and I think justly so) to operate with a forceps that has the blades of equal lengths, because when the instrument is applied, the occipital blade will prevent the facial passing sufficiently far over the face; its extremity may only reach the nasal bones, or be applied over the frontal sinuses, and, therefore, the bone might be crushed by the force employed in extraction. To avoid this, Dr. Davis has contrived a forceps with unequal blades, in such a manner that the curvature of the longer blade could be diminished or increased if necessary. More lately, Dr. Radford, of Manchester (who has had extensive experience in these cases of deformed pelvis), has invented a forceps with unequal blades for the same purpose. (Vide fig. 12, p. 247.) The majority of practitioners, however, employ only one kind of forceps for these two operations, which they call the *long forceps*. Dr. F. Ramsbotham's long forceps (vide fig. 9, p. 247) has a shank between the handles and the blades, and is so curved as to adapt itself to the axis of the brim of the pelvis. Dr. Rigby has introduced Brünigshäusen's long forceps,* (vide fig. 10, p. 247), an instrument somewhat longer than Dr. Ramsbotham's, having the second curve describing a larger circle. The lock between the handles also is differently constructed. The late Dr. Hamilton used a forceps about fourteen inches in length, with the second curve describing a

* "The most perfect lock is that of Professor Brünigshäusen, of Würzburg, first introduced by ourselves into this country, and commonly known by the instrument-makers as Professor Naegele's forceps."—*Rigby*, p. 137.

smaller circle than that of Dr. Ramsbotham, but it is very clear that Dr. Hamilton never employed it in this operation. (Vide fig. 13, p. 249). Dr. R. Lee observes: "When Dr. Hamilton was in London, twelve or thirteen years ago, the use of the *long forceps* was one of the subjects of conversation which I had with him; and I was astonished when he informed me that he had entirely, for some time, laid aside the short forceps. On further inquiry, however, it appeared that in *no case did he ever use the long forceps until an ear could be felt.*" *

I believe that Dr. Hamilton was not the only obstetrician who had spoken of and described, in very flattering language, *these* kind of long forceps operations. Before you venture to undertake such an operation, the utmost care should be taken in making a vaginal examination, lest you mistake the kind of deformity that prevents the head descending.

I cannot give you a stronger illustration of this than by quoting the first case of instrumental delivery reported by Dr. Lee, in his *Clinical Midwifery*: "On the 28th June, 1828, I (Dr. R. Lee) was present at the delivery of a woman aged thirty, who had been in labour nearly three days and nights, under the care of a midwife. It was the first child. The orifice of the uterus was not fully dilated, and it was very rigid; the vagina swollen and tender, the abdomen tense, and painful on pressure; tongue loaded, urgent thirst, countenance flushed, pulse rapid and feeble. The labour pains for ten or twelve hours had been gradually becoming more feeble and irregular. The head of the child was strongly compressed and much swollen, and the greater part was above the brim of the pelvis. *An ear could not be felt*, and the hollow of the sacrum was empty. It was determined by the practitioner who had charge of the case to attempt to deliver with the long forceps, and he observed, before proceeding to introduce the blades, that it was a case in which the superiority of the long over the short forceps would be observed in a striking manner, and that in less than a quarter of an hour the delivery would be safely and easily completed, and the life of the child preserved. The blades of the forceps were, however, introduced with great difficulty, and still greater was experienced in getting them to lock. Strong traction

* Lee's Theory and Practice of Midwifery.

was then made for several minutes, and the blades slipped off the head. *This happened several times* ; but the attempt to deliver was not abandoned till the operator was exhausted with fatigue. The head was then perforated, and extracted with the crotchet. Violent inflammation and sloughing of the vagina followed, and about three weeks after delivery it was ascertained that a large vesico-vaginal fistula existed." *

Dr. Lee, on this case, observes : " This was the first time I ever saw the forceps applied in actual practice ; and I was struck with the vast difference which exists between the application of the forceps to the head of an artificial fœtus put into a phantom, and the head of a living child. I was led to suspect what I now witnessed—that a dangerous degree of boldness and hardihood might readily be acquired by long practice upon a phantom, where this was not combined with attendance on cases of difficult labour." †

I feel the strongest conviction of the truth of these sentiments ; and for this reason would urge upon you the importance of closely observing those every-day cases which present to you no difficulties. It is only by educating your sense of touch so as to perceive the relations between the head and the pelvis, that you will detect any deviation from their ordinary proportions, or can estimate accurately the amount of disproportion that exists. It is by the same tact that you know with certainty the manner in which the forceps is applied, or can judge of the propriety of its application. It is only by such previous education you can hope to perform any instrumental delivery with success, and, therefore, the study of the presentations in these ordinary, but too often neglected, cases of natural labour, is a far safer mode of acquiring skill in the application of the forceps, than practising on those clumsy imitations of Nature that are commonly employed, under the fanciful names of "phantom," "mannikin," "dolly," etc., etc.

A very few observations only are required on those cases where the position of the head in the pelvic cavity is altered. *When the face is towards the pubis*, the ear can still be felt, and therefore the forceps can be applied in the manner already de-

* Lee's Clinical Midwifery, p. 6, 7.

. † Op. cit. p. 7.

scribed. In this case, it is still more necessary to dislodge the head, and to endeavour to rotate it into the correct position. If this cannot be done, the perinæum must be very carefully guarded from injury as the head is descending, because the pressure upon it is so great. When the face presents, the forceps may also be applied over the ears, and when the traction is made with the instrument, the handles should be directed downwards and backwards towards the perinæum, in order that the chin may the more readily be brought under the arch of the pubis. As soon as this is accomplished, they should be held in the axis of the vagina while the head is passing over the perinæum. In both these varieties, there is great danger of laceration of the perinæum.

In our description of these different operations, no allusion has been made to the os uteri—we have assumed that it is fully dilated. This is admitted by all practical writers to be essential to any such operation. It is unnecessary, therefore, to dwell upon such a point. We may conclude this subject in the words of Denman—"Before the completion of the first stage of labour—that is, before the os uteri be completely dilated, and the membranes broken—the use of the forceps can never come into contemplation, because the difficulties before occurring may depend upon causes which do not require their use, or, if required, they could not be applied with safety or propriety before those changes were made."

LECTURE XII.

OBSTETRIC OPERATIONS.

Operations to save the Mother only—Perforation—Importance of Auscultation—Conditions that authorise Perforation where the Child is alive—Mode of operating when the Head is fixed in the Brim or impacted in the Cavity—Comparative Merits of the Crotchet, and the Craniotomy Forceps—Operation when the Head is above the Brim, or can with difficulty be forced into it—Elizabeth Sherwood's case—Dr. Davis's Osteotomist—Baudeloque's Cephalo-tribe—Sigaultian Operation—Cæsarian Section—Induction of Premature Labour.

HAVING described those operations which are calculated to preserve the lives of the mother and the child, we must now turn our attention to those which are intended to save the mother only. When the head is so impacted in the pelvic cavity that it would be too hazardous to her safety to attempt delivery by the forceps, the alternative that remains is to perforate the head, to remove as much of the brain as possible, and to extract the child by means of the crotchet or craniotomy-forceps. So serious an operation requires the most mature consideration, especially if the child be alive; but should the death of the child take place before symptoms of danger to your patient present themselves, the operation may be undertaken without hesitation, because it is one much less calculated to injure the soft parts of the mother than that with the forceps, and it is more easily performed.

If, then, there be the least suspicion that the case may terminate in perforation, you cannot be too watchful in observing the symptoms. Your attention should be directed to two objects: first, to control, as far as possible, the inflammation which may arise; and, secondly, to observe carefully the pulsations of the fœtal heart. When you have heard them distinctly, observed the variations in their character, and find that they have ceased,

the operation may then be performed. But it would be advisable not to perforate immediately on their cessation, lest you might be deceived. It is preferable to wait for a short time, if the constitutional symptoms admit of it, and to examine again before you proceed to deliver. The greatest difficulty connected with this operation is *the time* when it must be performed. This is especially the case when the child is alive, and symptoms of inflammation are progressively advancing to a dangerous point. Fortunately these cases are rare; but when they do occur, the practitioner is placed in a dilemma: either he destroys the child—an expedient which he must have a natural repugnance to adopt—or if he attempt to deliver by instruments not destructive to it, he runs the risk of exposing the mother to the most serious dangers, without any certainty that he will succeed even in delivering the child, much less in saving its life. It is well, therefore, to consider the progress of such a case.

The head being tightly jammed in the pelvic cavity, a tumour is very rapidly formed on the presenting part. The vagina is hot, swollen, painful, at first dry, but afterwards moistened with an acrid serous discharge. The urethra is compressed, and the urine retained. The uterus is contracted about the child in the interval of the pains. If the fœtal heart should cease, the child may be removed, and these symptoms subdued without injury to the passages; but if the child be alive, there is still a hope, although certainly a distant one, that it may advance when the head is sufficiently compressed by the pelvis, and the swelling of the passages diminished by the antiphlogistic measures adopted. We are not authorised, therefore, to open the head, until there is some evidence of danger to our patient. It is here, then, that your difficulties begin. To pursue these symptoms: the inflammation, which had commenced in the vagina, extends to the uterus, and engages the constitution in an irritative fever. The surface of the uterus felt through the abdomen is not only hard, but very tender; the patient will not bear to have it touched. When the pains return, her agony is extreme, but in the interval she has no respite from suffering; she still complains of pain and soreness; the expression of her voice is altogether altered; the deep groan of the bearing pain is exchanged for a constant whine; a yellow, oily, offensive discharge flows from the vagina. The pulse is febrile; the tongue furred; the countenance pallid

and anxious; respiration laboured, and the stomach highly irritable. She constantly seeks for cold drinks, which are rejected as soon as taken. The symptoms of exhaustion soon follow; the discharges from the stomach are like coffee-grounds. The patient becomes extremely restless, tossing about the bed, and calling for air. Sudden chills on the surface alternate with clammy perspirations; the temperature of the vagina diminishes, and the *fæces* are discharged involuntarily. Blowing, in the act of respiration, coldness of the vagina and extremities, and the gradual cessation of the pains, immediately precede the death of the patient. In this train of symptoms, one evidence of danger is, that which shows the inflammation to have extended to the uterus. This might lead to softening of its structure, and consequently to its laceration. When you are satisfied that such is the case, you cannot temporize, the child must be removed. Another evidence is inflammation, increasing in the vagina, which may terminate in slough. This is still more important, if the typhoid character of the accompanying fever, with dry, brown tongue, and very rapid pulse, point out its erysipelatous tendency. In such cases, extensive gangrene very rapidly succeeds inflammation, so that sometimes the whole mucous membrane of the vagina has sloughed away.* Large portions, also, of the anterior or posterior walls of the vagina have been completely detached, and frightful openings made between the bladder, vagina, and rectum. To delay or vacillate, with such a risk before you, would be worse than dangerous: it would be criminal. Any such symptoms would fully authorise this operation, although the child be living. The same rule applies to cases of exhaustion.

With these observations, we shall proceed to demonstrate the manner of performing perforation. The position of the patient is the same as in the forceps operation. The preliminary measures to be adopted are similar, but much more attention is required, because it is so necessary to combat inflammation. We must assume, therefore, that the rectum is empty, and that the urine has been previously withdrawn from the bladder. It would be right, however, to examine the bladder carefully above the pubis before operating; sometimes the neck and fundus of the bladder are so compressed between the head and the pubis,

* Dr. Doherty on Adhesions of the Vagina, *Dub. Journ.*, vol. xxi. p. 66.

that if a short catheter be used, the urine is only removed from the lower segment, while the principal portion remains above. When this is the case, the head should be pressed back as much as possible, and a long gum-elastic catheter, of the size that will pass the urethra, introduced; great care should be taken not to use force in its passage, lest the instrument perforate the canal.

Having secured these essential points, two fingers of the right hand should be passed to the most depending part of the tumour that presents. The whole surface of the presentation should be carefully examined, in order to determine the degree of dilatation in the os uteri. If not fully dilated, you should observe its exact relation to the tumour,* and the distance of the edge of the os uteri from its centre; and if the os uteri be very thin and closely embraces the tumour, you should accurately define its margin, lest the degree of dilatation should escape your notice. Having thus made a very cautious examination, let the forefinger of the right hand remain applied to the centre of the presenting part, but rather to the pubic side, and with the left hand introduce the perforator, having the point (which is slightly curved) resting upon, and guarded by, the forefinger. When both fingers meet, the right hand may be withdrawn from the vagina, and the perforator forced through the tumour and bone, at the part where the forefinger had been applied. The stops arrest its further progress. The handles should then be separated, in order to break open the cranium; but care is required in doing so. The first two fingers of the left hand should rest on the stops of the instrument, to prevent the perforator slipping from the opening when the handles are raised, and to save the vagina from injury. With the common perforator, one handle is generally held by the operator, while the opposite is raised by an assistant; but if Naegele's perforator be used, an assistant is not required for this purpose. When the bone is sufficiently broken in one direction, the handles may be changed to the opposite, and a crucial opening made. The perforator should be passed into the cranium, and the brain completely

* This tumour, formed on the head by the pressure of the cervix of the uterus, is different from that which afterwards takes place when the head is in the cavity of the pelvis.

broken up. It may then be removed, but it would be advisable to have the forefinger of the right hand still within the opening, because it frequently happens, when the pressure of the head is very great, the bones so overlap each other, that the opening is again closed, which makes it difficult to introduce the crotchet. This latter step of the operation is often very troublesome, especially if the crotchet be properly curved, because the point of the instrument is so directed that it is difficult to get it introduced. In order to do so, the handle should be directed backwards towards the coccyx, when, with a little careful management, it may be passed through, and the remainder of the brain removed. As soon as this is accomplished, the handle of the crotchet should be held firmly in the right hand, while two or three fingers of the left are applied to the bone externally, partly for the purpose of protecting the vagina, if the crotchet should perforate, but chiefly to prevent it breaking through the bone. Some caution also is required in extraction. The force should be so applied that the bone may be held tightly between the flat part of the point and the fingers, without directing it much on the point itself. In this manner you may proceed, never using more force in extracting than is actually required. As the point of the crotchet frequently slips from its place, it is an advantage to have as much of the bone below it as possible, in order that it may not escape from the opening into the vagina; hence the advantage of perforating rather on the pubic side of the head than the lowest part of the presentation.

We have thought it necessary to detail to you the steps of this operation more particularly, because so much of your success depends upon the manner in which it is performed, and especially as respecting it there seems to be much misconception. This is evident in the construction of these instruments. You will find some crotchets having the shaft as straight as the handle, with the point sharp and spear-shaped. No doubt a crotchet of this kind could be fixed within the cranium without any difficulty, neither would there be the least danger that it would slip from its position; but as soon as the extracting force is applied, being directed completely on the point, there is great risk that it would break through the bone. When the crotchet is properly curved, this is altogether avoided, because, while the point is sufficiently fixed in the cranium to retain its hold, the principal force is


employed in compressing the bone against the fingers. These disadvantages induced the late Mr. Holmes and Dr. D. Davis to propose the craniotomy-forceps as a substitute—the revival of an instrument described in the early history of midwifery as the forceps. The craniotomy-forceps is formed of two blades or shafts, one of which has at the extremity a number of teeth, the other a corresponding number of openings, so that when the blades are applied together the teeth rest in these openings like sockets. Sometimes the shafts are separate, and may be united like the forceps, by a lock; others are joined together by a hinge-joint. The handles are strong, and the extremity of the blade is smooth and rounded. When the craniotomy-forceps is used, the blade, which is dentated, is introduced within the cranium, while the opposite blade is passed outside; when they are closed, the teeth perforate the bone, and are received by the openings mentioned, and thus the cranium is grasped firmly by the instrument. Extraction may then be made without difficulty, provided the bone does not give way. The craniotomy-forceps may be very easily introduced and applied to the head of the child; the teeth are guarded by the construction of the instrument; and there are no other sharp points to tear the vagina when applied, neither is there the same management required in extracting as there is with the crotchet. Its advantages, therefore, seem to be such as to make it supersede the use of the crotchet. Nevertheless, some of the most experienced practitioners dislike the instrument, because it is liable to one great objection from which the crotchet is free. The bone is so crushed by the craniotomy-forceps, that when extraction is made, it tears away just so much of the bone as was broken, and leaves the head behind. This is especially the case when the child has been dead some time, and putrescency has commenced. Thus the presenting part of the head may be torn up into fragments without being disturbed from its position. If the instrument fail in its purpose, there is great difficulty in applying the crotchet afterwards, and there is some danger lest the spiculæ of bone broken by the forceps should tear the passages: a lacerated wound of this kind may give rise to very serious inflammation, and therefore increase the hazard of the operation. To obviate this objection, some forceps are made without teeth, having only a strongly serrated edge in the blades, by which the bone is

grasped, and prevented from slipping. There is less risk of breaking through the bone with an instrument of this kind, and it is, therefore, less objectionable; but still there is more danger of separating the parietal bone at the sutures than with the crotchet, if the resistance be great. These observations on the comparative merits of the crotchet and craniotomy-forceps give you the result of my own experience in the use of these instruments. The objections stated have also occurred to other practitioners, in whose judgment I have the highest confidence, and therefore I feel the more certainty in their truth. Nevertheless, I should not wish you to infer that the craniotomy-forceps is either a useless instrument or one that should be altogether discarded from practice. On the contrary, in many cases, and especially in those where the head is much ossified, it may be employed with great advantage, if caution be used, and the instrument be properly selected. You should only conclude, from what has been said, that either instrument may be employed usefully in cases especially fitted for their application, but that in the majority of instances in which these kind of difficulties require the aid of instrumental assistance, it is safer to employ the crotchet.

In the description we have given of perforation, your attention has been directed to the operation *when the head of the child has either entered or passed through the brim of the pelvis*, and has there become impacted; but cases occasionally arise that present much greater difficulties even than these. There are instances *where the head cannot enter the brim of the pelvis*, in consequence of its extreme deformity; and in determining the mode of delivery, we are again involved in a cloud of controversial opinions, through which it is difficult to find out the true course to pursue. The object of one of these questions is to determine the limit of perforation, or, in other words, to decide what is the least possible space in the pelvis through which a child may be extracted by the crotchet. Secondly, when, in cases of extreme disproportion, this seems hardly to be accomplished, and at the same time, the attempt is extremely hazardous to the mother, the question arises, whether it should be at all attempted? When the result is so doubtful, so far as the safety of your patient is concerned, and the destruction of the child is certain, the advocates of the Cæsarian section claim for it the advantage that there is at least

a chance of preserving the child, while the risk to the parent can hardly be greater, and therefore they condemn any attempt to perforate under such circumstances.

Time would not permit us to enter into the discussion of these questions. It will be sufficient to point out to you the nature of them, and the different operations which have been suggested to meet the difficulty. The first question sprung up in the discussion of a very remarkable case that occurred in the practice of Dr. Osborne—the case of Elizabeth Sherwood, a cast of whose pelvis is before you. Dr. Osborne states, that “she was so deformed both in her spine and lower extremities, as never to be able to stand erect for one minute without the assistance of a crutch under each arm.” At the age of twenty-seven years, however, she became with child, and was admitted a patient into Store-street Hospital. A vaginal examination was made, and “immediately upon the introduction of the finger, I [Dr. Osborne] perceived a tumour equal in size, and not very unlike in feel, to a child’s head. However, it was instantly discovered that this tumour was formed by the basis of the os sacrum, and last lumbar vertebræ, which, projecting into the cavity at the brim, barely left room for one finger to pass between it and the symphysis pubis, so that the space from bone to bone at that part *could not exceed three quarters of an inch*. On the left side of the projection, quite to the ilium, which was about two inches and a half in length, the space was certainly not wider, and, indeed, by some of the gentlemen who examined her afterwards, it was thought to be rather narrower. On the right side, the aperture



Ovate Pelvis of Elizabeth Sherwood.

was somewhat more than two inches in length from the protuberance to the ilium, and, as it admitted the points of three fingers (lying over each other) in the widest part, it might, at the utmost,

be about an inch and three quarters from the hind to the fore part, but it became gradually narrower, both towards the ilium and towards the projection.”* Such was the pelvis through which Dr. Osborne determined, after consultation with Drs. W. Hunter, Denman, Bromfield, Walker, and Watson, to extract the child with the crotchet. “It was my duty (he proceeds) to perform the operation, which I began about eleven o’clock that night; after placing her, in the usual manner, close to the edge of the bed, on her left side, as the situation most commodious both for the patient and myself. Even the first part of the operation, which in general is sufficiently easy, was attended with considerable difficulty and some danger. The os uteri was but little dilated, and was awkwardly situated in the centre and most contracted part of the brim of the pelvis. The child’s head lay loose above the brim, and scarce within reach of the finger, nor was there any suture directly opposite to the os uteri. Having desired an assistant to compress the abdomen with sufficient force I introduced them (the perforating scissors) with the utmost caution through the os uteri, and, after repeated trials, at length succeeded in fixing the point into the sagittal suture near the posterior fontanelle. I very soon, and with great facility, penetrated into the cavity of the head, destroyed the texture of the cerebrum, with a common spoon extracted a considerable quantity, and breaking down the parietal bones, made an opening sufficient for the free discharge of what remained. In this state we left her”† for thirty-six hours, when, “upon examination, a small portion of the head was found squeezed into the pelvis; indeed there were some little detached bits of the parietal bones lying loose in the vagina. . . . Our intention, by delaying the extraction of the child six-and-thirty hours after opening the head, was, in the first instance, to allow the uterus opportunity, by its continued contractions, to force the head as low and as much within reach of the crotchet as the nature of the case admitted, and afterwards, to induce as great a degree of putrefaction as possible in the child’s body, by which means it would become soft and compressible, and afford the least possible resistance in its extraction. . . . I immediately deter-

* Osborne’s Essays, p. 242.

† Op. cit. p. 247.

mined to begin to make an attempt to extract the child. *I call it an attempt, for I was far from being satisfied in my own mind of the practicability.* . . . The os uteri being situated as before described, in the most contracted part of the brim of the pelvis, where the space was incapable of permitting the introduction of the curved point of the crotchet, without great difficulty and danger, my first endeavours were bent to draw the os uteri with my finger into the widest part of the brim of the pelvis, and to dilate it as much as possible. Both the removal of the os uteri, and such dilatation of it as the bones admitted, were effected without much trouble. I then introduced the crotchet, through the perforation, into the head, and by repeated efforts, made in the slowest and most cautious manner, destroyed almost the whole of the parietal and frontal bones, or the whole upper or presenting part of the head: and as the bones became loose and detached, they were extracted with a pair of small forceps, to prevent, as much as possible, laceration of the vagina in their passage through it. The great bulk of the head formed by the base of the skull, still, however, remained above the brim of the pelvis, and from the manner in which it lay, it was impossible to enter without either diminishing the volume, or changing the position: the former was the most obvious method, for it was a continuation of the same process, and, I trusted, would be equally easy in execution. I was, however, most egregiously mistaken and disappointed, being repeatedly foiled in every endeavour to break the solid bones which form the basis of the cranium, the instrument at first invariably slipping as often and as soon as it was fixed, or at least before I could exert sufficient force to break the bone. At last, however, by changing the position of the instrument, and applying the convex side to the pubis, I fixed the point, I believe, into the great foramen, and by that means became master of the most powerful purchase that the nature of the case admitted. Of this I availed myself to the utmost extent, slowly, gradually, but steadily, increasing my force till it arrived to that degree of violence which nothing could justify but the extreme necessity of the case, and the absolute inability, in repeated trials, of succeeding by gentler means. But even this force was to no purpose, for I could not perceive that I had made any impression on that solid bone, or that it had been the least advanced by all my exertions. I be-

came fearful of renewing the same force in the same way, and therefore abandoned altogether the first idea of breaking the basis of the cranium, and determined to try the second by endeavouring to change the position. . . . I therefore again introduced the crotchet in the same manner, and fixing it in the great foramen, got possession of my former purchase; then, introducing two fingers of the left hand, I endeavoured with them to raise one side of the fore part of the head, and turn it a little edgeways. Immediately and easily succeeding in this attempt, the two great objects were at once accomplished, for the position was changed, and the volume diminished. Continuing my exertions with the crotchet, I soon perceived the head advance, and, examining again, found a considerable portion of it had been brought into the pelvis. Every difficulty was now removed, and by a perseverance in the same means for a short time, the remaining part of the head was brought down, and out of the os externum.”*

We have detailed this case to you more at length, because it accurately describes an operation with the crotchet different from what we have described—one by which the vault of the cranium is quite broken up and removed, and the base of the skull is drawn obliquely through the contracted brim of the pelvis, the crotchet being fixed in the foramen magnum. It is also remarkable—we might say singular—in the fact, that a child could by any means be drawn through a pelvis so extremely distorted as to have the antero-posterior measurement reduced to three-quarters of an inch; and perhaps the most astonishing fact connected with the case was, that the woman recovered without a bad symptom, and sat up in seven days after such an operation. It is not surprising, therefore, that a warm controversy should have arisen; the advocates of the Cæsarian section condemning the attempt, and protesting against such a case being made a precedent for future crotchet operations; while the supporters of craniotomy claimed the case as a valuable proof of the superiority of perforation to hysterotomy, which latter was generally fatal to the parturient woman.

* Op. cit. p. 255.



Dr. Davis's Osteotomist.

Dr. D. Davis contrived several instruments to meet the difficulty of these cases—one was the *osteotomist*, a strong bone forceps intended to cut completely away the bones which form the vault, so as to leave the base of the cranium. Dr. Davis stated that this also may be broken and removed by the instrument, so as to prevent the necessity for that violent exertion which Dr. Osborne was obliged to have recourse to in drawing the cranium through the brim of the pelvis.*

Dr. Davis also contrived a double crotchet, for the purpose of extracting the body of the child, after the head, in this mutilated state, has been brought through the pelvis, and thus he anticipated that the osteotomist would “enable skilful operators to effect deliveries in cases of moderate distortions with much more facility to themselves, and proportionally less danger to their patients, than heretofore, but it will also have the effect of reducing almost to zero, the necessity of having recourse to that last extremity of our art, and the forlorn hope of the unhappy patient—the Cæsarian operation.”†

M. Bandeloque, jun., also, has invented an instrument for a similar purpose to the osteotomist, to break up the head, not by

* Dr. Campbell has invented an instrument—the *kephalepsalis*—for a similar purpose as Dr. Davis's *osteotomist*.

† Dr. Davis's *Obstetric Medicine*, p. 857.

cutting it away, but by crushing it together. *The cephalo-tribe* consists of two very strong blades, rough on the inside, and having handles, through which a screw passes. The handles are brought forcibly together by turning the screw, and the blades, by the same power, crush the bones that lie between them (vide fig. 22, p. 251).

Both these operations are intended to supersede the Cæsarian section, and both are liable to the same objection—viz., the extreme difficulty in using them in those cases where they are chiefly required, as well as the danger to which the passages would be exposed in such an operation, especially with the cephalo-tribe. In moderate distortions, where the crotchet or craniotomy-forceps may be used, the *osteotomist* and *cephalo-tribe* are unnecessary, because they possess no advantages which would lead us to employ them in preference to the former instruments. But when the distortion is extreme, and the alternative is the Cæsarian section, from the impossibility of delivery by the crotchet, they would be invaluable, if, as Dr. Davis expected, they could reduce hysterotomy to zero. Both instruments are new, and future experience must decide the question, but *prima facie* evidence seems against them. Look at the cephalo-tribe, and ask yourselves, How could that instrument be used in Elizabeth Sherwood's case? To me it seems impossible. Again, with regard to the osteotomist, it must be passed into the uterus, above the brim of the pelvis, and that part of the head within reach of the instrument cut away by it, until the bones are all removed. We question very much whether this could be done under the circumstances supposed, but admitting it to be possible, the difficulty of applying the crotchet to the broken cranium, lying loosely above the brim, must be very great. And if we fail, how are we to act? Are we, then, to have recourse to the Cæsarian section, for the purpose of delivering a mutilated child from the uterus? It is true, we are assuming a mal-adroit performance of the operation, which might arise from want of skill. It is right, however, to do so, and to consider the alternative in cases of failure, if we would properly appreciate the value of the improvement. This objection will, perhaps, appear with more force, from the caution used by Dr. Osborne to avoid such a difficulty in his operation. He first perforated the head, and then allowed his patient

to remain thirty-six hours in strong labour, in order that some part of the head (then a putrid mass) might be driven into the brim of the pelvis. He preferred leaving the patient so long in labour, under such unfavourable circumstances, rather than operate, while the head was yet above the brim of the pelvis. For these reasons we very much doubt whether the sanguine expectations of Dr. Davis will ever be realized.

Long before these instruments were invented, another operation was proposed, to supersede the Cæsarion section, which at first was attended with some success, received the approval and honours of the Academy of Medicine, Paris, excited the warmest enthusiasm in its favour, and has now become only a part of obstetric history. Sigault and Le Roy proposed to *divide the symphysis pubis*, and thus to force open the contracted brim. It is sufficient to say that this operation failed in its object, and proved to be so dangerous to the patient, that it has been discarded from practice; we shall not, therefore, dwell upon it, but proceed to the Cæsarion section itself.

This operation is based upon the third principle we have stated to you—viz., when, from the circumstances of the case, the safety of the mother seems to be more than doubtful, if not hopeless, the child must, if possible, be saved. *The Cæsarion section* is therefore indicated in those extreme cases. In the case of Elizabeth Sherwood, although the crutch succeeded, its success was the wonder of the professional world, and of none more than Dr. Osborne himself. It cannot, therefore, be taken as a rule to guide your practice. In order to decide upon the Cæsarion section, you should weigh carefully the probable result to the mother if the operation be not performed; and if it appear to you that perforation is impracticable, or so difficult to perform that the danger seems to be nearly as great to the patient as opening the uterus, you are then authorized to undertake the operation, because, if there be a probability that perforation will not ensure safety to the mother, you are certainly bound to consider the child, and to give it a reasonable chance for its life. If, for instance, the ratio of mortality from hysterotomy as compared with craniotomy, in these extreme cases, were equal, or as four to three, this slight difference would not, it appears to me, be sufficient to justify you in destroying the child. Taking this view of the operation, it would

be desirable to ascertain by statistical results its precise value, because no estimate of the operation can be formed from individual cases. It is difficult, however, to do so from reports, unless a careful attention is given to the circumstances under which the operation was performed. For instance, the results of the operation in Great Britain and on the Continent differ exceedingly. In the former, more than three-fourths of the patients died, and more than one-half the children were lost. In the latter, the mortality of the mothers was much less, and more children were saved. But in Great Britain the operation was performed, as a *dernier ressort*, after the patient had been several days in labour, and under the most unfavourable circumstances, while on the continent, it was generally undertaken in the first instance. The errors on the one side arose from unnecessary hesitation and delay, and I fear we must add that, on the other, there have been also some mistakes from precipitancy, and some needless operations performed.

Dr. Churchill has given the results of 409 cases of Cæsarian section :—

1st. Among British practitioners, in 40 cases, 11 mothers recovered, and 29 died, or nearly 3-4ths.

2nd. Out of 37 cases where the result to the child is mentioned, 22 were saved, and 15 were lost, or 1 in $2\frac{1}{2}$.*

3rd. Among Continental practitioners, out of 369 cases, 217 mothers recovered, and 152 died, or about 1 in $2\frac{1}{2}$.

4th. Out of 187 cases where the result to the child is given, 138 were saved, and 49 were lost, or nearly 1 in 4.

5th. Taking the entire number, which amounts to 409, we find that 228 mothers were saved, and 181 were lost, or about 1 in $2\frac{1}{2}$; and that out of 224 children, 160 were saved, and 64 lost, or 1 in $3\frac{1}{2}$.

More lately, Keyser, of Copenhagen, has applied himself to the same question; he has carefully examined and checked the accuracy of previous statistical researches on the Continent, and has arrived at a result differing slightly from that of Dr. Churchill. 338 cases are collected, of which 128 proved successful, and 210 unsuccessful. The mortality of the mother is, therefore, 0·62. The same data give 0·31 as the mortality of the children.

* Churchill's Operative Midwifery, p. 221.

Keyser has taken great pains to classify the cases according to the different circumstances under which the operations were performed, and has thus ascertained a remarkable fact respecting the degree of contraction in the pelvis:—" *The mortality was less in those cases in which it was extreme than in those in which it was more limited.* It was 0·47 in cases which were $1\frac{1}{2}$ (French) inches and less than this, and 0·66 in cases which were $1\frac{3}{4}$ and more than that measurement."* These latter cases are those in which we have stated that the operation is called for, and therefore, in the Continental practice, where it is performed under the most favourable circumstances, we may assume the mortality to the mother to be about 1 in 2, to the child 1 in 3. I do not think any fair conclusion can be drawn from British practice, because there were so many causes in operation against its success; neither can we ascertain the proportionate mortality where perforation is performed in these extreme distortions. We cannot, therefore, derive any rigid conclusion from a comparison of the results of both operations. But if the circumstances of the case be such that the risk to the patient is increased much beyond ordinary perforation, I do not think we should venture upon that operation in preference to the Cæsarian section. In the former case, the child must be sacrificed for a very doubtful advantage. In the latter, there is every reasonable chance of preserving the child, while the mother has at least an equal chance that she will recover. For these reasons we are not disposed to look upon the Cæsarian section with that horror with which some practitioners view it; nevertheless, the serious nature of the operation should be strongly impressed upon your minds, and every caution made use of that is required in capital operations. The strictest antiphlogistic measures should be previously used, to prevent inflammation, and the same means as to temperature, etc., adopted, that have been found so useful in those ovarian operations which have lately occupied public attention. The most essential point, perhaps, to attend to is, *the time* at which the operation is undertaken. It would be advisable to allow some time to elapse after labour commences, to satisfy yourself that the head cannot enter the brim of the pelvis, and thus to confirm

* London and Edinburgh Monthly Journal, No. LV., p. 542.

your previous diagnosis. At the same time, it would be highly improper to allow labour to proceed to such a length, as to hazard either inflammation, exhaustion, or the death of the child. It is for this reason that the operation has so often failed in British practice. In this respect, each case must be considered separately; but you may take it as a rule, admitting of many exceptions, that having previously ascertained, by examination *per vaginam*, the extreme distortion of the pelvis, if after the first twenty-four hours the head does not enter the brim, the operation may be performed. We have Keyser's evidence on this point, also, to prove that such delay would not be injurious to the patient:—"Regarding the time which intervened from the commencement of labour, M. Keyser divides his tables into three categories. In the first, the operation had been performed within the first twenty-four hours; in the second, in the interval between the twenty-fifth and seventy-second hour; in the third, more than seventy-two hours after the commencement of labour. In the first category, the mortality of the mothers was 0·67, that of the infants, 0·28. In the second, the result was respectively 0·55 and 0·33, and in the third, the mortality amounted to 0·72 and 0·60."* From these tables it appears that the middle period, between twenty-four and seventy-two hours, was less fatal to the mother. An operation of so grave a nature should never be undertaken without the aid and assistance of at least two professional men of reputation, if it be at all practicable to have their support.

The mode of operating we shall briefly explain. The rectum and bladder must be carefully emptied; the position of the placenta ascertained with the stethoscope, and the exact direction of the uterus observed. If it project forwards nearly in the middle line of the body, and the placenta be in its usual position at the back of the uterus, an incision may be made through the linea alba for about seven inches, commencing above the umbilicus and terminating about two inches above the pubis; the uterus is thus exposed, and the peritoneum along with it. An assistant should press with both hands firmly on the uterus, at either side of the wound, while the uterus is being divided. This must be

* London and Edinburgh Monthly Journal, p. 542.

done by cautious incisions, in the direction of the external wound, until the membranes are seen. These should be raised, and a small opening made in them to allow the liquor amnii to escape externally; the whole fluid may be removed by successive applications of sponges to the opening. The membranes should then be divided on a director the whole length of the wound, and while this is being done, a second assistant should be prepared to grasp and remove the child, while the first maintains pressure on the contracting uterus, to prevent as much as possible protrusion of the intestines or exposure of the peritoneum. The placenta then may be easily removed, the intestines replaced (they always protrude), and the wound united by several sutures; water-dressing and a broad bandage may be applied over the whole.

Lauvergat advised us to puncture the membranes previous to the operation, and this plan has certainly many advantages. The placenta can be heard much more distinctly; the size of the uterus is reduced, and the calibre of its vessels diminished; the amount of hæmorrhage may thus be lessened, and the external wound need not be so large. The peritoneum, also, is less likely to be exposed when the uterus contracts after the child is removed.

The dangers you have to apprehend from this operation are:

1st. The shock to the constitution, under which the patient may sink;

2nd. The hæmorrhage which may result from the operation; and

3rd. The inflammation of the peritoneum, in consequence of the sac being opened.

The subsequent treatment we shall consider, under these different heads in another part of the course.

If we have pointed out to you with sufficient clearness the different obstetric operations, the degrees of disproportion to which they are applicable, from the slightest to the most extreme,—from simple delivery by the vectis to the Cæsarian section,—if we have succeeded in defining the limits of each operation, we would now direct your attention to a means of obviating the necessity for those operations that involve the sacrifice of the child.

In cases of contracted pelvis, which prevent the passage of the full-grown child, if labour should take place at the seventh in place of the ninth month, you can readily perceive the possibility of the child (then much smaller) being safely delivered. The child is quite capable of supporting respiration at that period; therefore it has been proposed—and the practice is now very generally adopted—to *induce labour to take place* at the seventh month, when we are satisfied that a living child cannot be born at the ninth month.

The induction of premature labour is one of the greatest improvements in modern practice, because by its means the leading principle of obstetric operations may be carried out, and both mother and child preserved, in cases in which otherwise we could hardly hope for such a result. We shall not occupy your time with by-gone discussions on the propriety of prematurely forcing labour; it is sufficient to say, that its propriety—nay, its necessity—is admitted in the cases which we have described to you, and the only point to be determined, is the case in which the operation is required. We must recollect that, independently of other objections, we have a strong reason for not inducing premature action of the uterus if it can be avoided. The uterus is not prepared for such a change: the cervix is still unfolded, the connection between the uterus and the placenta is more intimate, the circulation in the uterus less easily diverted into other channels; consequently, you expose your patient to greater risk than at the conclusion of pregnancy, and thus you would not be justified in doing without a sufficiently powerful motive. The safety of the child is your justification; but you must have clear proof that it is in danger. You cannot trust to an examination of the pelvis only, because, unless distortion is great, it would be premature to say that the child cannot be delivered. The most certain evidence is the result of previous labours, and in the diseased pelvis you have generally sufficient proof of its necessity. Perforation may have been performed in the previous labour; or with every successive labour the contraction of the pelvis may have increased, so as to render the last more difficult than that which preceded it. If, in such a case, the previous delivery were completed with much difficulty by the forceps, you may fairly assume that the next will require perforation. Thus, you

will generally have sufficient evidence to guide you in these cases ; but remember the induction of labour is not suitable in first pregnancies.

Different modes of exciting the action of the uterus have been proposed :—1st. *By direct irritation*, as frictions over this organ, artificial dilatation of the os uteri with the fingers, or by the introduction of a sponge tent ;—2d. *By the specific action of ergot of rye* ;—3d, and lastly, *By deranging the connexion between the uterus and the ovum*, either by detaching the membranes from the sides of the uterus, or puncturing the membranes and allowing the liquor amnii to escape. Of these means the last is the most certain, but, at the same time, one which it would be preferable to avoid if other means were efficient for the purpose, because the liquor amnii would ensure a more favourable dilatation of the uterus, and the child be more secure. Ergot of rye is unsafe, because of the child, the preservation of which is your only motive for interfering ; therefore, artificial dilatation by a sponge tent may be first tried, and if it fail, the membranes may be ruptured with a stilette. The action of the uterus sometimes commences immediately, but it may not begin for twenty-four or forty-eight hours after the operation.

LECTURE XIII.

OBSTETRIC INSTRUMENTS.

THE VECTIS: History of—Invented by Roonhuysen, De Bruyn—Secret purchased by Visscher, and Van de Poll—Denman, Bland, Aitken, Lowder, Gaitskill.—THE FORCEPS: Invented by Dr. Paul Chamberlen, and used extensively by him and his Sons, Hugh and Peter—Forceps contrived by Giffard, Chapman, Gregoire—Smellie's Improved Forceps, the Parent of those at present used—Varieties in the Construction of different Forceps—Objects intended by them—Difference in the Principle adopted—Dr. Davis's Forceps, Dr. Denman's, Dr. Beatty's, Dr. Ziegler's.—THE PERFORATOR: Smellie's Scissors, Denman's, Naegele's—Holmes's Perforators—Crotchet—Craniotomy Forceps—Conclusion.

FROM the details of those instrumental deliveries, we wish now to draw your attention to the instruments themselves, and, giving you briefly their history, to point out the alterations and improvements which have been made in them. First, we shall speak of *the vectis* and *the forceps*.

Previously to the introduction of either of these instruments, the state of midwifery was such that it was sufficient that a labour was difficult, to assume the death of the child. No other operation was known or practised but perforation, and when a midwife (the accoucheur of that day) called for assistance, it was generally because the patient was in danger of her life. The medical man who undertook the operation of delivering the child, did so under the most unfavourable circumstances: the child must be destroyed, and perhaps the parent was not saved by the operation. It is not surprising, therefore, that the obstetrician (or man-midwife, as he was called) should have held a very humble rank amongst his professional brethren—like the plague, destruction seemed to follow in his path—and, consequently, he was an object rather to be shunned than sought after. Operative midwifery was in this

condition when, in the seventeenth century, two practitioners, one in Holland, the other in this country, contrived instruments by which delivery could be accomplished, and the child's life at the same time preserved. One of those inventors was Roonhuysen, a Dutch practitioner, who invented the vectis—the other, Dr. Paul Chamberlen, the inventor of the forceps. The introduction of steam did not produce a greater revolution in the commercial world than did these instruments in obstetric practice. Deliveries were effected safely that before would have been despaired of, and, not unlike our illustration, with a great economy of time. The reputation of both men soon rose to the highest pitch, but I regret to add they did not elevate their profession to the same degree. Governed by mercenary motives, the invention was kept secret by both, and all the aid that mystery could give was employed to magnify its importance. It served the intended purpose; the practice of both increased to the fullest extent, and consequently the number of patients delivered by these instruments was considerable. De Bruyn, one of Roonhuysen's pupil's, admits having delivered eight hundred women with the vectis; and from this you may form an estimate of the general number of operations. In fact, the practice of midwifery assumed a new character: formerly parturition was left to the efforts of nature, and very often far beyond what prudence would dictate. If she failed, and the patient was in danger of sinking under the inefficient efforts of the uterus, the child was dragged away by hooks and crotchets, in whatever way the practitioner could best accomplish it. Now, the principle was changed; art pushed nature aside; delivery became a question of mechanical skill; and, in these times, the principal merit of Roonhuysen and Chamberlen was, that by their invention they could not only deliver a woman where the natural efforts failed to do so, but also they could effect the delivery in a much shorter time than nature could generally accomplish, even where assistance was not so obviously required. Dr. Hugh Chamberlen boasted that "by this manual operation [the forceps] a labour may be dispatched (in the least difficulty) with fewer pains, and sooner"—than nature could—"to the great advantage, and without danger, both of woman and child." *

* Churchill's Operative Midwifery, p. 74.

These secret means of delivering women were sold from one to another, like patent medicines, until at length the secret made its way into the profession.

* We have stated that Roonhuysen invented the *vectis*, the use of which he taught to his son Roger, to Ruysch, and to Böckelman. They instructed De Bruyn, and, at length, after the secret had passed through three generations, two Dutch practitioners, Jacob de Visscher and Hugo Van de Poll, influenced by the true spirit of science and philanthropy, purchased the secret from De Bruyn's daughter for 5,000 livres, and at once made it known to the world. "Roonhuysen's lever consisted of a flat piece of iron, bent into a slight curve at both ends, and he generally employed it covered with soft leather." *



Roonhuysen's Vectis.

This simple contrivance was soon improved upon on the Continent by Titsing, Morand and Herbiniaux. Dr. Denman states, "that when the vectis was first known in this country [England], that described by Heister was preferred to those recommended by the surgeons of Amsterdam. The vectis used by Dr. Cole was like one blade of the forceps, somewhat lengthened and enlarged. That of Dr. Griffith was of the same kind, with a hinge between the handle and the blade; and that of Dr. Wathen was not unlike Palfyn's, but with a flat handle, and a hook at the extremity of the handle, which prevented its slipping through the hand, and might be occasionally used as a crotchet. Many other changes have been made in the construction of the instrument, but the vectis now generally used is of the following dimensions:—The whole length of the instrument before it is curved is $12\frac{1}{2}$ inches; the length of the blade before it is curved, $7\frac{1}{2}$ inches; the length of the blade when curved, $6\frac{1}{2}$ inches; the widest part of the blade is $1\frac{3}{4}$ inches. The weight of the vectis is $6\frac{1}{2}$ ounces. The handle is fixed in wood."† Dr. Bland, who was, equally with Denman, an advocate for the vectis, used an instrument much straighter, and one which could not be employed otherwise than as a lever.

* Churchill, p. 87.

† Denman's Introduction, p. 286.

Dr. Bland's vectis had the advantage of facility in the introduction; Denman's of security in its purchase on the head. Dr. Aitken, of Edinburgh, wished to combine both advantages, and invented what he called the living lever, from its motion resembling that of the finger. A screw was fixed in the handle, by turning which the blade might be curved to any extent. This instrument soon fell out of use; the mechanism was not sufficiently strong to keep the blade curved when any force was employed in extraction. Lowder's vectis is made with a hinge-joint between the blade and the handle; it can thus be carried very conveniently—unfortunately, too conveniently—in the pocket; there is, therefore, a great temptation to misapply the instrument. It had been in very general use when Mr. Gaitskill introduced an improved vectis. He says: "The vectis should be thirteen inches long; one-half to form the handle—the other the curve. The handle should be made of hard wood, rendered rough, for the purpose of obtaining a firmer hold, and made to screw on and off. When the instrument is made with a hinge-handle, it is very difficult to introduce; therefore, this construction of the instrument should never be adopted." *



Lowder's vectis, with hinge-handle.



Gaitskill's vectis, (handle screwed on.)

In the variety of these instruments you will observe a striking difference in their curvature: some are nearly straight, like Roonhuysen and Bland's; others, like Lowder's, are very much curved. The principle of the former is the lever; of the latter, the tractor. The mode in which Gaitskill applied the vectis has been already explained; and the instrument, in its construction, is adapted to his manner of using it; but it cannot be so conveniently employed in the way we have recommended. The

* London Medical Repository, pp. 823, 80, 81.

curve of the blade is too abrupt, and the blade itself rather too wide, to introduce conveniently on the pubic side of the pelvis. If the curvature were less and the blade narrower, it would be more suitable for the purpose indicated.

The forceps was invented by Dr. Paul Chamberlen, somewhere about the year 1650 (the precise date is uncertain); it was kept a secret from all, except his sons Peter and Hugh, for more than sixty years, and at length, in the year 1716, its principle transpired through some channels that have not been correctly ascertained.

We have already stated the boast of Dr. Hugh Chamberlen, that "by God's blessing and their own industry, his father, brother, and himself, had attained to, and long practised, a way to deliver women in this case, without any prejudice to them and their infants." Their success had led Dr. Hugh Chamberlen to calculate rather too confidently on the powers of the instrument; he seemed to think it omnipotent. In the year 1670, he went to Paris for the purpose of selling his secret, and, it is said, had offered it for 10,000 crowns to the first physician of Louis XIV. It was necessary, however, to determine its value; and a case of deformed pelvis soon after presented itself to Mauriceau. The woman had been altogether eight days in labour: on the fifth day the os uteri was fully dilated, but the narrowness of the pelvis was such, that the head could not advance. Mauriceau was again sent for at the end of three days more, and "declared to all the assistants that the delivery could not be effected; of which (says Mauriceau) they being fully persuaded, urged me to draw the child from the belly by the Cæsarian operation, which I would not undertake, knowing well that it is always very certainly mortal to the mother. *But after I had left the woman, without being able to afford her relief,* there arrived, unexpectedly, an English physician, named Chamberlen, who was then in Paris, and who, from father to son, practised midwifery in London, where he has since acquired the highest reputation in this art. This physician, seeing the woman in the condition I have described, expressed his astonishment that I, whom he pronounced and affirmed to be the most dexterous accoucheur in Paris, could not deliver her, and promised that he would do so in less than half a quarter of an hour, *whatever difficulty he might encounter.* He accordingly went to work,

and laboured upwards of three hours, without stopping to take breath, and then being thoroughly exhausted, and seeing the poor woman almost dead, he was compelled to abandon the case, and avow the delivery could not be effected, as I had declared. The woman died, undelivered, twenty-four hours after; and I found, on opening the body, which I did by performing the Cæsarian operation after death, that the whole uterus was torn and pierced in several places by the instruments which this physician had employed blindly without the guidance of his hand, which, being one-half larger than mine, could not be introduced.”* Chamberlen left Paris in disgust, or, as Mauriceau says, “he returned to England in a few days, seeing clearly that there were men in Paris more skilful in the art of midwifery than he.” The case quoted does not, however, prove any great skill on either side. Chamberlen may have killed the patient by his violence in using the forceps, but Mauriceau left her for eight days in labour, and, for all that we are informed to the contrary, would have allowed her to die without relief. The case, however, proves the indiscriminate use which Chamberlen made of the forceps. His secret was not sold, like Roonhuysen’s, so that the instruments invented by him remained unknown for a very long time.

Dr. Churchill states, “About this time [1716], or soon after, the secret appears to have been communicated to one or two, for Dr. R. W. Johnston, when speaking of the forceps, says, ‘Besides these, I have a pair of forceps which did belong to the late Mr. Drinkwater (late surgeon and man-midwife at Brentford), who began practice in 1668, and died in 1728. The size and form of this pair agree with those of Chapman and Giffard, save only that the hooks of the handles are turned outwards.’ And Mr. Chapman, in 1733, published a description and a plate of the instrument which he had used from the year 1726, stating it to be the instrument used by the Chamberlens, but without stating whence he procured it”† (vide fig. 7, p. 247). We have now sufficient proof that Chapman’s forceps was quite different from the Chamberlens’, and, consequently, the only way in

* Mauriceau, vol. ii. pp. 23, 24; Lee’s Lectures, p. 201.

† Churchill’s Op. Mid. p. 112.

which the secret was revealed to them was by an explanation of the principle, and not by any exhibition, of the instrument. Chamberlen's forceps was discovered by mere accident many years afterwards. It happened in this manner, as described by Mr. Causardine. The estate of Woodham Mortimer Hall, near Malden, was purchased by Dr. Peter Chamberlen, some time previous to 1683, and continued in his family till about 1715, when it was sold by Hope Chamberlen to William Alexander, wine merchant. In the year 1818 (more than a century afterwards), in an old chest, found in one of the chambers of this house, certain obstetric instruments were discovered, along with old coins, trinkets, gloves, fans, spectacles, etc. Mr. Causardine's description of these instruments is as follows:—"First we have a simple vectis with an open fenestrum; then we have the idea of uniting two of these instruments by a joint, which makes each blade seem as a fulcrum to the other, instead of making a fulcrum of the soft parts of the mother, and which also unites a power of drawing the head forward. This idea is, at first, by a pivot, which, being riveted, makes the instrument totally incapable of application. Then he goes to work again, and having made a notch in each vectis for a joint, he fixes a pivot in one only, which, projecting, is to be received into a corresponding hole in the other blade, after they have been applied separately. It may be observed, that although there is a worm in the projecting part of the pivot, yet there is no corresponding female screw in the hole to receive it. Every practical accoucheur will know that it is not easy, nor always possible, to lock the joint of the forceps with such accuracy as to bring this pivot and hole into opposite contact. This Chamberlen soon discovered, and next produced a more light and manageable instrument, which, instead of uniting by a pivot, he passes a tape through the two holes, and winds it round the joint, which method combines sufficient accuracy of contact, security, and mobility"* (vide fig. 1. p. 245).

The instruments here described were altogether different from those afterwards brought forward by Chapman and Giffard, about the year 1730. Chapman was the second practitioner

* Med. Chir. Trans. vol. ix. p. 183.

who gave lectures on midwifery in London. In them, he explained the new instrument, and the mode of its application. The forceps he employed resembled in the shape of the blades those at present in use; the handles were of steel and hooked at the extremity, the blunt hooks being turned inwards. The blades were united by a kind of mortice-lock, just like the blades of a pair of scissors. Giffard's were similar. Soon after Chapman published his account of the forceps, the inventive genius of the profession was busily employed in making improvements in it. Some of these, as Burton's, did not deserve the name, being rather more clumsy even than the original instrument. Others were only alterations, without the introduction of any new principle. But in 1752, Smellie's work appeared, and to him we are indebted for the forceps that is the basis of all the modern instruments. The practice before his time is best described in his own words:—"The common way of using them [the forceps] formerly, was by introducing each blade at random, taking hold of the head any how, pulling it straight along, and delivering with downright force and violence, by which means both the *os internum* and the *os externum* were often torn and the child's head much bruised. On account of these bad consequences, *they had been altogether disused by many practitioners*, some of whom endeavoured, in lieu of them, to introduce *divers kinds of fillets* over the child's head, but none of them can be so easily used, or have so many advantages, as the forceps, when rightly applied and conducted according to the directions that shall be laid down in the next section."

"Mr. Chapman, as mentioned in the introduction, was the first author who described the forceps, with the method of using them; and we find in the observations of Giffard several cases in which he delivered and saved the child by the assistance of this instrument. A forceps was also contrived at Paris, a drawing of which may be seen in the Medical Essays of Edinburgh, in a paper communicated by Mr. Butter, surgeon; but after Mr. Chapman had published a delineation of his instrument, which was that originally used by the Chamberlens, the French adopted the same species, which, among them, went under the denomination of Chapman's forceps. For my own part, finding in practice that, by the directions of Chapman, Giffard and Gregoire, at Paris, I

frequently could not move the head along without contusing it, and tearing the parts of the woman (for they direct us to introduce the blades of the forceps where they will easiest pass, and, taking hold of the head in any part of it, to extract, with more or less force, according to the resistance), I began to consider the whole in a mechanical view, and reduce the extraction of the child to the rules of moving bodies in different directions. In consequence of this plan, I more accurately surveyed the dimensions and form of the pelvis, together with the figure of the child's head, and the manner in which it passed along in natural labours; and, from the knowledge of these things, I not only delivered with greater ease and safety than before, but also had the satisfaction to find, in teaching, that I could convey a more distinct idea of the art in this mechanical light than in any other, and particularly give more sure and solid directions for applying the forceps, even to the conviction of many old practitioners, when they reflected on the uncertainty attending the old method of application. From this knowledge, too, joined with the experience and hints which have occurred and been communicated to me in the course of teaching and practice, I have been led to alter the form and dimensions of the forceps, so as to avoid the inconveniences that attend the use of the former kinds." Smellie introduced two very obvious improvements in the forceps, which have since been retained, and may be considered the essential attributes of the British forceps.* He first proposed the present mode of locking the forceps,† and also had the handles made of wood, in place of iron. (Vide fig. 2, p. 245). These alterations have not been followed in France. The forceps of Gregoire differed from Chapman's only in the iron handles; Gregoire's had their hooked extremities turned outwards; the method of locking was the same: these are still retained in Levret's forceps, which has been for many years in general use in France. So that Gregoire's forceps may be considered the parent of the French—Smellie's of the British forceps.

From the date of Smellie's forceps to the present day, these leading characters have, with a few exceptions, been continued

* Smellie's Midwifery, vol. i., p. 250—252.

† Dr. Rigby attributes the introduction of the lock to Chapman


in the construction of the British instruments, but they are almost the only points in which they agree. The varieties in other respects are almost endless: no two instruments are alike; and the zeal with which new changes are proposed and defended, the superiority which each inventor claims for his favourite instrument, and the constant introduction of some slight alteration, just sufficient to establish the parentage of the new forceps,—this desire for novelty would lead us to suppose that a new forceps was an essential introduction to practice. It would be only a tedious continuation of the history of this instrument to mention in detail all the different forceps proposed, or employed, with the names of the inventors; it is more desirable to point out to you the varieties suggested in the different parts of the instrument, and the objects which they are intended to accomplish.

The length of the forceps is generally about 11 inches— $4\frac{1}{2}$ for the handles, $6\frac{1}{2}$ for the blades. Those intended for the high operation exceed this. The late Dr. Hamilton's forceps (vide fig. 13, p. 249) was $13\frac{1}{2}$ inches in length; Brüninghäusen's, introduced by Dr. Rigby, 13 inches (vide fig. 10, p. 247); Dr. Radford's long forceps, $13\frac{1}{4}$ inches (vide fig. 12, p. 247). Those which are less than 11 or 12 inches are only intended to be used when the head is resting on the perinæum. This length is generally preferred by those practitioners who object to use this instrument for the purpose of shortening a labour which may be much prolonged. Aitken's short forceps (vide fig. 4, p. 245), Dr. Collins' (vide fig. 6, p. 245), and Denman's (vide fig. 3, p. 245), are about ten inches in length.

The length of the handles is very different, even in forceps which are made for a similar purpose; for instance, the handles of the long forceps used by Dr. Rigby are fully 6 inches; Dr. Radford's, only 3 inches. It is obvious that the greater the length of the lever, the greater the power gained; therefore, where much power is sought for, the handles must be long; when too great power is dreaded, the handles are made very short. *The length of the blades* is more uniform, being generally between 6 and 7 inches. The blades of Denman's forceps are about $6\frac{1}{2}$ inches; Haighton's long forceps, 8 inches in length. (Vide fig. 8, p. 247.) Some instruments have a shank between the handle and the blades, so that the lock may be quite external to the vulva, and the soft parts saved from injury in locking the blades. The blades,


however, differ extremely in their shape: some are straight, having only a single curve outwards from the lock; others, especially the long forceps, have a second curve backwards, to correspond with the axis of the brim of the pelvis. The latter was first suggested by Smellie and Pugh, and has been adopted by Drs. R. W. Johnston, Osborne, Haighton, Hamilton, Rigby, Ramsbotham, and others. The degree of the lateral curvature is very important. Some are made to embrace the head without compressing it; when such is the intention, the distance between the centre of the blades, where they are most apart, is about 3 inches, the distance between the extremities of the blades $1\frac{1}{2}$ to $2\frac{1}{2}$ inches, or what is just sufficient to prevent them slipping off the head. When compression is intended, the greatest distance between the blades is about $2\frac{1}{2}$ inches, that between the extremities half an inch, and sometimes even less. The *fenestræ* also vary: some are long, narrow and pyriform, in order that the blades may be the more easily introduced or withdrawn from the pelvis; other instruments have the *fenestræ* wide (Dr. Davis's remarkably so) and oval-shaped, so as, by the greater breadth of the blade, to embrace the head more completely, and at the same time to allow its widest part to pass through the blades. A few have no *fenestræ*. The blades are generally fixed immovably in the handles; but in Dr. Conquest's long forceps, one blade is screwed on, like Gaitskill's vectis. Dr. Hamilton and Dr. Davis made the handle of one part of the forceps moveable, in the same manner as Lowder's vectis. The object of these contrivances is to prevent the length of the handle interfering with the introduction of the instrument, which is sometimes the case with the long forceps, unless the patient is placed very much over the side of the bed. The exceptions to the manner of locking the blades are met with in the forceps recommended by Dr. Rigby, in which one blade has a fixed pivot, the other a notch which fits into it. The late Dr. Beatty contrived a forceps with a transverse opening in the shank of one blade, through which the other passed. (Vide fig. 16, p. 249.) Dr. Ziegler, of Edinburgh, has proposed another modification of the instrument, which I am informed is much used in Scotland. The *fenestra* of one blade is carried down to its handle, and, in introducing the instrument, this elongated *fenestra* is slipped over the handle of the other single blade

which is first introduced, and thus serves as a guide to the second.



Dr. Ziegler's Forceps.

These are the principal varieties introduced in the construction of the instrument. We have endeavoured to explain the objects intended by them. Among so many, each differing from the other, and all used by accoucheurs of acknowledged skill, it is no easy matter to determine which forceps is the best. In making a selection, therefore, of any, we must not be understood as wishing to depreciate the value of those we reject. In the construction of these instruments, two different principles seem to have been followed. Some have wished to render the mechanism of the forceps as perfect as possible; others have sought simplicity in its construction. The former have contrived instruments of great power, but which are not very easily applied; the latter have succeeded in the facility with which their forceps may be applied, although with some of them the power is extremely limited. Your late respected professor, Dr. Davis, contrived a forceps which you may take as an example of the former. In its mechanism it is perfect. It is exactly fitted to the head of the child, so as to lie close without bruising it; the fenestræ are made very wide, in order that the parietal protuberances may pass through them, and the blades are curved very much backwards, in order to correspond with the axis of the brim. If it be applied



Dr. Davis's Forceps.

to the head of the still-born child, the perfect manner in which the head is embraced is quite obvious ; therefore, when it is accurately passed over the head of a living infant arrested in the pelvis, there cannot be a greater extracting power ; but in order to accomplish these mechanical advantages, simplicity is sacrificed, and it is not easy either to introduce or to withdraw the blades. When the ear is near the pubis, and you wish to pass one blade over it, the fenestræ are too wide to allow it to pass behind the pubis ; the blade must first be passed along the plane of the ischium, and then brought gradually round to the ear at the pubis ; which is certainly inconvenient, and to the young practitioner may be embarrassing. Again, the second curve from before backwards, although receiving the support of many very experienced operators, seems to me to be liable to a similar objection. You may in your hurry introduce the wrong blade first ; or if you avoid (and of course you will be careful to avoid) this mistake, the curved blade passing in a direction different from the handle, you cannot be so certain that it is correctly applied ; on the other hand, the advantage claimed for the curve seems to me one founded much more on theory than practice. These objections only apply to the instrument in the hands of the inexperienced accoucheur ; but when, by patient attention, you have acquired that practical tact which is so necessary in this branch of your profession, and which experience will ultimately give you, you


will also acquire with it that facility in the application of the forceps which will render it a matter of indifference which you select, provided the instrument is at all properly constructed.

But, in commencing your career, we must advise you to avoid all complications in instruments—the simpler they are, the safer they will be in your hands. For this reason, we prefer forceps of the latter class, such as have been used by Smellie, Denman, and others, who employ the straight forceps, with the blades narrow, and the fenestræ pyriform. These can always be easily introduced or withdrawn; the handle is a perfect guide to the direction in which the blade is passing, and the only point which it is necessary to secure is, that the blades do not slip when extraction is made. This cannot take place if the blades be properly curved: their curvature is one of the most essential points in the construction of these forceps. Their extremities should be sufficiently close to retain their position on the head without bruising the face, and, at the same time, the space between the blades should be sufficiently wide to prevent much compression of the cranium.

Experience can alone decide between the merits of similar instruments on a point of this kind; and having used several of these forceps, that which I have found to be the best, and one which I am disposed to recommend to you, is one used by Dr. Beatty, Professor of Midwifery to the College of Surgeons in Ireland, and described by him in the *Dublin Journal*, vol. xxi. The whole length of the forceps is $12\frac{1}{2}$ inches, the distance between the extreme points of the blades $1\frac{1}{8}$ inch; the greatest distance between the blades is 3 inches; the breadth of the blade $1\frac{3}{8}$ inch.* This instrument is well calculated for that operation which I have described as being intermediate between the operations with the long and short forceps, viz., when the head is in the cavity of the

* These measurements are those given by Dr. Beatty, in the *Dublin Journal*, but the instrument which is in my possession, and which I have found so useful, varies a little from this. The entire length is eleven inches and a half; the handle, four inches and a half; blade, seven; greatest breadth between blades, three inches; between extreme points, one inch and an eighth. To this instrument I have added a shank to the blades about one inch in length, when it is necessary to introduce them high within the pelvic cavity. It is right to state that the length is taken in a direct line from the extremity of the handle to the extremity of the *curved* blade..

pelvis, without touching the perinæum. When the head is resting on the perinæum, a shorter forceps would be more convenient.



Beatty's Forceps.

Beatty's Forceps modified.

nient—for instance, that of Dr. Collins, which is about 10 inches in length. (Vide fig. 6, p. 245.) But when the head is fixed in the brim of the pelvis, the forceps of Dr. Radford, having unequal blades, appears to be the most suitable. (Vide fig. 12, p. 247.) In giving you this account of an instrument of so much importance as the forceps, we wish to avoid leading you into what seems to us a great error, viz., a belief that by mechanical skill in the construction of the instrument, a great deal more can be accomplished by it than what is really the case. Hence every practitioner has his favourite forceps, and no little boast is sometimes made of the deliveries accomplished by it. But we would beg of you to remember that Smellie and Denman with their simpler forceps had as much success as Hamilton and Davis, with their more ingenious instruments, and *that the success of an operation depends much more on the hands that use the forceps than on the instrument itself.*

It is necessary that the forceps should be sufficiently well formed to embrace the head conveniently, without bruising any part. It should be capable of being easily locked and unlocked, the blades introduced or withdrawn without difficulty, and the handles of sufficient length to hold the forceps firmly, but no

more; you should avoid making the length of the handles a means of increasing the power of the instrument. If you secure these points in the forceps you select, you have gained every advantage the instrument can give you; for the rest, your success must depend upon your previous education.

It is unnecessary to dwell at any length upon those instruments which are used for perforation and extraction of the head. This operation was the only one known before the forceps was invented, and at first was never attempted until the child was dead; it was then drawn away by "hooks and crotchets." If the head were too large, it was opened by some pointed instrument (Raynald used a penknife), and the hook placed inside. In course of time these instruments were brought to a more definite form, and reduced to the present instruments employed.

The first attempt at a *perforator* was made by Sir Fielding Ould, who invented an instrument which he called "*terebra occulta*." The point was rounded, had a cutting edge, and was concealed within a sheath, to prevent it injuring the passages. In 1752, Smellie proposed a scissors with a short cutting edge on the outside, terminating in a blunt stop (vide fig. 19, p. 251). By this means he intended, not only to perforate the bone, but also to cut away the broken fragments. Denman modified this instrument to its present shape. He retained the external cutting edge, widened the stops, to prevent them passing within the opening, and made the internal cutting edge a flat surface. Thus Smellie's *scissors* became Denman's *perforator* (vide fig. 21, p. 251).


The *perforator* remained unaltered from Denman's time until the present day, when Naegele suggested an improvement in the hinge-joint. In Denman's *perforator*, it is necessary to open the handles for the purpose of opening the blades, so as to break the bone. This is often very inconvenient, and an assistant is generally required for the purpose. Naegele, however, altered the hinge-joint in such a manner, that by pressing the handles together the blades were opened; thus, with Naegele's *perforator*, one hand can accomplish what requires two with Denman's. In order to prevent the handles of Naegele's *perforator* closing too soon, while being introduced, a straight steel rod passes from one to the other, which can be very easily removed; but, if the catch between the rod and handle be not nicely fitted,

if it be too tight or too loose, this object may be defeated ; and if the handles are not kept quite apart while the perforator is being introduced, the points separate from each other too soon (vide fig. 20, p. 251). The late Mr. Holmes endeavoured to obviate these accidents—first, by altering the handles, removing the steel rod, and changing their direction, so that, by pressing fully against them, the blades were kept closed ; but lest they might open, the point of the perforator is attached only to one blade (vide fig. 23, p. 251).



Crotchet, having the correct curve.

The crotchet has undergone but little alteration. That invented by Mesnard, and adopted by Smellie, is still very generally used. We have already stated to you the importance of having the shaft of the instrument properly curved, and not, as is often the case, made quite straight, which is only suitable for that extreme operation which we have described in Elizabeth Sherwood's case, and which you should not attempt. Very lately, Dr. Churchill has introduced some improvements in this instrument (vide fig. 25, p. 251). The handle is placed at right angles with the shaft of the instrument, something like the handle of a boot-hook, and the point of the perforator is notched, so as to have two short points to hold the bone, in place of one long one, which might pass through it.



Craniotomy Forceps.

The craniotomy forceps is sometimes employed in preference to the crotchet. We have already stated the objection to that instrument when made as originally designed by Dr. Davis. The strong points of the blade, in forcing through the bone, too

often break it up into fragments, without moving the head. To avoid this, Mr. Coxeter, of Grafton street, has, at my suggestion, made one, by which the bone can be very firmly held without being broken. The surface of the internal blade is strongly serrated with rough lines, which correspond with grooves in the opposite blade, and the bone being held between them can hardly slip from their grasp.

I must now bring to a conclusion this very difficult subject—the management of difficult labours; one which I always feel a reluctance to enter upon, because I cannot go through it without coming into collision with some opposing opinions. In the principles I have laid down I have wished to avoid dogmatism, and have endeavoured to place before you, as impartially as was in my power, the facts upon which the reasoning that led me to them is founded. I think you will find them to be correct, and the safest for you to adopt, when you enter upon practice; but if hereafter, when your experience and manual skill are improved, you find that you can venture on a bolder course than what I have advised, I shall only be too happy to receive the benefit of that experience, and, in return, to be taught by you.

“———— Si quid novisti rectius istis
Candidus imperti; si non, his utere mecum.”

Recollect, however, that you must be cautious how you venture upon “a meddlesome midwifery,” and *that there are no circumstances to justify violence in performing an operation.*

MEASUREMENT OF FORCEPS *a*

WHICH HAVE BEEN CHIEFLY USED IN BRITISH PRACTICE.

NAME.		LENGTH.			BREADTH.			Length of shank.	Second curve.	AUTHORITY.
		Whole.	BLADE.	Of handle.	Greatest between blades	Between points.	Of blade.			
		A long curve.	Direct.							
Chamberlen's forceps		11	7 $\frac{1}{2}$	6 $\frac{3}{4}$	4 $\frac{1}{2}$	2 $\frac{1}{12}$	1	1 $\frac{1}{2}$	0	Measured from } vide the original <i>b</i> } fig. 1.
Giffard's	do.	12 $\frac{1}{2}$	7 $\frac{1}{2}$	6 $\frac{3}{4}$	5 $\frac{1}{2}$	2 $\frac{1}{4}$	2	1 $\frac{1}{2}$	—	Mulder
Chapman's	do.	15	9 $\frac{3}{4}$	9	5 $\frac{1}{2}$	2 $\frac{1}{4}$	0	1 $\frac{1}{2}$	—	Do. vide fig. 7.
Smellie's short	do.	11 $\frac{1}{2}$	6 $\frac{1}{4}$	6	5 $\frac{1}{2}$	2	0	1 $\frac{1}{2}$	—	Do. vide fig. 2.
Do. long	do.	12 $\frac{1}{2}$	7 $\frac{1}{2}$	7	4 $\frac{3}{4}$	2	0	1 $\frac{1}{2}$	—	Do.
Pugh's short	do.	11	7 $\frac{1}{2}$	7	3 $\frac{3}{4}$	2 $\frac{1}{4}$	1	—	2d c.	Do.
Do. long	do.	14	8 $\frac{3}{4}$	7 $\frac{1}{4}$	5 $\frac{1}{2}$	2 $\frac{1}{4}$	1	—	2d c.	Do.
Johnston's	do.	11	6 $\frac{1}{2}$	6	4 $\frac{1}{2}$	2	0	—	2d c.	Do.
Lowder's	do.	11 $\frac{1}{2}$	6 $\frac{1}{2}$	6	4 $\frac{5}{8}$	2	—	—	0	Do.
Osborne's	do.	11 $\frac{1}{4}$	6 $\frac{3}{4}$	6	4 $\frac{1}{2}$	2	—	—	—	Do.
Denman's	do.	10 $\frac{3}{4}$	6 $\frac{1}{2}$	6 $\frac{1}{4}$	4 $\frac{1}{2}$	2 $\frac{1}{2}$	—	—	—	Do.
Do.	do.	10	7	6 $\frac{1}{2}$	3 $\frac{3}{4}$	2	—	—	—	Measured, <i>d</i> vide fig. 3.
Do.	do.	11 $\frac{3}{8}$	0	7	4 $\frac{3}{8}$	2 $\frac{1}{2}$	1	—	—	Ramsbotham <i>e</i>
Haighton's	do.	13 $\frac{1}{2}$	0	8 <i>f</i>	5 $\frac{1}{2}$	2 $\frac{3}{8}$	—	3 $\frac{3}{8}$	2d c.	Radford, <i>g</i> vide fig. 8.
Hopkins'	do.	12	8	5 $\frac{1}{2}$	5	2 $\frac{1}{4}$	—	1 $\frac{1}{2}$	2d c.	Measured, vide fig. 18.
Hamilton's	do.	13 $\frac{1}{8}$	0	7 $\frac{3}{8}$	5 $\frac{1}{2}$	2 $\frac{5}{12}$	—	—	2d c.	Radford, vide fig. 13.
Aitken's	do.	10 $\frac{1}{2}$	0	6 $\frac{1}{2}$	4 <i>k</i>	3	1 $\frac{1}{12}$	1 $\frac{5}{12}$	—	Measured, vide fig. 4.
Davis's	do.	11 $\frac{1}{2}$	0	6 $\frac{1}{2}$	4 $\frac{1}{2}$	2 $\frac{5}{8}$	0	1 $\frac{3}{8}$	2d c.	Davis, vide page 236.
Conquest's short	do.	11	7 $\frac{1}{2}$	6	4 <i>m</i>	2 $\frac{8}{12}$	—	1 <i>m</i>	0	Measured, vide fig. 5.
Do. long	do.	14 $\frac{1}{2}$	0	6 $\frac{1}{2}$	6 <i>n</i>	2 $\frac{1}{2}$	1	2	—	Radford
Ramsbotham's long do.	do.	12 $\frac{1}{2}$	0	8 $\frac{1}{2}$	4 $\frac{1}{2}$	2 $\frac{1}{2}$	0	1 $\frac{1}{2}$	2d c.	Ramsbotham, vide fig. 9.
Brüninghausen's	do.	13	8	7	6	2 $\frac{1}{2}$	—	—	2d c.	Measured, vide fig. 10.
Beatty's, W. (late)	do.	11	7 $\frac{1}{2}$	6 $\frac{1}{2}$	4	2 $\frac{1}{2}$	—	—	0	Do. vide fig. 16.
Beatty's, T.	do.	12 $\frac{1}{2}$	0	8	4 $\frac{1}{2}$	3	—	—	0	Beatty
Do. with shank	do.	12 $\frac{1}{2}$	0	6 $\frac{1}{2}$	4 $\frac{1}{2}$	3	—	1 $\frac{1}{2}$	—	Measured } vide p. 238.
Churchill's	do.	12	8	7 $\frac{1}{2}$	4 $\frac{1}{2}$	2 $\frac{1}{2}$	—	—	—	Do.
Collins'	do.	10	0	5 $\frac{1}{2}$	4 $\frac{1}{2}$	2	1 $\frac{1}{2}$	—	—	Collins, vide fig. 6.
Ziegler's	do.	12 $\frac{1}{2}$	0	5	4 $\frac{1}{2}$	2	1	3	—	Ziegler, vide p. 235.
Radford's long	do.	13 $\frac{3}{4}$	0	10 $\frac{3}{4}$ <i>s</i>	3	2	0	3 $\frac{1}{2}$	0	Radford, vide fig. 12.
Do. the short blade		13	0	10	3	2	0	3 $\frac{1}{2}$	0	Do.
J. Y. Simpson's forceps <i>t</i>		13	6 $\frac{3}{12}$	6	5	2 $\frac{1}{2}$	1 $\frac{2}{12}$	2	2d c.	Measured, vide fig. 14.

a The measurements of these forceps have been taken either from the authorities stated, or from actual measurement, viz., 1. The whole length of the instrument, taken in a direct line from the extremity of the handle to the end of the blade. 2. The length of blade along the curve. 3. The length of the blade in a direct line, from lock to point. 4. Length of handle. 5. The greatest space between the blades. 6. The same between the points. 7. The breadth of the single blade. 8. The length of the shank.

b This forceps seems to have been the last and the most improved of four made by Chamberlen.

c Greatest breadth of blade near the lock.

e Ramsbotham, Principles, etc., p. 288.

g Radford's Essays, "On the Long Forceps," p. 3.

i The handle attached to one blade by a hinge-joint.

k A moveable roller is placed between the handles.

l Fenestræ very wide.

n One handle screwed on to the blade.

o Introduced and used by Dr. Rigby. A pivot lock.

p There is a transverse opening in the shank, through which the opposite blade passes.

q The measurement is here given in one-eighth fractions, to correspond with Dr. Beatty's.

r The blades being unequal, the measurement of each is given. The locks are reversed.

s The measure of the blade includes the shank, which forms a ring.

t Knees above joint to prevent unlocking; joint loose to allow lateral motion.

FIG. 1.

The most improved of five instruments made by Chamberlen, copied from the original.

FIG. 2.

Smellie's forceps covered with leather.

FIG. 3.

Denman's forceps, copied from a fac simile of one in possession of the Dublin Lying-in Hospital.

FIG. 4.

Aitken's forceps. A small roller is fixed in one handle, which is moved by a screw, and is so adapted as to keep the handles separated when the instrument is grasped.

FIG. 5.

Conquest's forceps, with wide fenestræ. The shanks twisted.

FIG. 6.

Collins' forceps, blades straight.




FIG 1.—Chamberlen's Forceps.



FIG. 2.—Smellie's Forceps.




FIG. 3.—Denman's Forceps.



FIG. 4.—Aitken's Forceps.




FIG. 5.—Conquest's Forceps.



FIG. 6.—Collins' Forceps.

FIG. 7.

Chapman's forceps, from which Gregoire's and the French forceps seem to be derived.

FIG. 8.

Haighton's forceps; blades very light and slightly curved backwards; handles long.

FIG. 9.

Dr. F. Ramsbotham's forceps, with shank and second curve.

FIG. 10.

Brüninghäusen's forceps, used by Dr. Rigby.

FIG. 11.

Davis's forceps, with unequal blades.

FIG. 12.

Radford's forceps, with unequal blades; handles short; shank forms a ring.




FIG. 7.—Chapman's Forceps.




FIG. 8.—Haighton's Forceps.



FIG. 9.—Ramsbotham's Forceps.



FIG. 10.—Rigby's do.



FIG. 11.—Davis's do.



FIG. 12.—Radford's Forceps.

FIG. 13.

Hamilton's forceps; blades strongly curved backwards; one handle moveable on the blade.

FIG. 14.

Dr. J. Y. Simpson's forceps. Knees above the joint to prevent them unlocking. Joint so loose as to allow lateral motion to a considerable degree. Handle so as to use the instrument as a tractor rather than a compressor.

FIG. 15.

A modification of Ziegler's forceps, which had been copied from a pair supposed to be Dr. Z.'s. Inventor unknown.

FIG. 16.

The late Dr. W. Beatty's forceps, having a transverse opening in the shank of one blade sufficiently wide to admit the other

FIG. 17.

Churchill's forceps. Blades straight and the fenestræ short. The greatest distance between the blades near the extremity.

FIG. 18.

Hopkins' forceps. Blades having wide fenestræ and second curve; shank forms a ring; handles long.




FIG. 13.—Hamilton's Forceps.




FIG. 14.—Simpson's Forceps.




FIG. 15.




FIG. 16.—W. Beatty's Forceps.




FIG. 17.—Churchill's Forceps.




FIG. 18.—Hopkins' Forceps.

FIG. 19.

Smellie's scissors, with rounded stops, and having the outer edge like the inner, sharp.

FIG. 20.

Naegele's perforator modified. The point is formed by one blade, but so as to leave the other of sufficient length to prevent it escaping from the opening made in the cranium.

FIG. 21.

Denman's perforator. The inner cutting edge of Smellie's made flat, the stops not rounded.

FIG. 22.

Baudeloque's cephalotribe. The blades forced together by the power of a screw.

FIG. 23.

Holmes' perforator. The blades closed by separating the handles. The point attached to one blade.

FIG. 24.

Craniotomy forceps. One blade having transverse ridges adapted to similar grooves on the opposite blade.

FIG. 25.

Churchill's crotchet. Point, double. Handle, transverse.



FIG. 19.
Smellie's Scissors.




FIG. 20.
Naegele's Perforator.




FIG. 21.
Denman's Perforator.




FIG. 22.
Baudeloque's Cephalotribe.



FIG. 23.
Holmes' Perforator.



FIG. 24.
Craniotomy Forceps.



FIG. 25.—Churchill's Crotchet.

SUMMARY

• OF THE

PRINCIPLES AND RULES

LAI D DOWN IN THE PRECEDING LECTURES,

ARRANGED AS APHORISMS,

TO WHICH SOME OF DENMAN'S ARE ADDED.

APHORISMS.

SECTION I.

MECHANISM OF NATURAL LABOUR.

I. Labour is the action of the uterus to expel its contents when the fœtus is sufficiently mature to sustain respiration, p. 55.

II. Labour is called natural "if the head of the child present, if the labour be completed in twenty-four hours, and if artificial assistance be not required" (Denman), p. 56.

III. Labour may be divided into three stages. The first stage is dated from the opening of the os uteri to its complete dilatation. The second stage commences when the os uteri is perfectly dilated and terminates in the expulsion of the child. The third stage is occupied with the expulsion of the placenta, p. 58.

IV. The dilatation of the mouth of the uterus (the first stage) is effected chiefly by the muscular fibres of the fundus acting upon it through the medium of the liquor amnii, which conveys accurately to the os tinæ the whole power of the uterus (pp. 61—64), moderates and equalizes the force employed, and dilates the mouth of the uterus without exciting irritation. The pains caused by this action are called *grinding pains*.

V. The order of uterine action is from the fundus to the mouth of the uterus, p. 64.

VI. When the os tinæ offers any unusual resistance to the fundus, the force of the fundus is not increased, the action is frequently renewed, and may be for a time suspended, in order to diminish irritation, so that the object of nature is to accomplish her purpose by time rather than by force, p. 72.

VII. When the dilatation of the uterus is completed, *the second*

stage of labour begins. There is then a marked difference in the uterine contraction. Not only is the entire force of the uterus employed, but it is aided by the muscles that bound the abdominal cavity. The pains that accompany this action are called “bearing pains,” p. 73.

VIII. The manner in which the head passes through the pelvis in its most usual position, is as follows—

1st. When the head is above the brim of the pelvis, the forehead and the occiput are nearly on the same level, but when the head enters the brim, the occiput descends lower than the sinciput, and glides a certain distance along the plane of the ischium, against which it rests. The forehead then advances more rapidly at the opposite side of the pelvis, until it is arrested by the convergence of the ischium and the shorter sacro-ischiatic ligament. The occiput then again descends obliquely along the ischio-pubic ramus, and emerges with part of the parietal bone beneath the pubic arch.

2d. Simultaneous with this motion, there is a very slight rotation on the longitudinal axis of the head, by which that side of the head which is next to the pubis descends lower than that near the sacrum, so that the parietal protuberance of the pubic side becomes the presenting part.

3d. As the head is so advancing through the pubic cavity, the shape of the pelvis obliges it to pass in a spiral direction: hence the head, which may enter the pelvis in the oblique or transverse measurement, is turned as it descends towards the conjugate axis.

4th. When the head escapes from the outlet, the occiput rests against the ramus of the pubis and ischium, and becomes a fixed point, round which the remaining portion of the head successively passes out. In some instances, the occiput rests directly against the arch of the pubis, and the head is expelled in the conjugate axis of the outlet. According to the more usual course, it passes out obliquely, pp. 74, 75, 76.

IX. There are generally *four* positions of the head in which the parietal bone presents. 1st. Left occipito-cotyloid. 2d. Right occipito-cotyloid. 3d. Left fronto-cotyloid. 4th. Right fronto-cotyloid. Two in which the face presents. 1st. Right mento-cotyloid. 2d. Left mento-cotyloid, pp. 77—79.

X. The four positions of the vertex may be known by the anterior and posterior fontanelles, and by the ear.

XI. When the head begins to press on the perinæum, the mucous secretion is increased in the vagina, the head advances and retires with the pains for some time, until the perinæum is sufficiently stretched. The pains then become more powerful, and force the head through the vulva, there is then the greatest danger lest the perinæum be lacerated, pp. 84, 85.

XII. When the head is expelled, the face turns towards the right thigh in the first position—towards the left thigh in the second position. In the latter case, the arms frequently lacerate the perinæum, p. 86.

XIII. As soon as the expulsion of the child is accomplished, the uterus ceases to contract for some time, the interval varying from five to fifteen minutes; then the contractions are renewed for the expulsion of the placenta. This constitutes *the third stage of labour*, p. 86.

XIV. If the abdominal muscles contract sufficiently, or if artificial support be given to the fundus during the contractions of the uterus to expel the child, the placenta is generally separated at the same time, and may be either expelled from the vulva or into the vagina, p. 87.

XV. If the uterus be without support, the placenta may be retained, either from deficient irritation or irregular contraction. Hæmorrhage does not usually accompany retention caused in this way, p. 88.

SECTION II.

MANAGEMENT OF NATURAL LABOURS.

XVI. The “show” is a slight sanguineous discharge, arising from the ruptured vessels of the mouth of the uterus, when it first dilates. As soon as this takes place, labour properly begins. The patient has now entered upon the first stage, p. 91.

XVII. When called upon to attend a patient, *the summons should be instantly responded to*. Your introduction should not be

abrupt; and if called to a patient for the first time, still greater caution should be exercised, pp. 92, 93.

XVIII. During the early part of the first stage, it is not necessary nor advisable to remain in the room with your patient. It will be sufficient to remain in an adjoining room, until there is occasion to make a vaginal examination, p. 94.

XIX. When the pains have continued for some time regular and frequent, an examination may then be made. The patient, loosely attired in her night-dress, should lie on the bed on her left side, as near to the edge as possible, having the knees drawn up towards the abdomen. You should then wait until the pain returns, and when it is about to cease pass the forefinger of the right hand, anointed with cold cream, or any unctuous substance, within the vagina. Examine whether it be rigid or relaxed, dry or moist. Examine the rectum through the posterior wall, and fundus of the bladder through the anterior. Introduce the middle finger, in order to examine the os uteri. Pass both fingers along the sacral side of the vagina, and when you cannot advance them further, direct them forward towards the pubis; you then feel what seems to be the irregular folds of a flaccid bag, projecting into the vagina: examining this with caution, the edge of the os uteri may be traced; if the finger be passed within it, you will sometimes feel the head; you may often fail, and yet the head present. The dilatability of the os uteri, its direction, and the degree to which it is opened, should be ascertained, pp. 95, 96.

XX. If the membranes are ruptured early, an examination should always be made, lest the funis or the shoulder present. Before the fingers are withdrawn from the neighbourhood of the os uteri, examine the distance of the sacrum; and as they are being withdrawn, ascertain if possible the space in the pelvic cavity, p. 96.

XXI. In order to make a sufficiently careful examination, a little time may be required, during which the pains may return; you should then cease until they subside, noting only those points which have been mentioned; but make it a rule not to withdraw from your examination until you have perfectly satisfied yourself as to the character of the labour. Having accomplished this, a second examination during this stage, unless it be prolonged, would be unnecessary, p. 96.

XXII. The first stage of labour is not always completed before the second begins. The grinding pains merge into the bearing pains so gradually as to require some attention to observe the change. When the bearing pain comes on, the patient is obliged to grasp firmly whatever is within her reach, she retains her breath more than before, and sometimes makes an involuntary effort to bear down. Her voice also alters, its tone is more subdued, and she seems more patient of her sufferings than before. Sometimes the complete dilatation of the os uteri is marked by the constitutional symptoms; there may be a slight rigor or vomiting, perhaps a strong inclination to go to stool, p. 100.

XXIII. The patient must now remain in bed, which should be properly prepared for her reception. A skin of morocco leather, or a broad piece of India-rubber cloth, must be placed next the bed, to protect it from being stained, and a blanket folded very wide, and enclosed in a soiled sheet, placed underneath the hip of the patient, as she lies upon her left side. They should be so fastened together, that the whole may be removed at once, pp. 100, 101.

XXIV. When the bearing pains become decided, a sheet should be fastened to the bed-post, and so within reach of the patient that she may grasp it firmly. The nurse should support her feet by pressing a pillow against them. Small pillows between the knees are very inconvenient, p. 102.

XXV. During the second stage, the patient often becomes fatigued and thirsty. You must imperatively forbid heating drinks of all kinds to be given to the patient. A free ventilation of the apartment should be secured, and as few persons as possible permitted to remain in the room, p. 102.

XXVI. In the second stage of labour more than one examination is necessary, but if examinations be repeated too often, the passages become dry and irritated. *The first object* of the examination is to determine the proportion between the head and the pelvis; *the second object* is to ascertain the exact position of the head; *the third object* is to note the progress made by the head, pp. 105, 106, 107.

XXVII. In order first *to determine the proportion*, the finger should be passed, in the interval of the pains, between the pubis and the head, and moved round on either side. The ear can be felt if there be sufficient space for the head to pass, but if the

head be high up in the pelvis, the finger can only just touch it, p. 105.

If the ear cannot be reached, and there seem to be a disproportion, its degree may be judged by feeling the scalp. When the head is only slightly compressed, the scalp is simply folded or puckered by the closing of the sutures; as the compression increases, these folds merge into one, which ultimately forms a distinct tumour. This continues to enlarge, so that in cases of *impaction* of the head it is sometimes of great magnitude, p. 105.

XXVIII. *To ascertain the exact position of the head*, pass the finger to the sagittal suture. This traverses the presenting part of the head, and looks towards the sacrum. Direct it along this suture towards the left obturator space and plane of the ischium, a fontanelle will then be felt: if it be the posterior (a small triangular space or a point where three sutures meet), it is the first position; if it be the anterior fontanelle (a large membranous space bounded by four sides or undefined), it is the third position. When the fontanelle is on the right side of the pelvis, the second position may be distinguished from the fourth in the same manner, pp. 78, 79.

XXIX. If there be a doubt of, or difficulty in, feeling the fontanelle, the ear will determine the side of the pelvis to which the occiput is applied, the lobe of the ear being felt on the corresponding side, p. 80.

XXX. You must be prepared to support the perinæum the moment it suffers any degree of distension. Sit behind the patient as she lies upon her left side, the back of the chair being towards the head of the bed, and while the head of the child is passing through the pelvic cavity press moderately *with the left hand* over the hip of the patient. *With the right hand* support the perinæum. A single fold of a fine napkin should be placed along the edge of the perinæum, and the right hand so applied that the fold of skin between the fore-finger and thumb should correspond to it, the fore-finger and thumb passing on either side of the vulva, and the palm of the hand resting against a thicker fold of the napkin, applied to the posterior part of the perinæum, pp. 107, 108.

XXXI. The perinæum should not be *pressed against too soon*; rather wait till you feel the head protruding with each pain

through the vulva. The pains are very unequal, and the object of being prepared to support the perinæum early is to resist too violent distension by them. Support the perinæum against the powerful pains. It is not necessary to press strongly when the pains are weak, pp. 108, 109.

XXXII. When the head is passing the vulva, do not draw the perinæum backwards nor push the head too rapidly forwards. In either case the widest part of the head would be forced too soon over the perinæum. When the head is delivered, examine carefully lest the funis be coiled round the neck. If it be a single coil it is sufficient to draw down more of the funis and loosen it. If two or more, one must be brought over the head. When this cannot be done the funis must be divided, which should never be done if it be possible to avoid it, p. 109.

XXXIII. When the shoulders are passing the perinæum, caution must be used lest the arms should lacerate it, especially in second positions of the head. Sometimes the shoulders require to be assisted, which may be done by placing the forefinger of the right hand within the axilla of the child's arm, on the pubic side, and guarding the perinæum carefully with the left hand, p. 110.

XXXIV. As soon as the shoulders and thorax of the child are delivered, it can respire, and is so far beyond danger; no haste should, therefore, be used in extracting the body and lower limbs; it is preferable to allow the uterus gradually to expel them, and while it is doing so, the left hand should be immediately applied over the fundus, in order to maintain a moderate pressure upon the uterus while it is descending towards the pelvis, p. 110.

XXXV. Pressure should then be maintained either by a temporary bandage or by the hand of an assistant, and the funis tied and divided. Apply a strong ligature of housewife thread, bobbin, or narrow tape, about two inches from the umbilicus, and a second about an inch further. You must be careful to see the part of the funis you are dividing, lest the fingers or any part of the child should be in your way, and also to examine the cut surface of the umbilical portion. The blood should be squeezed out of the vessels, and the surface wiped with a napkin, for the purpose of detecting any oozing of hæmorrhage that might take place if the funis were not properly tied, pp. 110, 111.

XXXVI. The hand may be applied to the fundus and a very moderate pressure is often sufficient to expel the placenta completely out of the vagina; but if not, it can be drawn out by the funis quite easily, directing the traction in the axis of the vagina. But if the uterus should not obey the stimulus at first, do not persevere; it is always more advisable to wait for some time than to use too much irritation. Neither should you attempt to remove the placenta by the funis alone, p. 111.

XXXVII. After delivery, a bandage is necessary to support the pelvis by compressing it as much as possible, and to support the uterus by moderate and equable pressure over the whole abdomen. The bandage should be rather more than a yard long and half a yard wide. It should first be drawn evenly over the pelvis; its lower edge, when so placed, being about an inch below the trochanter. This margin should be drawn as tightly as the patient will bear and pinned securely below the right trochanter. The bandage should be again drawn and pinned in a similar manner across the crista of the ilium, so that the pelvis may be embraced by this portion of the bandage, about three inches in width, as tightly as possible. The remainder of the bandage should be drawn and pinned with moderate tightness, but equally from the pelvis to the diaphragm, so that the whole abdomen be included in it without projecting over the bandage, p. 112.

XXXVIII. When the bandage is applied, the folded sheets that had been placed under your patient during labor should be removed, and replaced by others dry and warm, in order that she may be induced to sleep. She must not be disturbed afterwards, p. 113.

XXXIX. If the placenta be not separated in the first instance, wait for an hour, and again, by pressure, excite the action of the fundus. This should be brought as nearly as possible toward the centre of the pelvis, and grasped firmly by both hands. As soon as the fundus becomes hard, strong pressure upon it is generally sufficient to cause the expulsion of the placenta: if it should not, do not use any violence; rather let the nurse, under your direction, maintain the fundus in the same position, while you pass the fingers, and, if necessary, the hand, into the vagina, in order to stimulate the uterus to contraction, pp. 114, 115.

XL. In order to remove the placenta, the funis should be

held firmly in the left hand, and the fingers of the right passed along it within the vagina. Sometimes this alone excites contraction; but if not, all the fingers, in a conical form, may be introduced within it as far as the os uteri. You will often find a large portion lying at the upper part of the vagina, you may even feel the insertion of the funis, but do not attempt to withdraw it; pass the hand still towards the os uteri, and by irritating it, the portion of the placenta that lies within the cervix is often detached. If this be not sufficient, withdrawing the hand along the vagina for a short distance will excite contraction; but if both means fail, the fingers must be introduced within the os uteri, when the upper part of the placenta may be grasped and the whole removed. The assistant should press on the fundus while the hand is being withdrawn. The placenta should never be drawn away by the lower portion, p. 115.

SECTION III.

DIFFICULT LABOURS.

XLI. Difficult labours are divided into *tedious* and *laborious labours*. When labour is prolonged beyond the average period (twenty-four hours) without being at any time unusually severe, it is called *tedious*. When, without reference to time, there is a powerful struggle carried on by the uterus to overcome some unusual resistance, it is called *laborious*, p. 118.

XLII. When labour is tedious from *over-distension* of the uterus, puncture the membranes within the os uteri as high as you can reach, so that the liquor amnii may escape gradually. When the uterus is thus relieved, the pains increase in strength and frequency, and labour proceeds rapidly to its conclusion, p. 119.

XLIII. When *extreme obliquity of the uterus* is the cause of delay, mere change of position is sometimes sufficient to remove this difficulty. If not, or the uterus hang very much over the pubis, a bandage may be applied so as to draw the fundus gradually upwards, pp. 119, 120.

XLIV. When the fundus of the uterus is thrown very much

forward, the os uteri is drawn upwards and backwards towards the promontory of the sacrum ; and when labour begins, the anterior lip, being the most dependant part, often becomes so thin as to resemble the membranes. It might be mistaken for them ; and the cause of delay supposed to be rigidity of the membranes. The supposed membranes—that is, the uterus—might be accordingly punctured. If the presenting surface be carefully examined, the os uteri will be felt towards the promontory of the sacrum, p. 121.

XLV. When a patient is suffering from *hysterical excitement* great caution is necessary. Conversation should not be permitted within hearing of the patient ; none but her immediate friend and nurse should be allowed in the room ; strict attention should be paid to the evacuations, p. 122.

XLVI. *Mental despondency* is obscurely marked by symptoms of nervous shock—a rapid and feeble pulse—a tendency to chill—coldness of the extremities—constant watchfulness—listless expression—feeble pains ;—these are the most dangerous cases. Stimulants are necessary : ergot of rye, to hasten the action of the uterus—and the forceps, if the second stage be at all prolonged, pp. 126, 127.

XLVII. When the os uteri is *rigid*, and the woman of a plethoric habit, a free depletion from the arm, followed by nauseating doses of tartarized antimony, will often relieve it. If she be hysterical, or leuco-phlegmatic, local depletion and cathartic enemata are preferable, p. 128.

XLVIII. When the head is *arrested with the face towards the pubis*, the correction may sometimes be made with the fingers, but it is preferable to introduce the vectis on the pubic side of the pelvis, to press back the head in the intervals of the pains, and to rotate it gradually towards the sacral side. If the *face-presentation* be in a similar manner *arrested*, as few vaginal examinations as possible should be made, merely such as are sufficient to determine the arrest : when this is ascertained, recourse must be had to the forceps. When *the longitudinal axis of the head is fixed across the pelvic cavity*, the anterior fontanelle is very distinct. In the interval of the pains, press up the forehead, and so retain it until the following pain returns, when the occiput will descend, p. 134.

XLIX. *The hydrocephalic head* may be known by the increased

size of the fontanelles—the mobility and separation of the sagittal suture—the great overlapping of the parietal bones during a pain, if the head have entered the pelvic cavity, and the general looseness of the bones. If the head be arrested, perforation is required ; but the head thus enlarged often passes through the pelvis without assistance, pp. 137, 138.

L. *When the head is above the brim* of the contracted pelvis, and it is desirable to determine the degree of disproportion, the simplest and most efficient pelvi-meter is the hand of the practitioner. In ordinary vaginal examinations, when one finger only is introduced, the superior part of the sacrum is quite out of reach. When the point of the finger touches the upper part of the sacrum, it is evidence that the sacrum is pressed too much forwards ; if it touch the promontory, or just below it, it is very doubtful whether the head can pass. In some cases, when the cavity and outlet are very wide, the whole hand may be introduced within the pelvis ; and if the brim be contracted, the degree to which the fingers are compressed will determine the amount of diminution. In the standard pelvis the fingers, when in the brim may be slightly separated from each other. In a slight contraction, through which the head may pass, they are pressed close together. In greater disproportion the fingers overlap each other ; and in extreme cases, three, two, and sometimes even one finger only can be introduced ; but, excepting in such extreme cases, do not hastily assume, even with every evidence in this way obtained, that the head cannot pass the brim, until time and close observation of the labour removes every doubt, pp. 143, 144.

LI. *When the head is fixed in the brim of the OVATE pelvis*, the outlet and cavity being wide, the head may be delivered with the long forceps, p. 146.

LII. The term “*arrest*” is applied to cases where, although the head ceases to advance, the cause either does not depend upon disproportion in the pelvis, or when disproportion exists, it is not so great as to render the delivery of a living child impossible. When the head is pushed back, the finger can pass with facility between the head and the pelvis, the ear may be touched, the parietal bones do not overlap each other strongly, and the scalp is only puckered. If a tumour be formed on the presenting part, it is diffused, increases slowly, and seldom attains any magnitude, pp. 146, 147.

LIII. The term "*impaction*" is employed when the head not only ceases to advance, but when there is every evidence that its further progress is beyond the power of the uterus. The head cannot be easily displaced, it is impossible without force to pass the finger between it and the pubis; the ear cannot be felt; the urethra is compressed, the parietal bones are strongly overlapped, and a tumour grows very rapidly to a great size, often completely obscuring the characters of the presentation; the vagina is also swollen and congested, pp. 146, 147.

LIV. Difficult labours become dangerous to the patient, when they give rise either to inflammation of the passages or exhaustion of the uterus. Inflammation of the vagina may terminate in slough; inflammation of the uterus, in softening of its proper structure: the former may cause fistula, the latter rupture of the uterus. Exhaustion gives rise to sudden and dangerous hæmorrhage which may be fatal, and sometimes causes fatal syncope without hæmorrhage. In either case—whether inflammation be present, or the premonitory symptoms of exhaustion show themselves—labour must be brought to a conclusion, p. 149.

LV. In cases of slight deviation from the standard pelvis, where there is every evidence of space sufficient for the head ultimately to pass through the pelvis, if nature be allowed time for the purpose, interference is improper, although the period of twenty-four hours may be exceeded. But if the head be arrested, and so remain for four hours, notwithstanding the pains are regular and strong, the forceps may be used. If the pains be only feeble or absent, the action of the uterus may be suspended; although the head may be arrested for four hours during that suspension, it may advance when the action of the uterus is renewed. Assistance is not therefore required, until it is found that the uterus fails to do so, pp. 156, 157.

LVI. When the head is not arrested, but at the same time advances so extremely slowly, in consequence of the narrowness of the pelvis, that it seems to be arrested, there is no increased danger either to the mother or to the child* in leaving the deliv-

* Feeble constitutions are seldom able to endure very protracted suffering; cases, consequently, sometimes occur which are exceptions to the above rule, in which the premonitory symptoms of exhaustion appear before labour is concluded. The slightest indication of exhaustion should be attended to, and, when present, delivery effected.

ery to the natural efforts. The forceps is not therefore called for, p. 164.

LVII. *When the head is impacted* in the pelvic cavity, it cannot be delivered by the forceps without such injury to the passages as might endanger the mother's life. The probability of preserving the child's life is not sufficiently certain to justify an attempt which might be so hazardous. In the great majority of these cases, the death of the child takes place naturally; and it may be removed before symptoms dangerous to the mother present themselves, p. 176.

LVIII. If the reverse occur, and danger to the mother—whether from exhaustion or extending inflammation—be indicated before the death of the child, then perforation is called for, rather than render the risk to the mother a certainty by the dangers that result from a forcible extraction by the forceps, p. 176.

LIX. *When the head is retarded at the outlet* of the pelvis, it may arise from the perinæum being rigid, or from the arch of the pubis being too narrow, so that the head cannot pass out between the ischio-pubic rami. The strictest attention is necessary to subdue any tendency to inflammation in the perinæum. Fomentations must be sedulously employed, and, if necessary, depletion by leeches. Sometimes the vectis may be passed on the pubic side of the head, to assist its advance; but where the difficulty arises from narrowness of the pubic arch, the forceps is preferable, pp. 176, 177.

SECTION IV.

OBSTETRIC OPERATIONS.

LX. The rules that govern the application of instruments are founded upon three principles:

1st. *To preserve the lives of the mother and child.* If this be doubtful—

2nd. *To preserve the life of the mother without reference to the child.* When this cannot be done—

3rd. *To save the child if possible,* p. 187.

LXI. The instruments used for the *first object* are the *vectis* and the *forceps*. For the *second*, the *perforator*, the *crotchet*, the *craniotomy forceps*, the *osteotomist*, the *cephalotribe*. For the *third*, the Cæsarian section is performed, pp. 186, 187.

LXII. The *vectis* is intended to act as an extractor, to assist the feeble action of the uterus, to correct mal-positions of the head, or to overcome any unusual resistance of the perinæum. It is not, therefore, an instrument of much power, and its use is limited to the removal of slight impediments to the passage of the head, p. 187.

LXIII. The *forceps* is more generally used in the practice of midwifery, and is an instrument of much more extensive application. It may be employed when the head is at the outlet, in the cavity, or in the brim of the pelvis. The *short forceps* is used in the two former operations, the *long forceps* in the latter, pp. 191, 192.

LXIV. The following general rules must be observed before these instruments are applied:—

1st. It has long been established as a general rule, that instruments are never to be used in the practice of midwifery; the cases in which they are used, are, therefore, to be considered merely as exceptions to this rule [Denman].*

2nd. But such cases can very seldom occur in the practice of any one person,† and when they do happen, neither the *forceps* nor any other instrument is ever to be used *in a clandestine manner* [Denman].

3rd. The first stage of labour must be completed, that is, the os uteri must be dilated and the membranes broken, before we think of applying the *forceps* or the *vectis* [Denman].

4th. The difficulties which attend the application and use of the *forceps* are far less than those of deciding upon the proper time when, and the cases in which, they ought to be applied [Denman].

5th. The lower the head of the child has descended, and the longer the use of the *forceps* is deferred, the easier in general will their application be, the success of the operation more certain, and the hazard of doing mischief less [Denman].

* Aphorisms.

† An obstetric physician, who is frequently consulted, will of necessity meet more cases of this kind than another.

6th. Care is also to be taken that we do not, through aversion to the use of instruments, too long delay that assistance we have the power of affording with them [Denman].

7th. A rule for the time of applying the forceps has been from the following circumstances: that, after the cessation of the pains, the head of the child should have rested, for *six hours*,* in such a situation as to allow the use of the forceps before they are used [Denman].

8th. But this, and every other rule intended to prevent the rash and unnecessary use of the forceps, must be subject to the judgment of the person who may have the management of any individual case [Denman].

LXV. Before the vectis is applied, you must first observe those preliminary measures necessary in all obstetric operations. The urine should be withdrawn with an elastic gum catheter of rather large size (No. 10), and without the stilette. It is always safer to use a catheter of this kind, because there is less risk of injuring the urethra, if it should be compressed, than if the unyielding silver catheter be employed. An enema should also be administered, and when the bowels are relieved, the patient, lying on her left side, must be drawn as near to the edge of the bedstead as possible. The pelvis must be raised more than usual, and if the patient has been lying on a bed, and not on a mattress, it would be advisable to place a hair cushion under the hips. The vectis should be placed in warm water, and anointed, p. 189.

LXVI. Introduce two fingers of the right hand between the head and the symphysis pubis. Passing them on either side of the symphysis, the tip of the ear will be felt without difficulty. The finger must remain applied to it while the vectis is being introduced. It should be held about the middle, between the two forefingers and thumb of the left hand, and the handle directed downwards and backwards towards the coccyx, in order that the blade may lie flat upon the head, when the instrument is passing between it and the fingers of the right hand. When the blade is so applied, press it gently forwards with a slightly oscillating motion, until the edge reaches the lobe of the ear,

* *Four hours* are sufficient for this purpose, and the cessation of the pain is not essential.

which is now placed exactly between the finger and the vectis. The handle must therefore be depressed still more, in order that the edge may pass over the ear. When this is safely accomplished, the finger may be withdrawn, and the vectis passed forward to its proper position, pp. 189, 190.

LXVII. "Then, grasping the handle of the instrument firmly in the right hand, wait for the accession of a pain" [Denman], (which, although absent before, almost always returns when this new irritation is applied to the uterus), and draw steadily with it. "When the pain ceases let the instrument rest, and on its return repeat the same kind of action, resting and acting in imitation of the manner of the pains" [Denman], p. 190.

LXVIII. Carefully avoid using the vectis as a lever; and in order to do so the more certainly, it is better to pass two fingers of the left hand between the head and the perinæum, and to grasp the shank of the instrument with the remaining fingers; counter-pressure is thus made similar to the forceps, and the vectis may be used solely as a tractor, p. 190.

LXIX. When the head begins to advance, and to press strongly on the perinæum, the introduced fingers may be withdrawn, and the vectis maintained in its position, rather for the purpose of acting with it if the pains should again become feeble, than to extract the head by its means, if the uterus be sufficient to expel it: thus the perinæum will be better secured from injury, pp. 190, 191.

LXX. When the vectis is used to correct mal-positions of the head, it is better to operate with one not too much curved. One blade of the forceps will often answer in these cases, p. 191.

LXXI. *The operation with the forceps, when the head is resting on the perinæum*, may be undertaken, if there be any diminution in the transverse measurement of the outlet of the pelvis. The pubic blade with the lock looking upwards, must be passed over the ear in a similar manner to the vectis; and when so placed, the handle may be raised towards the pubis, and there maintained by an assistant in its *exact position*. Pass two fingers of the left hand between the head and the perinæum; hold the sacral blade lightly by the handle with the right hand; and, taking the lock of the introduced blade as your guide, endeavour to direct it so along the introduced fingers, that the edge of the sacral may pass along the lock of the pubic blade. As the sacral

blade passes forwards, and the locks approach each other, the handle of the pubic blade should be taken in the left hand, and drawn slowly towards the perinæum. In this manner the locks will glide together, and the instrument be applied without much difficulty, p. 193.

LXXII. When the pains return, the handles of the forceps should be held firmly, the perinæum being carefully supported by an assistant, and traction made at first very moderately, carefully observing the action of the uterus; and as you perceive that the pains are inefficient, the force may be increased. With each effort the handles may be drawn, first with a slightly waving motion to either side, and then upwards towards the pubis, in order that the head may pass in the axis of the vagina. When the head advances, and you are satisfied that the difficulty is overcome, it is better to leave the rest to the uterus, because there is less risk of injuring the perinæum. For the same reason, it would be advisable to unlock the forceps, and to withdraw the sacral blade, leaving the pubic to act with as a vectis, pp. 193. 194.

LXXIII. *When the head is arrested in the cavity of the pelvis*, the forceps may be introduced in the same manner as before; but if the ear be out of reach, the presentation must be carefully examined. The lambdoid suture may be traced passing up from the posterior fontanelle towards the ear. The trochanter also will give the direction of the plane of the ischium. Both will assist in guiding the pubic blade. The passage of the sacral blade is rather more difficult; its advance is frequently checked as it approaches the brim of the pelvis. If such should happen, be very careful not to use force in pressing it forwards. It is better to act with the pubic blade for a short time as a vectis, and if the head advance, even slightly, the opposite blade will frequently glide into its place, p. 197.

LXXIV. When the forceps is applied, dislodge the head from its situation in the first instance, because some accidental displacement may be the cause of difficulty, which the uterus cannot correct. When the head is relieved, it will glide into the correct position, and may be delivered without difficulty. If, however, the head remain arrested, the forceps must be seized firmly; and in order to secure your hold, a coarse napkin might be placed loosely round the handles. A steady and powerful traction should

be maintained so long as the pain continues ; and when it ceases, *the grasp of the instrument must be at once released, and remain so until the succeeding pain*, when the same steady traction may be renewed. As the head advances more easily, and approaches the perinæum, leave it as much as possible to the uterus, pp. 197, 198.

LXXV. When the forceps is in the pelvic cavity, the shank of the instrument lies between the tubers of the ischia. When the head is in the hollow of the sacrum, the handles are directed towards the pubis. It is advisable rather to follow than to guide the direction of the head, because as it descends it will *naturally* change its position, which might be prevented by attempting too hastily to turn it, pp. 198, 199.

LXXVI. *The operation when the head is fixed in the brim of the pelvis* differs from either of the preceding operations ; the blades are applied over the occiput and face of the child, and not over the ears. Two fingers, and as much as possible of the right hand, should be passed behind the trochanter towards the centre of the ilium on the superior side of the pelvis ; and if the anterior fontanelle be felt distinctly, the longer blade of the forceps (if they be unequal) should be passed over the fontanelle to the face of the child : the shorter blade may then be passed in the opposite direction over the occiput, guiding it by the lock of the introduced blade, p. 199.

LXXVII. When the forceps is properly applied, the handles look downwards and backwards towards the perinæum, and in the axis of the brim ; traction must be made in this direction, and when the pain commences, the handles of the instrument should be held as in the former case, and the force gradually increased according to the resistance. Two or three trials will generally succeed in extricating the head from the brim of the pelvis, when it will rapidly advance without assistance : the forceps might be removed, but it is safer not to do so, lest any impediment should delay its further progress, p. 199.

LXXVIII. Before *perforation* is performed, the bowels should be relieved, and the urine removed from the bladder, which should be carefully examined above the pubis, because sometimes the neck and fundus of the bladder are so compressed between the head and the pubis, that if a short catheter be used, the urine is only removed from the lower segment, while the principal portion remains above. When this is the case, the head should be pressed

back as much as possible, and a long gum-elastic catheter, of the size that will pass the urethra, introduced; great care should be taken not to use force in its passage, lest the instrument might perforate the canal, pp. 206, 207.

LXXIX. Two fingers of the right hand should be passed to the most depending part of the tumour that presents. The whole surface of the presentation should be carefully examined, in order to determine the degree of dilatation in the os uteri. If not fully dilated, observe the exact relation of the os uteri to the tumour on the head. If the os uteri be very thin, and closely embrace the tumour, its margin must be defined, lest the degree of dilatation escape your notice, p. 207.

LXXX. Let the forefinger of the right hand remain applied to the centre of the presenting part, but rather to the pubic side, and with the left hand introduce the perforator, having the point (which is slightly curved) resting upon, and guarded by, the forefinger. When both fingers meet, the right hand may be withdrawn from the vagina, and the perforator forced through the tumour and bone at the part where the forefinger had been applied. The first two fingers of the left hand should then rest on the stops of the instrument, to prevent the perforator slipping from the opening when the handles are raised, and to secure the vagina from injury. When the bone is sufficiently broken in one direction the handle may be changed to the opposite, and a crucial opening made. The perforator should be passed into the opening, and the brain completely broken up, p. 207.

LXXXI. When the perforator is withdrawn, the forefinger of the right hand should still remain within the opening, and the crotchet introduced, having the handle directed backwards towards the coccyx, when, with a little careful management, it may be passed through, and the remainder of the brain removed. The handle of the crotchet should then be held firmly in the right hand, while two or three fingers of the left are applied to the bone externally. The force should be so applied that the bone may be held tightly between the flat part of the point and the finger, without directing it much on the point itself, p. 208.

LXXXII. When the craniotomy forceps is used, the blade, which is rough or dentated, is introduced within the cranium, while the opposite blade is passed outside. When they are closed, the teeth perforate the bone, and are received by corresponding

openings in the opposite blade. Extraction may then be made without difficulty, provided the bone does not give way. This is less likely to take place when the blade is not made with teeth, p, 209.

LXXXIII. The induction of premature labour may be effected by introducing a sponge tent within the os uteri, so as to dilate it. If this fail, puncture the membranes with a stilette. Uterine action generally commences within forty-eight hours, p. 223.

LXXXIV. The induction of premature labour is not applicable to first pregnancy, unless there be very clear proof of great distortion.

The most certain evidence is the result of previous labours, where perforation had been required, or when, with every successive labour, the contraction of the pelvis may have so increased as to render the last labour more difficult than the preceding one: so that, ultimately, perforation might be necessary, unless in this way obviated, p. 223.

INDEX.

	Page
ABDOMEN, pendulous	41
diminution in size of	58, 60
use of bandaging	88
how to bandage	112, 113
Adhesion of placenta	89
Ala of pelvis	14
Amnii, liquor, use of	60, 61, 63, 64
effect of premature	
escape of	98
effect of sudden	
discharge of	119
gradual escape of	121
Amnion, dropsy of	118
Anatomy of pelvic bones	14
Angle, pubic,	21 n. 28
Aphorisms	255
Arm presentation	98, 99, 106
Arrest, causes of	106, 145, 146
and impaction of head	146
use of forceps in	195
see Head	
Articulation, anatomy of sacro-	
iliac	18
of sacrum	
& coccyx	18
sacro-ver-	
tebral	18
Auscultation, use, in pregnancy	
of	171, 172
in perforation	204
Axes of pelvis, see Pelvis	

B.

Bandage, use of	88
application of	112
Baudeloque, measurements by	32
on rickets	49
callipers of	51
on positions of head	77
experiments with for-	
ceps	169
jun. cephalotribe	216, 251
Beatty, Dr., on ergot of rye	153 n.
forceps by	237

	Page
Bed, how to be prepared	101
Bell, Sir C., on uterine fibres	58, 61, 62
on the abuse of the	
forceps	176 n.
Biparietal axis, true position of	30
Births, danger in male and female	135
Bladder, state of, previous to la-	
bour	91
prolapse of	180
Blundell, Dr., division of labours	
by	56
on use of forceps	174
Boivin, measurements by	32
pelvimeter by	52
Bone, labour delayed by deposit	
of	16, 17, 40
law of growth of	39
inference from prominence of	70
Bregma, situation of the	29
Brünigghäusen's forceps	200
Burns, Dr., measurements by	28, 32
division of labours by	56
on rupturing the	
membranes	98
on using forceps	156

C.

Cæsarian section in mollities os-	
sium	45
in osteo-sarco-	
ma	184
attempts to su-	
persede	216, 217
statistics of	218-220
contrasted with	
perforation	219
reason of failure	
of	220
mode of oper-	
ating	220
dangers of	221
Campbell, Dr., on using forceps	156
Catheter, sort to be used	189
Cephalotribe described	216
objections to	216

- | | Page | | Page |
|---|---------------|---|------------------|
| Chamberlen, Dr. P., invents forceps | 225, 228 | Deweese, Dr., division of labours | 56 |
| Child, management of new born | 110 | on uterine contraction | 67, 68 |
| signs of death of | 147, 172 | Diarrhœa previous to labour | 91 |
| Chin, position, in pelvis, of child's | 29 | Difficult labour, see Labour. | |
| Churchill, Dr., division of labours | 56 | Digital measurements | 52, 143 |
| by | 56 | Dilatation, see Uteri Os. | |
| on Cæsarian section | 218 | Distortion of pelvis | 33, 43 |
| crotchet by | 240 | causes of | 44—47 |
| Clark, Dr. J., on conjugate axis | 30 | Doherty, Dr., on rigidity of vagina | 178, 179, 180 n. |
| Cloquet, measurements by | 32 | Dress and position of the patient | 101 |
| Coccyx, advantage of mobility of | 17 | Drew, Dr., pelvic tumours by | 184 |
| Collins, Dr., on face-presentation | 84 | Drinks proper for patient | 102 |
| on hydrocephalic head | 136 | Dublin lying-in-Hospital, cases at | |
| on using forceps, | 157, 159, 175 | position of head | 81 |
| on stethoscope | 172 | face-presentation | 84 |
| short forceps by | 233 | mental despondency | 124, 125 |
| Conjugate axis of pelvis | 30 | head too large | 136 |
| Conquest, Dr., short forceps by | 194 | Duties of accoucheur, first | 92, 93 |
| Constipation previous to labour | 91 | | |
| Contouly's pelvimeter | 51 | E. | |
| Contraction of uterus | 53—61 | Ear, diagnostic position of child's | 78, 80, 195 |
| Corset, ill effects of the | 88 | as a test of the labour | 105 |
| Craniotomy forceps described | 209, 240 | Ergot of rye, use of, | 153, 154 |
| Crotchet, see Perforation. | | Examination, consequence of frequent | 53, 105 |
| mode of employing the | 207, 208 | first vaginal | 94 |
| and craniotomy forceps | 209 | when to be made | 95 |
| double, by Dr. Davis | 215, 216 | objects gained by, | |
| by Dr. Churchill | 240 | in first stage | 95, 96 |
| D. | | early, when necessary | 99 |
| Davis, Dr., on pelvimeters | 51, 52 | objects gained by, | |
| division of labours by | 56 | in second stage | 105—107 |
| on forceps in impaction | 174 | Exhaustion of uterus | 149 |
| craniotomy forceps by | 209, 240 | symptoms of | 152 |
| osteotomist by | 215 | treatment of | 152—154 |
| forceps by | 234, 235 | Expulsion of child, sudden | 34 |
| Death of child, signs of | 147, 171, 172 | F. | |
| Deformities in pelvis | 33, 43 | Face, see Presentation. | |
| causes of | 44—47 | Fatigue distinguished from exhaustion | 149 |
| Delay, arguments for and against, | 162, 163 | Feelings in pregnancy, change of the | 126 |
| danger of, when forceps required | 195—197 | Fistula from use of forceps | 163, 164 |
| Denman, Dr., division of labours | 56 | Flatus previous to labour | 90 |
| by | 56 | Fontanelle, anterior | 29 |
| on stages of labour | 57 | distinction of anterior and posterior | 79, 80 |
| on using forceps | 157 | Forceps, long, in arrest of head | 146 |
| short forceps by | 194, 233 | opinions when to be used | 156, 157 |
| perforation by | 239 | list of operations with | 158, 165 |
| Dentition, cause of late | 47 | effects from employing | 163, 171, 175 |
| Despondency, effects of, on labour | 123 | | |
| treatment of | 126, 127 | | |
| Development of pelvis arrested | 35 | | |

	Page
Forceps in impaction of the head	167, 168, 170
Baudeloque's experiments with	168, 169
an extractor	170 n.
long and short, peculiar uses of	191, 192
when head on perinæum	192, 193
directions for locking	193
how to be used in extraction	194
rules in using	194, 195
danger of delay when requisite	195—197
long, different kinds of	200
practice with, on phantoms	202
in different positions of the head	202, 203
craniotomy	209, 240
history of the	225, 228—233
Smellie on the	231, 232
dimensions of various	233, 234
by Dr. Davis	234, 235
by Dr. Beatty	237
measurements and drawings of various	243—249
Forehead, see Presentation.	
Funis, prolapse of,	99
coiled around the neck	109
time and mode of tying the	110, 111

G.

Gaitskill's vectis	187, 188, 227
Gifford's forceps	229, 231
Gravity, influence of, on form of pelvis	44
Grinding pains, see Pains.	
Growth of pelvis	35, 36
bone, law of	39

H.

Hæmorrhage, internal, how caused	34, 89
Hand, inference from shape of	41
presentation	98, 106
Hardy, Dr., on ergot of rye	153, 154 n.
Head, action of planes of ischium in passage of the	15, 16
passage of, arrested by bony deposits	16
see Rotation.	
motion of, at outlet	22
a lever of third order	22
comparative largeness of human	22
tumour of, see Tumour.	
different positions of	28
see Measurements.	
passage, how effected	30

Head, passage of, through pelvis	75, 76
positions of	76, 77
first position of	77
second position of	78
third and fourth positions of	79
at perinæum	85
expulsion of, in various positions	86
misplacement of	134
male and female, contrasted	135
too much ossified	135
hydrocephalic	136
diagnosis and treatment of	137
how and where arrested	142
arrested above the brim	142
within the brim	145
in pelvic cavity	146
arrest and impaction of	147
general treatment of impaction of	176
retarded at outlet	176
resting on perinæum, operation when	192
in pelvic cavity	194—199
fixed in brim	145, 199
above brim	210
Holmes, Mr., perforator by	240, 251
Hunter, Dr. W., on uterus	62
on using forceps	157
Hydrocephalic head, see Head	
Hysteria in labour	121
treatment of	122
Hysterotomy, see Cæsarian section.	

I.

Impaction of head	147, 166
use of forceps	166, 175
in	166, 175
conclusions on	
treatment of	176
see Head.	
Inertia of uterus	87, 88
Inflammation of the passages	149
symptoms of	150, 151
treatment of	151
of perinæum	176
Interference, danger from pre-mature	99, 155
Instruments, effects of using	143
time for using	156, 157
employed, list of	186
rules in using	187
Ischiatic foramina, contents of	
sacro-	20
Ischium, boundaries of two planes of	15
advantages of their inclination	16
spine and tuber of	16

K.

Keyser on Cæsarian section	218, 219
Knox, Dr., on deformed pelvis	50

L.	Page	Measurements of pelvis . . .	Page
Laborious labour, see Labour.		iliac spinous processes . . .	14
Labour, see Parturition.		of first plane of pelvis . . .	24
definition of	55	of plane of brim	25
divisions of	55, 56	cavity	26
division adopted	56	of outlet	26
definition of natural	56	depth of cavity	28
stages of	57	transverse, of outlet	29
1st stage, mechanism of 64—69		of child's head—	
commencement of	71, 91	of longitudinal axis	28
premonitory symptoms		transverse	28
of	80	of eighteen healthy	
duration and manage-		pelves	31
ment of	94—102	by various authors	32
causes of delay in	118	of diseased pelvis	54
2nd stage, mechanism of		digital	143, 144
73—77		Measuring pelvis, modes of, 50, 51, 52	
management of	104—111	Mental despondency, see De-	
causes of delay in	123	spondency.	
3rd stage, separation of		Merriman, Dr., division of labours	
placenta	86	by	56
retention of	87—89	stages of labour by	57
management of	111, 115	on ovarian tumours 180, 181	
difficult, definition of	117	Miscarriage, influence of pelvis on	19
subdivisions of	118	Mollities ossium, effects of,	43, 140
tedious	118	symptoms of	48
causes of	118	Monro, Dr., measurements by	32
laborious	133	Moreau, measurements by	32
causes of	133	Mortality in forceps operations 158, 159	
management of	155	in unassisted labour 160—162	
use of forceps in	157—177	Muscles of ilium	14
perforation in	173	action of pelvic	46
premature, when neces-		action of abdominal	87, 88
sary	222	Muscular fibres of uterus	58—62
modes of in-			
ducing	223		
mechanism of natural,			
Aphorisms	255—257		
management of	257—263		
difficult, Aphorisms, 263—267			
Lee, Dr., measurements by	32		
on hydrocephalic head	137		
forceps cases by	158		
long forceps case by	201, 202		
Le Roy, proposed operation by	217		
Lever of third order, child's head			
acting as a	22		
Ligaments, broad	15		
short sacro-ischiatic	16		
sacro-coccygeal	18		
subpubic	20		
internal and great sacro-			
ischiatic	20		
Liquor amnii, see Amnii.			
Lowder's vectis	227		

	Page
Ould, Sir F., discrimination of	76
perforator by	239
Ovarian tumour	179
treatment of	180, 181
P.	
Pains, after	34
cause of severe	89
grinding and bearing	73, 92, 100
fruitless	150
Parietal protuberance, see Biparietal axis	
Parturition, common cause of difficult	113
difference of, in town and country	16, 49, 141
how difficult or easy in same person	25
difficult from shape of pelvis	40
see Labour.	
Patient, dress and position proper for the	101, 120
management of, after delivery	112, 113
fruitless attempts of the	150
effects of forceps on	170, 171
Pelvimeters described	50—52
estimation of	143
Pelvis, effects of shape of	13
anatomy of bones of	14
difference in alæ of male and female	14
cavity of	14
two planes of	14, 15
divisions of—true, false	19, 20
effects of too wide or narrow brim	19, 33—35
boundaries of cavity of	20
shape and boundaries of brim and of outlet	21
mechanism of the	22
use of measuring	22
see Measurements.	
axes of brim and outlet	23, 24
intersection of axes	24
planes of the	24
proportions of the	26, 27
irregularities and distortion in the	33
development of, arrested	35
growth of	36, 37
contrast of male and female	38, 39
varieties, in axis, of brim	42
cordiform	42, 140
ovate	43, 139
forces acting on the	44
obliquely ovate	50
effects of masculine	138, 139
softened	139
characteristics of	141

	Page
Perforator, list of operations with the	158
in impaction of head	167, 168, 173
when preferable to forceps	176
symptoms requiring	204—206
mode of performing	206—208
of Dr. Denman	239
of Nægele	239
of Mr. Holmes	240
Perinæum	
subject to laceration	22, 86
distension of	84—86
how to support	107—109
when support is necessary	108
inflammation and treatment of	176
operation when head resting on	192, 193
Placenta, expulsion of	86, 87
causes of retention of	87—89, 114
adhesions of	89
how to be separated	111
extraction of	115
Placental murmur	119
Polypus, obstruction from	182
treatment of	182
Positions of head	76
expulsion of various	85
first	77
second	78
third and fourth	79
modes of distinguishing the	79, 80
management of	106
fronto-cotyloid	106
Practice, cause of difference in	141, 142
Practitioner, duties of obstetric	92, 93
Premature labour, see Labour.	
Presentation of vertex	77, 78, 79
forehead	81, 82, 134
face	82, 83, 134
necessity of distinguishing	83
consequence of	84
shoulders	94
hand and arm	98, 99, 106
funis	99
Presenting part	23
Proportion between head and pelvis, how ascertained	105
Promontory, see Sacrum.	
Puberty, change in pelvis at	37
Pubic angle	21 n. 23
Pubis symphysis, anatomy of	17
deposition of bone at	40

- | | |
|--|---|
| <p>Q.</p> <p>Quadrupeds, pelvis of . . . 36</p> <p>R.</p> <p>Radford, Dr., forceps by . . . 200, 233</p> <p>Ramsbotham, Dr., measurements by . . . 32</p> <p> on position of head . . . 77, 78</p> <p> on using forceps . . . 157</p> <p> case of polypus by . . . 182</p> <p> on impaction of head . . . 196</p> <p>Reflex nervous function . . . 64</p> <p>Rickets, effects of . . . 43, 139</p> <p> cause of . . . 47</p> <p>Riecke, on use of forceps . . . 174, 175</p> <p>Rigby, Dr., measurements by . . . 32</p> <p> cases quoted by . . . 35</p> <p> division of labours by . . . 56</p> <p> Wigand quoted by . . . 65</p> <p>Roonhuysen invents vectis. . . 225</p> <p>Rotation of Child's head in its passage . . . 15, 16, 22, 30</p> <p> how effected . . . 20, 21, 26, 27</p> <p>S.</p> <p>Sacrum, anatomy of the . . . 16</p> <p> promontory of, its importance . . . 17, 18</p> <p> above the brim . . . 24, 25</p> <p> influence of its position . . . 25</p> <p>Shaw, Mr., on immature pelvis . . . 37</p> <p>Sherwood, E. case of . . . 211—214</p> <p>Shoulders, direction of the . . . 85</p> <p>Shoulder-presentation . . . 94</p> <p>Show, nature of the . . . 91</p> <p>Sigault, proposed operation by . . . 217</p> <p>Simpson, Professor, on male births . . . 135</p> <p>Smellie on forceps . . . 231, 232</p> <p> scissors by . . . 239</p> <p>Softened pelvis, see Pelvis.</p> <p>Solly, Mr., case by . . . 48</p> <p>Spine, disease of . . . 16</p> <p> curvature of . . . 49</p> <p>Stages of labour . . . 57</p> <p>Stage first, mechanism of . . . 64—69</p> <p> second . . . 73—77</p> <p> third . . . 86</p> <p> see Labour.</p> <p>Stature no criterion of pelvis . . . 35</p> <p>Stethoscope, see Auscultation.</p> <p>Suppiot, Madame, case of . . . 43</p> | <p>Symphysis pubis, difference in male and female . . . 17</p> <p> see Pubis.</p> <p>Syncope, how caused . . . 34</p> <p>T.</p> <p>Table of pelvic proportions . . . 31</p> <p> various pelvic measurements . . . 32</p> <p> diseased pelvis . . . 54</p> <p> position of the head . . . 77</p> <p> comparative view of operations . . . 158, 165</p> <p>Tedious labour, see Labour.</p> <p>Thompson, E., case of . . . 48</p> <p>Tumefaction of child's face . . . 83</p> <p>Tumour of head as a symptom . . . 105, 205, 207</p> <p>Tumours, obstruction by . . . 177</p> <p> fibrous . . . 183</p> <p> case of . . . 183</p> <p> pelvic . . . 184</p> <p> see Ovarian Polypus.</p> <p>U.</p> <p>Urethra, position of orifice of the . . . 96</p> <p>Uteri, prolapsus, how caused . . . 19, 34</p> <p> stricture of cervix . . . 34</p> <p> inversio . . . 34, 35</p> <p> os, how dilated . . . 61, 62</p> <p> muscularity of . . . 62</p> <p> views on dilatation of . . . 68</p> <p> various conditions of . . . 70</p> <p> dilatable . . . 69</p> <p> rigid . . . 70</p> <p> causes of . . . 71</p> <p> inflammation of . . . 71</p> <p> symptoms of dilatation of . . . 100</p> <p> displacement of . . . 120, 121</p> <p> artificial . . . 63, 121</p> <p> treatment of rigidity of . . . 128</p> <p> inflammation of . . . 126, 128</p> <p> hypertrophy of . . . 130</p> <p> cervix, toughness of . . . 130</p> <p> treatment of . . . 130</p> <p> os, cartilaginous state of . . . 131</p> <p> treatment of . . . 131</p> <p> agglutination of . . . 177</p> <p> absence of . . . 177</p> <p> in forceps operations . . . 203</p> <p>Uterus, muscular fibres of . . . 58—61, 67</p> <p> contraction of . . . 58—61</p> <p> order of contraction of . . . 64</p> <p> action of, when os rigid . . . 71</p> <p> inertia of . . . 87, 88, 114</p> <p> effects of irregular contraction of . . . 88</p> <p> contraction of, after placenta expelled . . . 111</p> |
|--|---|

	Page
Uterus, suspended action of	114
irregular contraction of	115
over-distension of	118
symptoms	
of	119
obliquity of	119
cause of	120
inflammation and ex-	
haustion of	149
see Inflammation.	
symptoms of exhaustion	
of	152
treatment of	152—154

V.

Vagina, how inverted	34, 35
rigidity and stricture	
of	177
treatment of	178, <i>n.</i> 179, 180
Vaginal examination, see Examination.	

	Page
Vectis described	187, 226, 227
Gaitskill's rules for applying	187, 188
cases when advantageous	189
how to apply	189, 190
possible ill effects of	190, 191
history of the	225, 226
Roonhuysen's	226
Lowder's and Gaitskill's	227
Velpeau, measurements by	32
on stages of labour	57
Ventilation to be secured	102
Vertex, see Presentation.	

W.

Wigand's views on uterine action	65
Wood, Mr., operation by	48

Z.

Ziegler, Dr., forceps by	235
--------------------------	-----



CATALOGUE

OF

B O O K S

ON

MEDICINE, ANATOMY, SURGERY,

MIDWIFERY, CHEMISTRY,

AGRICULTURE, &c., &c., &c.

FOR SALE BY

SAMUEL S. & WILLIAM WOOD,

No. 261 PEARL STREET,

(OPPOSITE UNITED STATES HOTEL,)

NEW YORK.

1846.

MEDICAL CATALOGUE.

- ABERCROMBIE, J. Pathological and Practical Researches on Diseases of the Brain and the Spinal Cord. 8vo.
- Pathological and Practical Researches on Diseases of the Stomach, Intestinal Canal, Liver, and other Viscera of the Abdomen. 8vo.
- Inquiries concerning the Intellectual Powers, and the Investigation of Truth. 12mo.
- Philosophy of the Moral Feelings. 12mo.
- ABERNETHY, J. Physiological Lectures. 8vo.
- Surgical Observations, containing a classification of Tumours, with cases to illustrate the history of each species;—an account of Diseases which strikingly resemble the Venereal Disease; and various cases illustrative of different surgical subjects. 8vo.
- Surgical Works, containing constitutional origin and treatment of Local Diseases, on Aneurisms, Diseases of the Urethra, Injuries of the Head, Puncturing the Urinary Bladder, Tic Douloureux, Removal of loose substances from the Knee-Joint, Hæmorrhoidal Diseases, Fistulæ in the Perinæum, Tumours, Chronic and Lumbar Abscesses, &c. 2 vols. 8vo.
- ACCUM, F. Chemical Amusement, comprising a series of curious and instructive experiments in Chemistry. 12mo.
- Culinary Chemistry, exhibiting the Scientific Principles of Cookery. Plates. 12mo.
- ACKERLY, G. Management of Children in Sickness and in Health. 12mo.
- ADAMS, J. Observations on Morbid Poisons, Chronic and Acute, comprehending Syphilis, Yaws, Sivvens, Elephantiasis, and the Anomala confounded with them; and Acute Contagions, particularly the Variolous and Vacciné. 4 coloured plates. 4to.
- ALCOCK, T. Lectures on Practical and Medical Surgery, comprising observations and reflections on Surgical Education; on the Investigation of Disease, ordinary Duties of the Surgeon, &c. 12 plates. 12mo.
- ALCOTT, W. A. Mother's Medical Guide in Children's Diseases. 18mo.
- Young Mother; or Management of Children as to Health. 12mo.
- ALIBERT, J. L. Treatise on Malignant Intermittents, with Notes, &c., by Charles Caldwell, M.D. 8vo.
- ALISON, W. P. Outlines of Pathology, and Practice of Medicine. 8vo.
- American Gentleman's Medical Pocket Book, and Health Adviser; containing a statement of the modes of curing every disease to which he is liable; directions in case of Accidents on the Road or at Sea, &c. 32mo.
- American Lady's Medical Pocket Book, and Nursery Adviser; containing rules for preserving the health of unmarried Females; directions to pregnant and lying-in Women; an account of their diseases, &c. 12mo.
- AMUSSAT's Lectures on Retention of Urine, caused by Strictures of the Urethra; and on the Diseases of the Prostate. Translated by James P. Jervcy, M.D. 3 plates. 8vo.

- Anatomia Britannica.** A System of Anatomy and Physiology, selected from the works of Haller, Albinus, Monro, Winslow, Hunter, Soemmerring, Scarpa, Cruikshank, Masgagni, Murray, Walter, Sabatier, Meckel, Zinn, &c. 3 v. 12mo.
- Anatomical Examinations.** A complete Series of Anatomical Questions with Answers: the Answers arranged so as to form an elementary system of Anatomy; to which are added Tables of the Bones, Muscles, and Arteries. 12mo.
- Anatomical Remembrancer; or Complete Pocket Anatomist,** containing a concise description of the Bones, Ligaments, Muscles, and Viscera; the distribution of the Nerves, Blood-Vessels and Absorbents; the arrangement of the several Fasciæ; the Organs of Generation in the Male and Female, and the Organs of the Senses. 32mo.
- ANDERSON, W.** System of Surgical Anatomy. On the structure of the Groin, Pelvis and Perineum, as connected with Inguinal and Femoral Hernia, Tying the Iliac Arteries, and the operation of Lithotomy. 9 plates. 4to.
- ANDRAL, G.** Treatise on Pathological Anatomy; translated by Drs. Townsend and West. 2 vols. 8vo.
- Essay on the Blood in Disease; translated by Drs. Meigs and Stillé. 8vo.
- Medical Clinic, or Reports of Medical Cases—Diseases of the Abdomen, the Chest, and the Encephalon. With Observations, &c., by Dr. Spillan. 3 v. 8vo.
- ANDREWS, M. W.** Practical Observations on the Application of Lunar Caustic to Strictures in the Urethra; with remarks on its use in Strictures in the Oesophagus. 8vo.
- ANNESLEY, J.** Sketches of the most prevalent diseases of India; comprising a Treatise on the Epidemic Cholera of the East; Statistical and Topographical Reports of the diseases in the different divisions of the army under the Madras Presidency; the Annual rate of Mortality, &c.; Practical Observations on the effects of Calomel on the Alimentary Canal, and on the Diseases most prevalent in India; also, an Inquiry into Chronic Inflammations and Abscess of the Liver. Coloured plate. 8vo.
- ARAN, F. A.** Practical Manual of the Diseases of the Heart and Great Vessels: translated by W. A. Harris, M.D. 12mo.
- ARETÆUS** on the Causes and Signs of Acute and Chronic Disease: translated by T. F. Reynolds, M.D. 8vo.
- ARMSTRONG, JOHN.** Memoir of his Life and Medical Opinions: to which is added an Inquiry into the facts connected with those forms of Fever attributed to Malaria or Marsh Effluvium: by Francis Boott, M.D. 2 vols. 8vo.
- Art of Preserving Health: a Poem.
- Practical Illustrations of Typhus Fever, of the Common Continued Fever, and of Inflammatory Diseases. 8vo.
- Facts and Observations relative to the Fever commonly called Puerperal: Practical Illustrations of the Scarlet Fever, Pulmonary Consumption, and Measles: with observations on the efficacy of Sulphureous Waters in Chronic Complaints. 8vo.
- Lectures on the Morbid Anatomy, Nature, and Treatment of Acute and Chronic Diseases: edited by Joseph Rix. 8vo.
- Same work, with an Account of the Life and Writings of Dr. Armstrong, by John Bell, M.D. 2 vols. 8vo.
- ARNOTT, N.** Elements of Physics; or Natural Philosophy, general and medical: in plain or non-technical language. With additions by Isaac Hays, M.D. 8vo.
- ASHWELL, S.** A Practical Treatise on the Diseases peculiar to Women: with Notes by Paul B. Goddard, M.D. 8vo.
- On Incision in cases of Occlusion, and Rigidity of the Uterus.
- ASSALINI, P.** Observations on the Disease called the Plague, on the Dysentery, the Ophthalmia of Egypt, and on the means of prevention: with some remarks on the Yellow Fever of Cadiz, and the description and plan of a Hospital for the reception of patients affected with epidemic and contagious diseases. Translated by Adam Neale. To which is added a Letter concerning the Seasoning or Yellow Fever of the West Indies: by George Pinchard, M.D. 12mo.

- ATKINS, D. Medical and Surgical Cases and Observations. Plates. 8vo.
- AVERILL, C. A Short Treatise on Operative Surgery, describing the principal Operations as they are practised in England and France. For the use of Students in operating on the Dead Body. With additions by John Bell, M.D. 12mo.
- BAILLIE, M. Morbid Anatomy of some of the most important parts of the Human Body. 8vo.
- BAKEWELL, R. An Introduction to Geology: intended to convey a practical knowledge of the science, and comprising the most important recent discoveries; with explanations of the Facts and Phenomena which serve to confirm or invalidate various Geological Theories. With an Appendix by Prof. B. Silliman. Plates. 8vo.
- BALL, I. Analytical View of the Animal Economy, calculated for the Students of Medicine as well as private gentlemen. With allegories and moral reflections drawn from the subject, to awaken the mind to an elevated sense of the great Author of Nature. 12mo.
- BAMFIELD, R. W. Essay on Curvatures and Diseases of the Spine, including all the forms of Spinal Distortion: to which the Fothergillian gold medal was awarded by the Medical Society of London. Edited by J. R. Mitchell, M.D. 8vo.
- BANCROFT, E. N. An Essay on the Disease called Yellow Fever, with observations concerning Febrile Contagion, Typhus Fever, Dysentery, and the Plague. With Notes by John B. Davidge, M.D. 8vo.
- BARD, S. Compendium of the Theory and Practice of Midwifery, containing practical instructions for the management of Women during Pregnancy, in Labour, and in Child-bed. Many engravings. 8vo.
- BARNES, J. The New London Chemical Pocket Book, or Memoranda Chemica; adapted to the daily use of the Student. Engravings. 12mo.
- BARRY, E. A Treatise on the Three different Digestions and Discharges of the Human Body, and the Diseases of their principal organs. 8vo.
- BARTLETT, E. Essay on the Philosophy of Medical Science. 8vo.
 — History, Diagnosis, and Treatment of Typhoid and of Typhus Fever: with an essay on the diagnosis of Bilious Remittent and of Yellow Fever. 8vo.
 — Sketches of the Character and Writings of eminent living Surgeons and Physicians of Paris. With portraits of Broussais and Dupuytren. 12mo.
- BARTLETT, J. S. Physician's Pocket Synopsis: conveying an accurate and concise view of the History, Character, Symptoms, and Treatment of the various Medical and Surgical Diseases incident to the Human frame. Compiled from the best authorities, with references to the most approved modern writers: together with the natures, properties, and doses of the simple and compound agents employed in Pharmacy. Revised by Henry Coley. 18mo.
- BARTON, W. P. C. Outlines of Lectures on Materia Medica and Botany, delivered in Jefferson Medical College, Philadelphia. 2 vols. 12mo.
 — Vegetable Materia Medica of the United States; or Medical Botany: containing a Botanical, General, and Medical History of Medicinal Plants indigenous to the United States. 50 coloured plates. 2 vols. 4to.
- BARWELL, MRS. Infant Treatment; with Directions to Mothers for self-management before, during, and after pregnancy. Addressed to Mothers and Nurses. Revised by a Physician of New York under the approval and recommendation of Valentine Mott, M.D. 12mo.
- BATEMAN, T. A Practical Synopsis of Cutaneous Diseases, according to the arrangement of Dr. Willan, exhibiting a concise view of the diagnostic symptoms, and the method of treatment. Plate. 8vo.
- BAUDELOCQUE, A. C. Treatise on Puerperal Peritonitis, to which was awarded the Prize of the Royal Society of Medicine of Bordeaux. Translated by G. S. Bedford, M.D. 8vo.
 — An Abridgment of Heath's Translation of Baudelocque's Midwifery: with Notes by William P. Dewees, M.D. Plates. 4vo.

- BAYLE, A. L. J. & H. HOLLARD. A Manual of General Anatomy, containing a concise description of the elementary tissues of the Human Body. Translated by S. D. Gross, M.D. 8vo.
- BAYLE, A. L. J. An Elementary Treatise on Anatomy. Translated by A. S. Doane, M.D. 18mo.
- BAYNTON, T. Descriptive Account of a New Method of treating Old Ulcers of the Legs. 8vo.
- BEACH, W. American Practice of Medicine; being a Treatise on the Character, Causes, Symptoms, and Treatment of Diseases of Men, Women, and Children, of all Climates, on Vegetable or Botanical Principles, &c. Many engravings. 3 vols. 8vo.
- The Family Physician; or the Reformed System of Medicine, on Vegetable or Botanical Principles; being a Compendium of the "American Practice." Many engravings. 8vo.
- BEALE, L. J. A Treatise on the Distortions and Deformities of the Human Body: exhibiting a concise view of the nature and treatment of the principal Malformations and Distortions of the Chest, Spine, and Limbs. 4 plates. 8vo.
- BEAUMONT, W. Experiments and Observations on the Gastric Juice, and the Physiology of Digestion. Engravings. 8vo.
- BECHE, H. T. DE LA. Geological Manual. With engravings. 8vo.
- BECK, T. R. & J. B. Elements of Medical Jurisprudence. 2 vols. 8vo.
- BECLARD, P. A. Elements of General Anatomy, or a Description of every kind of Organs composing the Human Body. With Notes by Joseph Tognò, M.D. 8vo.
- BEDDOES, T. Contributions to Physical and Medical Knowledge, principally from the West of England. 8vo.
- Observations on the Nature and Cure of Calculus, Sea Scurvy, Consumption, Catarrh, and Fever. 8vo.
- BEDINGFIELD, J. A Compendium of Medical Practice, illustrated by interesting and instructive cases, &c. With Notes by Stephen W. Williams, M.D. 8vo.
- BEGIN, L. J. The French Practice of Medicine: being a translation of L. J. Begin's Treatise on Therapeutics, with notes and observations illustrative of the treatment of diseases in the climate of North America: by Xavier Tessier. 8vo.
- Application of the Physiological Doctrine to Surgery. Translated by W. S. Reynolds, M.D. 8vo.
- BELINAYE, H. Compendium of Lithotripsy; or an account of removal of the Stone from the Bladder without incision; with a series of statistical tables, and numerous wood-cuts representing the most important instruments and improvements to the present time. 8vo.
- BELL, BENJAMIN. A Treatise on Gonorrhœa Virulenta, and Lues Venerea. 8vo.
- BELL, CHARLES. A System of Operative Surgery, founded on the Basis of Anatomy. Many engravings. 2 vols. 8vo.
- A System of Dissections, explaining the Anatomy of the Human Body, with the manner of displaying the parts, the distinguishing the natural from the diseased appearances, and pointing out to the Student the objects most worthy of attention, during a course of dissections. 2 vols. 18mo.
- The Hand, its Mechanism and Vital Endowments, as evincing design. Engravings. A Bridgewater Treatise. 12mo.
- The Nervous System of the Human Body, embracing the Papers delivered to the Royal Society on the subject of the Nerves. 10 Plates. 8vo.
- The Anatomy and Philosophy of Expression, as connected with the Fine Arts. Many very fine engravings. 8vo.
- A Series of Engravings explaining the Course of the Nerves. 9 plates. 4to.
- Engravings of the Arteries; illustrating the Anatomy of the Human Body, and serving as an introduction to the Surgery of the Arteries. 10 coloured plates. 8vo.
- Observations on Injuries of the Spine and of the Thigh Bone. 9 plates. 4to.
- Institutes of Surgery; arranged in the order of the Lectures delivered in the University of Edinburgh. 8vo.

- BELL, JOHN & CHARLES. The Anatomy and Physiology of the Human Body. Numerous engravings. 3 vols. 8vo.
- BELL, JOHN. The Principles of Surgery as they relate to Wounds, Ulcers, Fistulæ, Aneurisms, Wounded Arteries, Fractures of the Limbs, Tumours, the Operations of Trepan and Lithotomy. With commentaries, &c., by Charles Bell. Many engravings and coloured plates. 4 vols. 8vo.
- Discourses on the Nature and Cure of Wounds. Plates. 8vo.
- Engravings of the Bones. 12 plates. 4to.
- Engravings of the Muscles and Joints. 16 plates. 4to.
- A Practical Dictionary of Materia Medica; including the composition, preparation, and uses of medicines; and a large number of extemporaneous formulae; together with important toxicological observations. 8vo.
- Health and Beauty: an Explanation of the Laws of Growth and Exercise; through which a pleasing contour, symmetry of form, and graceful carriage of the body are acquired; and the common deformities of the spine and chest prevented. 18mo.
- Lessons on the Human Frame, designed for Schools and Families. Engravings. 18mo.
- On Regimen and Longevity; comprising materia alimentaria, national dietetic usages, and the influence of civilization on health and the duration of life. 12mo.
- On Baths and Mineral Waters. 12mo.
- BELL, J., AND W. SPOKES. Lectures on the Theory and Practice of Physic. 2 v. 8vo.
- BELL, J., AND D. F. CONDIE. All the material Facts in the History of the Epidemic Cholera; being a Report of the College of Physicians of Philadelphia, to the Board of Health; and a full account of the causes, post mortem appearances, and treatment of the disease. 8vo.
- BELL, T. The Anatomy, Physiology, and Diseases of the Teeth. 11 plates. 8vo.
- BEMENT, C. N. American Poulterer's Companion: a Practical Treatise on Breeding, Rearing, Fattening, and General Management of the various species of Domestic Poultry. 68 engravings. 12mo.
- BERTIN, R. J. Treatise on the Diseases of the Heart and Great Vessels. Edited by J. Bouillaud, and translated by C. W. Chauncy, M.D. 8vo.
- BERZELIUS, J. J. The Kidneys and Urine: translated by M. H. Boyé and F. Leam. ing, M.D. 8vo.
- Use of the Blow-pipe in Chemistry and Mineralogy. Translated by J. D. Whitney. 4 plates. 8vo.
- BICHAT, X. General Anatomy applied to Physiology and Medicine: translated by George Hayward, M.D. 3 vols. 8vo.
- Additions to the General Anatomy, by P. A. Beclard; translated by G. Hayward, M.D. 8vo.
- Pathological Anatomy. The last course of X. Bichat, from an autographic manuscript of P. A. Beclard: with an account of the life and labours of Bichat, by F. G. Boisseau. Translated by Joseph Tognio. 8vo.
- Physiological Researches on Life and Death: with Notes by F. Magendie. Translated by F. Gold and G. Hayward, M.D. 8vo.
- A Treatise on the Membranes in general, and on different Membranes in particular: with a historical notice of the life and writings of the author, by M. Husson. Translated by John G. Coffin, M.D. 8vo.
- BIGELOW, H. J. Manual of Orthopedic Surgery, being a Dissertation which obtained the Boylston Prize for 1844, on the following Question: "In what cases and to what extent is the Division of Muscles, Tendons, or other parts proper for the relief of Deformity or Lameness?" 6 plates. 8vo.
- BIGELOW, J. Florida Bostoniensis: A Collection of Plants of Boston and its vicinity, with their generic and specific characters, principal synonyms, descriptions, places of growth, and time of flowering, &c., with a Glossary of Botanical Terms. 12mo.
- American Medical Botany, being a collection of the native medicinal plants of the United States, containing their Botanical History and Chemical Analysis,

- and Properties and Uses in Medicine, Diet, and the Arts. 60 coloured plates. 3 vols. 8vo.
- Elements of Technology, taken chiefly from a course of Lectures delivered at Cambridge, on the Application of the Sciences to the Useful Arts. Plates. 8vo.
- BILLARD, C. M. A Treatise on the Diseases of Infants, founded on recent clinical observations and investigations in Pathological Anatomy, made at the Hospice des Enfants trouvés: with a dissertation on the viability of the child. With Notes by Dr. Ollivier, of Angers; and an Appendix by James Stewart, M.D. 8vo.
- BILLING, A. First Principles of Medicine. 8vo.
- BIRD, G. Urinary Deposits, their Diagnosis, Pathology, and Therapeutical Indications. 8vo.
- BLACK, S. Clinical and Pathological Reports. 8vo.
- BLACKALL, J. Observations on the Nature and Cure of Dropsies, and particularly on the presence of the coagulate part of the blood in dropsical urine: with an Appendix containing several cases of Angina Pectoris, with Dissections, &c. 8vo.
- BLACKLOCK, A. Treatise on Sheep; with the best means for their Improvement, General Management, and Treatment of their Diseases. 8 plates. 18mo.
- BLANDIN, P. H. A Treatise on Topographical Anatomy; or the Anatomy of the Regions of the Human Body, considered in its relations with Surgery and Operative Medicine. 8vo. With 4to Atlas of 12 plates.
- BLANE, SIR GILBERT. Elements of Medical Logic, illustrated by practical proofs and examples. 12mo.
- BLEGEBOURGH, R. Facts and Observations respecting the Air-Pump Vapour-Bath in Gout, Rheumatism, Palsy, and other Diseases. Plate. 18mo.
- BLUNDELL, J. Lectures on the Principles and Practice of Midwifery: edited by Charles Severn, M.D. 8vo.
- Observations on some of the more important Diseases of Women: edited by Thomas Castle, M.D. 8vo.
- BOERHAAVE, H. Academical Lectures on the Theory of Physic: being a genuine translation of his Institutes and Explanatory Comment, collated and adjusted to each other, as they were dictated to his Students at the University of Leyden. 6 vols. 8vo.
- An Account of the Life and Writings of Herman Boerhaave, with portrait. 8vo.
- BOISSEAU, F. G. Physiological Pyretology; or a Treatise on Fevers, according to the principles of the New Medical Doctrine. Translated by J. R. Knox, M.D. 8vo.
- BOOTH, J. C., AND M. H. BOYE. Encyclopedia of Chemistry, Theoretical and Practical: presenting a complete and extended view of the present state of Chemical Science, with its applications to Medical Science, Agriculture, the Arts, and Manufactures. Numerous engravings. 8vo.
- BORREMAN'S Plate of the Nerves; 4½ feet long: on Roller.
- BOSTOCK, J. An Elementary System of Physiology. 3 vols. 8vo.
- BOSWELL, —. Poultry-Yard; or a Practical View of the best method of Selecting, Rearing, and Feeding the various species of Domestic Fowl. 18mo.
- BOUILLAUD, J. New Researches on Acute Articular Rheumatism in general; and especially on the Law of Coincidence of Pericarditis and Endocarditis with this Disease, as well as on the Efficacy of the method of treating it by repeated Blood-lettings at short intervals. Translated by James Kitchen, M.D. 8vo.
- BOUSSINGAULT, M. Rural Economy in its relations with Chemistry, Physics, and Meteorology. Translated by Geo. Lowe. 12mo.
- BOYER, BARON. A Treatise on Surgical Diseases, and the Operations suited to them. Translated by A. H. Stevens, M.D. Plates. 2 vols. 8vo.
- The Lectures of Boyer on Diseases of the Bones, arranged in a systematic Treatise, by A. Richerand. Translated by M. Farrell, M.D.: with Notes, &c., by J. Harishorne, M.D. Plates. 8vo.

- BRANDE, W. T.** A Dictionary of Science, Literature, and Art: comprising the History, Description, and Scientific Principles of every branch of Human Knowledge; with the derivation and definition of all the Terms in general use. Numerous engravings. 8vo.
- BREE, R.** A Practical Inquiry into Disordered Respiration; distinguishing the species of Convulsive Asthma, their causes and indications of cure. 8vo.
- BRERA, V. L.** A Treatise on Verminous Diseases, preceded by the Natural History of Intestinal Worms, and their origin in the Human Body. With Notes by J. Bartoli. Translated by John G. Coffin, M.D. 5 plates. 8vo.
- BREWSTER, D.** New Edinburgh Encyclopedia, conducted by David Brewster, LL.D., assisted by upwards of one hundred Gentlemen in Europe, most eminent in Science and Literature; and now improved for the greater satisfaction and better information of the People of the United States, in the Civil, Religious, and Natural History of their country; in American Biography; and in the great discoveries in Mechanics and the Arts. Many plates. 20 vols. 4to.
- A Treatise on Optics: with an Appendix by A. D. Bache, A.M. Numerous engravings. 12mo.
- BRICHETEAU, I.** Medical Clinics of the Hospital Necker; or, Researches and Observations on the Nature, Treatment, and Physical Causes of Diseases. 8vo.
- Bridgewater Treatises on the Power, Wisdom, and Goodness of God, as manifested in the Creation. Many plates. 7 vols. 8vo.
- BRIGIAM, A.** An Inquiry concerning the Diseases and Functions of the Brain, the Spinal Cord, and the Nerves. 12mo.
- Remarks on the Influence of Mental Cultivation, and Mental Excitement upon Health. 18mo.
- Observations on the Influence of Religion upon the Health and Physical Welfare of Mankind. 12mo.
- A Treatise on Epidemic Cholera; including a historical account of its Origin and Progress, to the present period. With Map showing its progress. 12mo.
- BROACKES, W.** Diseases of the Alimentary Canal and Constipation, treated Homœopathically: with Notes by G. Humphrey, M.D.: also an Essay on Homœopathic Diet. 18mo.
- BRODIE, SIR B. C.** Pathological and Surgical Observations on the Diseases of the Joints. 8vo.
- Lectures on the Diseases of the Urinary Organs. 8vo.
- BROUSSAIS, F. J. V.** On Irritation and Insanity, wherein the relations of the Physical with the Moral conditions of Man, are established on the basis of Physiological Medicine. Translated by Thomas Cooper, M.D. 8vo.
- Principles of Physiological Medicine, in the form of Propositions; embracing Physiology, Pathology, and Therapeutics, with Commentaries on those relating to Pathology. Translated by Isaac Hays, M.D., and R. E. Griffith, M.D. 8vo.
- History of Chronic Phlegmasiæ, or Inflammations, founded on Clinical experience and Pathological Anatomy, exhibiting a view of the different varieties and complications of these diseases, with their various methods of treatment. Translated by Isaac Hays, M.D., and R. E. Griffith, M.D. 2 vols. 8vo.
- BROWN, J.** The Elements of Medicine; or a Translation of the Elementa Medicinæ Brunonis: with Notes, Illustrations, and Comments. 8vo.
- BROWNE, W. A. F.** What Asylums were, are, and ought to be: being the substance of five Lectures delivered before the Managers of the Montrose Royal Lunatic Asylum. 12mo.
- BUCHAN, W.** Domestic Medicine, or a Treatise on the Prevention and Cure of Diseases, by Regimen and Simple Medicines: with observations on Sea-Bathing, and the use of Mineral Waters. 8vo.
- BUCKLAND, W.** Geology and Mineralogy considered with reference to Natural Theology. A Bridgewater Treatise. 69 plates. 2 vols. 8vo.
- BUEL, JESSE.** The Farmer's Companion; or Essays on the Principles and Practice of American Husbandry. Engravings. 12mo.

- BUIST, R.** American Flower-Garden Directory; containing Practical Directions for the Culture of Plants in the Flower-Garden, Hot-House, Green-House, Rooms, or Parlour Windows, for every month in the year; with a Description of the Plants most desirable in each; Nature of the Soil, &c. 12mo.
- The Rose Manual, containing accurate descriptions of all the finest varieties of Roses, properly classed in their respective families, their character and mode of culture, with directions for their propagation, and destruction of Insects. Engravings. 12mo.
- BULL, T.** Hints to Mothers for the Management of Health during the period of Pregnancy, and in the Lying-in Room; with an exposure of popular errors in connexion with those subjects. 18mo.
- BURDELL, H. & J.** Observations on the Structure, Physiology, Anatomy, and Diseases of the Teeth. 8 engravings. 8vo.
- BURKE, J. F.** The Dispensing Chemist's and Medical Pupil's Assistant; containing Latin directions, with their translations, for every species of Medical Prescription: with several forms of ordinary Prescriptions, and a succinct account of the properties and doses of the various Medicines in common use. 12mo.
- BURKE, W.** A Popular Compendium of Anatomy; or a concise and clear description of the Human Body: with the Physiology or Natural History of the various actions and functions of its different organs and parts: with plates. Containing also an Essay on Suspended Animation, with the proper means to be used for the recovery of drowned persons. 12mo.
- BURNE, J.** A Treatise on the Causes and Consequences of Habitual Constipation. 8vo.
- BURNS, ALLAN.** Observations on the Surgical Anatomy of the Head and Neck: with a Life of the Author, and additional cases and observations, by G. S. Pattison. 10 plates. 8vo.
- BURNS, JOHN.** Dissertations on Inflammation; containing Dissertations on some of the Laws of the Animal Economy, and the History, Causes, and Consequences of Simple Inflammation. 8vo.
- Principles of Midwifery, including the Diseases of Women and Children. With Notes, &c., by T. C. James, M.D. 12 plates. 8vo.
- Observations on Abortion; containing an account of the manner in which it takes place, the causes which produce it, and the methods of preventing or treating it. 12mo.
- Principles of Surgery; containing the doctrine and practice relating to Inflammation and its various consequences, Tumours, Aneurisms, Wounds, and the States connected with them; Surgical Anatomy of the Human Body, and its application to injuries and operations. 2 vols. 8vo.
- Popular Directions for the Treatment of the Diseases of Women and Children. 12mo.
- Anatomy of the Gravid Uterus. 8vo.
- BUSH, G.** A Treatise on the Malformation, Injuries, and Diseases of the Rectum and Anus. Plates. 8vo. with 4to vol. of plates.
- CABANIS, P. J. G.** An Essay on the Certainty of Medicine: translated by R. La Roche, M.D. 8vo.
- CALVERT, G.** A Practical Treatise on Hæmorrhoids or Piles, Strictures, and other important Diseases of the Rectum and Anus. Plate. 8vo.
- CAMPER.** The Works of the late Professor Camper, on the connexion between the science of Anatomy and the arts of Drawing, Painting, Statuary, &c. &c., containing a treatise on the natural difference of features in persons of different countries and periods of life; and on beauty, as exhibited in ancient sculpture; with a new method of sketching heads, national features, and portraits of individuals, with accuracy. By T. Cogan, M.D. 18 plates. 4to.
- CARLISLE, SIR A.** The Means of preserving Health and prolonging Life, applied to Hereditary diseases; the affections of Children; and the disorders of Old Age: comprising the result of fifty years' experience, derived from Hospital and private practice. 8vo.

- CARMICHAEL, R. An Essay on Venereal Diseases, and the uses and abuses of Mercury in their treatment. With Notes by G. Emerson, M.D. 5 plates. 8vo.
- CARPENTER, G. W. Essays on some of the most important articles of the *Materia Medica*, comprising a full account of all the new proximate principles, and the popular medicines lately introduced in practice, detailing the formulas for their preparation, their habitudes and peculiarities, doses, and modes of administration, &c. &c. 8vo.
- CARPENTER, W. B. Principles of Human Physiology, with their chief applications to Pathology, Hygiene, and Forensic Medicine. Especially designed for the use of Students. With Notes by Meredith Clymer, M.D. 216 engravings. 8vo.
- A Popular Treatise on Vegetable Physiology. 64 engravings. 12mo.
- Dissertation on the Physiological Inferences to be deduced from the Structure of the Nervous System in the Invertebrated classes of Animals. 8vo.
- CASTLE, T. Introduction to Medical Botany; comprehending the Elements and Glossology of Botany; the Linnæan artificial and natural systems; the natural system of Jussieu; and several comprehensive tables of the properties, uses, and doses of Medicinal Plants. Coloured plates. 18mo.
- Essay on Poisons; embracing their Symptoms, Treatment, Tests, and Morbid Appearances. To which are added the Means for treating cases of Suspended Animation. 21 coloured plates. 18mo.
- A Manual of Surgery, founded upon the principles and practice lately taught by Sir Astley Cooper, Bart. F.R.S., and Joseph Henry Green, F.R.S. 12mo.
- CAZENAVE, A., AND H. E. SCHEDEL. A Practical Synopsis of Cutaneous Diseases, from the most celebrated authors, and particularly from documents afforded by the Clinical Lectures of Dr. Bielt, Physician to the Hospital of St. Louis, Paris. 8vo.
- Celsi *Medicinæ libri octo ex recensione Leonardi Targæ*. With Notes by E. Milligan, M.D. 8vo.
- The same work in Latin and English: with a Life of the Author, Tables of Weights and Measures, Explanatory Notes, &c., by Alex. Lee, A.M. Surg. 8vo.
- CELSUS, A. C. Of Medicine, in eight books. Translated, with Notes, &c., by James Greive, M.D., and revised by G. Futvoye. 12mo.
- CHAILLY, M. A Practical Treatise on Midwifery. Translated and edited by Gunning S. Bedford, M.D. 216 engravings. 8vo.
- CHAPMAN, N. Lectures on the more important Diseases of the Thoracic and Abdominal Viscera. 8vo.
- Lectures on the more important Eruptive Fevers, Hæmorrhages, and Dropsies, and on Gout and Rheumatism. 8vo.
- Discourses on the Elements of Therapeutics and *Materia Medica*. 2 v. 8vo.
- CHAPTAL, J. A. Chemistry applied to Agriculture. 12mo.
- CHASE, H. Treatise on the Radical Cure of Hernia by Instruments. 32 engravings. 8vo.
- CHAVASSE, P. H. Advice to Wives on the management of themselves, during the periods of Pregnancy, Labour, and Suckling. 18mo.
- Advice to Mothers on the management of their offspring, during the periods of Infancy, Childhood, and Youth. 18mo.
- CHELIUS, J. M. A System of Surgery. With Notes, &c., by John F. South, Surg., and References to American Surgery by George W. Norris, M.D. 2 vols. 8vo.
- CHESELDEN, W. Anatomy of the Human Body. 40 plates. 8vo.
- Engravings of the Human Bones, selected and reduced from the large Plates of Cheselden: with explanations, &c. 14 plates. 12mo.
- CHEYNE, G. Practical Rules for the Restoration and Preservation of Health; and the best means of invigorating and prolonging Life. 18mo.
- CHEYNE, J. Essays on the Diseases of Children, with Cases and Dissections. 2 vols. 12mo.
- CHITTY, J. A Practical Treatise on Medical Jurisprudence, with so much of Anatomy, Physiology, Pathology, and the Practice of Medicine and Surgery, as are essential to be known by members of Parliament, lawyers, coroners, magistrates,

- officers in the army and navy, and private gentlemen ; and all the Laws relating to Medical Practitioners. Engravings. 8vo.
- Cholera. Report of the Royal Academy of Medicine, to the Minister of the Interior, upon the Cholera-Morbus: published by order of the French Government. Translated by John W. Sterling, M.D. 12mo.
- Report on Spasmodic Cholera, prepared by a committee under the direction of the Counsellors of the Massachusetts Medical Society. 2 maps. 8vo.
- CHRISTISON, R. A Treatise on Poisons in relation to Medical Jurisprudence, Physiology, and the Practice of Physic. 8vo.
- Granular Degeneration of the Kidneys, and its connection with Dropsy, Inflammations, and other Diseases. 8vo.
- CHURCHILL, F. The Diseases of Females, including those of Pregnancy and Child-bed. With Notes by Robert M. Huston, M.D. Engravings. 8vo.
- Theory and Practice of Midwifery. With Notes and Additions by Robert M. Huston, M.D. 116 engravings. 8vo.
- CIVALE, On the Medical and Prophylactic Treatment of Stone and Gravel: with a memoir on the Calculi of Cystine. Translated by Henry M. Smith, M.D. 8vo.
- CLARIDGE, R. T. Hydropathy ; or the Cold Water Cure, as practised by Vincent Priessnitz, at Graefenberg, Silesia, Austria. 8vo.
- CLARK, BRACY. Hippodomania, or the True Structure, Laws, and Economy of the Horse's Foot : also Podophthora, or a ruinous defect in the principle of the common shoe detected, and demonstrated by experiments ; with a proposition for a new principle of shoeing, &c. 13 plates. 4to
- CLARK, SIR J. The Sanative Influence of Climate: with an account of the best places of resort for invalids in England, the South of Europe, &c. 8vo.
- A Treatise on Pulmonary Consumption ; comprehending an inquiry into the causes, nature, prevention, and treatment of Tuberculous and Scrofulous Diseases in general. 8vo.
- CLARKE, SIR C. M. Observations on those Diseases of Females which are attended by Discharges. Plates. 2 vols. 8vo.
- CLARKE, SIR A. The Young Mother's Assistant ; or a Practical Guide for the prevention and treatment of the Diseases of Infants and Children. 12mo.
- An Essay on Diseases of the Skin ; containing practical observations on Sulphureous Fumigations, in the cure of Cutaneous complaints. 12mo.
- CLARKE, J. A Treatise on the Mulberry Tree and Silkworm ; and on the Production and Manufacture of Silk. Engravings. 12mo.
- CLATER, F. Every Man his own Cattle Doctor: containing the Causes, Symptoms, and Treatment of all the Diseases incident to Oxen, Sheep, and Swine ; and a sketch of the Anatomy and Physiology of Neat Cattle. Edited by William Youatt. With Additions by J. S. Skinner. Numerous illustrations.
- CLATER, F. & J. Every Man his own Farrier ; containing the causes, symptoms, and most approved methods of cure of the Diseases of Horses. With Additions by J. S. Skinner. 12mo.
- CLEGHORN, F. Observations on the Epidemical Diseases of Minorca. With Notes by Benjamin Rush, M.D. 12mo.
- CLENDINNING, J. The Croonian Lectures for 1838. Experiments and Observations relating to Pathology and Pathological Relations of the Heart. 8vo.
- CLUTTERBUCK, H. Lectures on Blood-Letting ; delivered at the General Dispensary, Aldersgate Street. 8vo.
- COATES, R. Popular Medicine ; or Family Adviser ; consisting of outlines of Anatomy, Physiology, and Hygiene, with such hints on the Practice of Physic, Surgery, and the Diseases of Women and Children, as may prove useful in Families when regular Physicians cannot be procured. 8vo.
- Physiology for Schools. 57 engravings. 12mo.
- COCKS, W. P. A Concise Treatise on Operative Surgery, describing the methods adopted by the English, Continental, and American Surgeons ; for the use of junior practitioners and students. 12 plates. 8vo.

- COLHOUN, S. Essay on Suspended Animation. 12mo.
- COLLES, A. Treatise on Surgical Anatomy. With Notes by J. P. Hopkinson, M.D. 8vo.
- Lectures on the Theory and Practice of Surgery. Edited by Simon M'Coy, Esq., F.R.C.S.I. 8vo.
- Practical Observations on the Venereal Disease, and on use of Mercury. 8vo.
- COLLIN, M. Manual for the use of the Stethoscope: a short Treatise on the different methods of investigating the Diseases of the Chest. Translated by W. N. Ryland, M.D. 4 plates. 12mo.
- COLLINS, R. A Practical Treatise on Midwifery, containing the result of 16,654 Births, occurring in the Dublin Lying-in Hospital, during a period of 7 years. 8vo.
- COLOMBAT, DE L'ISERE. A Treatise on the Diseases and Special Hygiene of Females: translated by Charles D. Meigs, M.D. 49 engravings. 8vo.
- A Treatise on the Diseases and Hygiene of the Organs of the Voice: translated by J. F. W. Lane, M.D. 18mo.
- COMBE, G. Constitution of Man considered in relation to external objects. 12mo.
- A System of Phrenology. With upwards of 100 engravings. 12mo.
- Observations on Mental Derangement, being an application of the Principles of Phrenology to the Elucidation of the Causes, Symptoms, Nature, and Treatment of Insanity. 8vo.
- COMBE, A. The Principles of Physiology applied to the Preservation of Health, and to the Improvements of Physical and Mental Education. 18mo.
- A Treatise on the Physiological and Moral Management of Infancy. For the use of Parents. 18mo.
- Physiology of Digestion, considered with relation to the Principles of Dietetics. 18mo.
- COMSTOCK, J. L. Elements of Chemistry, in which the recent Discoveries in the Science are included, and its Doctrines familiarly explained. 76 engravings. 12mo.
- A System of Natural Philosophy: in which the principles of Mechanics, Hydrostatics, Hydraulics, Pneumatics, Acoustics, Optics, Astronomy, Electricity, Magnetism, Steam Engine, Electro-Magnetism, Electrotypes, Photography, and Daguerreotype, are familiarly explained. 258 engravings. 12mo.
- Outlines of Physiology, both Comparative and Human; in which are described the Mechanical, Animal, Vital, and Sensorial Organs and Functions, &c. Also the application of these principles to Muscular Exercise, and Female Fashions and Deformities. 137 engravings. 12mo.
- An Introduction to Mineralogy. 186 illustrations. 12mo.
- Outlines of Geology: intended as a Popular Treatise on the most interesting parts of the Science. 97 illustrations. 12mo.
- An Introduction to the Study of Botany, including a treatise on Vegetable Physiology and descriptions of the most common Plants in the Middle and Northern States. 247 engravings. 12mo.
- CONDIE, D. F. A Practical Treatise on the Diseases of Children. 8vo.
- CONOLLY, J. An Inquiry concerning the Indications of Insanity, with Suggestions for the better protection and care of the Insane. 8vo.
- CONWELL, W. E. E. Observations, chiefly on Pulmonary Disease in India; and an Essay on the Use of the Stethoscope. Also, a Code of Medical Regulations for East India Company's Establishment of Surgeons, &c. &c. 4to.
- COOK, C. T. Observations on the Efficacy of White Mustard Seed. 18mo.
- COOKE, J. A Treatise on Nervous Diseases. 8vo.
- COOPER, SIR ASTLEY. The Anatomy and Surgical Treatment of Abdominal Hernia. Edited by C. A. Key. 26 plates. 8vo.
- Observations on the Structure and Diseases of the Testis. Edited by B. B. Cooper, F.R.S. 24 plates. Including the Anatomy of the Thymus Gland. 5 plates. 8vo.
- The Anatomy and Diseases of the Breast. To which are added his various Surgical Papers, now first published in a collected form. 35 plates. 8vo.
- The Principles and Practice of Surgery, founded on the most extensive Hos-

- pital and private Practice, during a period of nearly fifty years. Edited by A. Lee, M.D. 32 plates. 3 vols. 8vo.
- A Treatise on Dislocations and Fractures of the Joints. Edited by B. B. Cooper, F.R.S. Numerous engravings. 8vo.
- Lectures on the Principles and Practice of Surgery, with additional Notes and Cases. By F. Tyrrell, Esq. 4 plates.
- On Spermatocoele, or Variocele of the Spermatic Cord. 8vo.
- COOPER, SIR A., AND B. TRAVERS. Surgical Essays; on Dislocations, Iritis, Ligature of Aorta, Phymosis and Paraphymosis, Exostosis, Wounds and Ligatures of Veins, Unnatural apertures in the Urethra, Encysted Tumours, &c. 21 plates. 8vo.
- COOPER, SAMUEL. The First Lines of the Theory and Practice of Surgery; including the Principal Operations. With Notes and Additions, by Willard Parker, M.D. 2 vols. 8vo.
- A Dictionary of Practical Surgery: comprehending all the most interesting improvements, from the earliest times to the present period: an account of the Instruments and Remedies employed in Surgery, the Etymology and Signification of the principal Terms, &c. &c. With a Supplementary Index by D. M. Reese, M.D. 8vo.
- Same work, with Notes by John Syng Dorsey, M.D. 2 vols. 8vo. 1816.
- A Treatise on the Diseases of the Joints. 18mo.
- COPELAND, T. Observations on some of the Principal Diseases of the Rectum and Anus; particularly Stricture of the Rectum, the Hæmorrhoidal Excrescence, and the Fistula in Ano. 12mo.
- COPLAND, J. A Dictionary of Practical Medicine; comprising General Pathology, the Nature and Treatment of Diseases, Morbid Structures, &c. Edited by Charles A. Lee, M.D. 3 vols. 8vo. (*In course of publication*.)
- CORFEE, G. A Popular Treatise on the Kidney; its hitherto unknown Functions, and its Diseases, in connection with the Circulating Animal Oils, &c. Plate. 8vo.
- CORVISART, J. N. An Essay on the Organic Diseases and Lesions of the Heart and Great Vessels. Translated by Jacob Gates, M.M.S.S. 8vo.
- COSTER, J. The Practice of Medicine, according to the Principles of the Physiological Doctrine.
- Manual of Operative Surgery: translated by George Fife, M.D. 12mo.
- Manual of Surgical Operations; containing the new methods of operating devised by Lisfranc. Translated by John D. Godman, M.D. 12mo.
- COULSON, W. On Deformities of the Chest. 4 plates. 12mo.
- On Diseases of the Bladder. 8vo.
- COURTENAY, F. B. A Practical Treatise on the Causes, Nature, and Treatment of Strictures of Urethra; with a Review of the Different Modes of Cure, &c. 8vo.
- COX, J. M. Practical Observations on Insanity: with Remarks on Medical Jurisprudence as connected with Diseased Intellect. 12mo.
- COXE, E. J. A Practical Treatise on Medical Inhalation, with numerous cases demonstrating the curative powers of the local application of various remedies in Bronchitis, Consumption, and other diseases of the Respiratory Organs, &c. 18mo.
- CRAIGIE, D. Elements of Anatomy, General, Special, and Comparative. 14 plates. 4to.
- CROLY, H. Irish Medical Directory for 1843, including Notices of the Literary and Scientific Institutions of Ireland. With Notes, Historical, Biographical, and Bibliographical. 18mo.
- CROWTHER, B. Practical Remarks on Insanity; to which is added a Commentary on the Dissection of the Brains of Maniacs; with some account of Diseases Incident to the Insane. 8vo.
- CRUVEILHIER, J. The Anatomy of the Human Body. Edited by Granville Sharp Pattison, M.D. 302 engravings. 8vo.
- CULLEN, W. First Lines of the Practice of Physic. With Notes, &c., by Charles Caldwell, M.D. 2 vols. 8vo.
- Practice of Physic: with Notes, &c., by R. J. Thornton, M.D.

- CULLEN, W. Treatise of Materia Medica. With large additions, including many new articles, &c., by Benj. Smith Barton, M.D. 2 vols. 8vo.
- A Synopsis of Nosology. Translated by John Thompson, M.D. To which is added Willan's Classification of Cutaneous Diseases. 12mo.
- CULPEPER, N. The Complete English Family Physician: being a Botanical and Astrological Practice of Physic. With Additions by G. A. Gordon, M.D. Very numerous coloured plates. 3 vols. 8vo.
- CURLING, T. B. A Practical Treatise on the Diseases of the Testis, and of the Spermatie Cord, and Scrotum. Edited by P. B. Goddard. Engravings. 8vo.
- A Treatise on Tetanus, being the Essay for which the Jacksonian Prize for the year 1834 was awarded, by the Royal College of Surgeons, London. 8vo.
- CURRIE, JAMES. Memoir of his Life, Writings, and Correspondence. Edited by his son William Wallace Currie. 2 vols. 8vo.
- CURTIS, J. H. A Treatise on the Physiology and Diseases of the Ear; containing a Comparative View of its Structure and Functions, and of its various Diseases, arranged according to the Anatomy of the organ, or as they affect the external, the intermediate, and the internal Ear. Plate. 8vo.
- Observations on the Preservation of Health, in Infancy, Youth, Manhood, and Age; with the best means of improving the Moral and Physical condition of Man, prolonging Life, and promoting Human Happiness. 12mo.
- CURTIS, J. T., AND J. LILLIE. Epitome of Homœopathic Practice. 18mo.
- CUTBUSH, E. Observations on the Means of preserving the Health of Soldiers and Sailors, &c., with Remarks on Hospitals and their internal arrangement. 8vo.
- CUTLER, T. The Surgeon's Practical Guide in Dressing, and in the Methodic Application of Bandages. Numerous engravings. 18mo.
- CUVIER, BARON. The Animal Kingdom arranged in conformity with its Organization. Translated by H. M'Murtrie, M.D. 20 plates, 4 vols. 8vo.
- Cyclopedia of Practical Medicine: comprising Treatises on the Nature and Treatment of Diseases, Materia Medica, and Therapeutics, Medical Jurisprudence, &c., &c. Edited by John Forbes, M.D., A. Tweedie, M.D., and J. Conolly, M.D. Revised by R. Dunglison, M.D. 4 vols. 8vo.
- Cyclopedia of Anatomy and Physiology: consisting of a series of Dissertations on the most important subjects of Human Anatomy, General, Surgical, and Morbid; of Physiology; of Comparative Anatomy; and of Animal Chemistry. Edited by R. B. Todd, M.D. Numerous engravings. 8vo. (In course of publication.)
- Cyclopedia of Practical Surgery; comprising a series of Dissertations on Operative Medicine. Edited by William B. Costello, M.D. Numerous engravings. 8vo. (In course of publication.)
- DANA, S. L. A Muck Manual, for Farmers. 18mo.
- DANA, J. D. A System of Mineralogy, including an extended Treatise on Crystallography. Numerous illustrations. 8vo.
- DARRACH, W. Drawings of the Anatomy of the Groin; with Anatomical Remarks. 4 plates. 8vo.
- DARWIN, E. Zoonomia; or the Laws of Organic Life. 2 vols. 8vo.
- DAVIDSON, W., AND A. HUDSON. Essays on the Sources and Mode of Action of Fever. 8vo.
- DAVIES, J. Selections in Pathology and Surgery; or an Exposition of the nature and treatment of Local Disease; exhibiting New Pathological Views; and pointing out an important practical improvement. 8vo.
- DAVIS, D. D. Acute Hydrocephalus, or Water in the Head, an Inflammatory Disease, and curable equally by the same means with other Diseases of Inflammation. 8vo.
- Elements of Obstetric Medicine; with the Description and Treatment of some of the principal Diseases of Children. 8vo.: with Atlas of 70 plates 4to.
- DAVY, J. Researches, Physiological and Anatomical. 8vo.
- DE FOE, DANIEL. History of the Plague in London, in 1665. 12mo.

- DELEUZE, J. P. F. Practical Instruction in Animal Magnetism. 12mo.
- DENDY, W. C. The Book of the Nursery: Precepts for the management of Infants, and for the prevention and domestic treatment of the Diseases incidental to Childhood. 18mo.
- Practical Remarks on Diseases of the Skin, on the external Signs of Disorder, and on the Constitutional Peculiarities during Infancy and Childhood. 8vo.
- DENMAN, T. Aphorisms on Practical Obstetrics; embracing the Application and Use of the Forceps and Vectis, Prematural Labours, Labours attended with Hæmorrhage, Convulsions, &c. Engravings. 32mo.
- An Introduction to the Practice of Midwifery: with a Biographical Sketch of the Author. Notes and Emendations by John W. Francis, M.D. 17 plates. 8vo.
- DE SALLE, E. A Synoptical Table of the Mineral and Vegetable Poisons, together with the Symptoms which they produce, the Treatment required, and the Reagents that recognise them. To which are added the Morbid Appearances observed on Dissection: by William Bennet, M.D.
- DESLANDES, L. Manhood: the causes of its premature decline; with directions for its perfect restoration; addressed to those suffering from the destructive effects of excessive indulgence, solitary habits, &c. 18mo.
- DESRUVELLES, H. M. J. Memoir on the Treatment of Venereal Diseases without Mercury, employed at the Military Hospital of the Val-de-Grace. With Notes, &c., by G. J. Guthrie, Esq. 8vo.
- DESAULT, P. J. A Treatise on Fractures, Luxations, and other affections of the Bones. Edited by Xav. Bichat, and translated by Charles Caldwell, M.D. Plates. 8vo.
- The Surgical Works, or Statement of the Doctrine and Practice of P. J. Desault, Surgeon-in-chief of the Great Hospital of Humanity, at Paris, by Xavier Bichat: translated by E. D. Smith, M.D. Vol. 1, Diseases of the Soft Parts. Vol. 2, Diseases of the Urinary Passages. Plates. 8vo.
- DETMOLD, W. An Essay on Club-Foot, and some analogous Diseases. 4 plates. 8vo.
- DEWEES, W. P. A Compendious System of Midwifery, chiefly designed to facilitate the inquiries of those who may be pursuing this branch of study. 18 plates. 8vo.
- A Treatise on the Diseases of Females. 12 plates. 8vo.
- A Treatise on the Physical and Medical Treatment of Children. 8vo.
- An Essay on the Means of Lessening Pain, and facilitating certain cases of Difficult Parturition. 8vo.
- A Practice of Physic, comprising most of the Diseases not treated of in "Diseases on Females," and "Diseases of Children." 8vo.
- DICK, R. Derangements, Primary and Reflex, of the Organs of Digestion. 8vo.
- Same work, with an Addition containing Notices of Brandy and Water, Homœopathy, the Cold-Water Treatment, Liebig's new views in Animal Chemistry, with a new method of treating cases of Functional Neuralgia. 8vo.
- DICKSON, S. Fallacies of the Faculty; being the Spirit of the Chrono Thermal System. 8vo.
- DICKSON, S. H. Essays on Pathology and Therapeutics; being the substance of the Course of Lectures delivered in the Medical College of South Carolina. 2 vols. 8vo.
- DIX, J. H. Treatise on Strabismus, or Squinting, and the new mode of treatment. Plate. 12mo.
- DOANE, A. S. Surgery Illustrated: compiled from the Works of Cutler, Hind, Velpeau, and Blasius. 52 plates. 8vo.
- DORSEY, J. SYNG. Elements of Surgery, for the use of Students: with Notes by J. Randolph, M.D. 23 plates. 2 vols. 8vo.
- DOWNING, A. J. Treatise on the Theory and Practice of Landscape Gardening, adapted to North America, with a view to the improvement of Country Residences. Engravings. 8vo.
- Designs for Cottage Residences adapted to North America, including Elevations and Plans of the Buildings, and Designs for Laying out Grounds, &c. Engravings. 8vo.

- DOWNING, A. J. Fruits and Fruit Trees of America : or the Culture, Propagation, and Management, in the Garden and Orchard, of Fruit Trees generally ; with Descriptions of all the finest varieties of Fruit, Native and Foreign, cultivated in this country. 212 engravings. 12mo.
- DRAPER, JOHN WILLIAM. A Treatise on the Forces which produce the Organization of Plants : with an Appendix, containing several Memoirs on Capillary Attraction, Electricity, and the Chemical Action of Light. 4 plates. 4to.
- DRUITT, R. The Principles and Practice of Modern Surgery : with Notes, &c., by Joshua B. Flint, M.D. 153 engravings. 8vo.
- DUCAMP, T. A Treatise on Retention of Urine, caused by Strictures in the Urethra ; and of the means by which Obstructions of this canal may be effectually removed. Translated by William M. Herbert, M.D. 5 plates. 8vo.
- DUFFIN, E. W. The Influence of Physical Education in producing and confirming in Females, Deformity of the Spine. Plate. 12mo.
- DUNCAN, A. Observations on the distinguishing symptoms of three different species of Pulmonary Consumption, the Catarrhal, the Apostematous, and the Tuberculous ; with some remarks on the Remedies and Regimen best fitted for the prevention, removal, or alleviation of each species. With an Appendix, on the preparation and use of Lactucarium, or Lettuce-Opium. 12mo.
- DUNGLISON, R. The Medical Student ; or Aids to the Study of Medicine. 12mo.
- Human Health : or the Influence of Atmosphere and Locality ; Change of Air and Climate, Seasons, Food, Clothing, Bathing, and Mineral Springs, Exercise, Sleep, Corporeal and Intellectual Pursuits, &c. &c., on Healthy Man ; constituting Elements of Hygiene. 8vo.
- Observations on the Condition of the Insane Poor. 8vo.
- New Remedies ; Pharmaceutically and Therapeutically considered. 8vo.
- Human Physiology. Upwards of 300 illustrations. 2 vols. 8vo.
- The Practice of Medicine : a Treatise on Special Pathology and Therapeutics. 2 vols. 8vo.
- General Therapeutics and Materia Medica, adapted for a Medical Text Book. 2 vols. 8vo.
- A Dictionary of Medical Science, containing a concise account of the various subjects and terms ; with the French and other Synonymes ; Notices of Climate, and of celebrated Mineral Waters, Formulæ for various officinal and empirical preparations, &c. 8vo.
- DUNSFORD, H. The Practical Advantages of Homœopathy, illustrated by numerous cases. 8vo.
- DUPARCQUE, F. A Treatise on the Functional and Organic Diseases of the Uterus. Translated by Joseph Warrington, M.D. 8vo.
- DUPUYTREN, BARON. Clinical Lectures on Surgery, delivered at Hotel Dieu, in 1832. Translated by A. Sydney Doane, M.D. 8vo.
- EATON, A., AND J. WRIGHT. North American Botany, comprising the native and common cultivated Plants north of Mexico ; genera arranged according to the artificial and natural methods. 8vo.
- EBERLE, J. A Treatise on the Theory and Practice of Medicine. 2 vols. 8vo.
- Notes of Lectures on the Theory and Practice of Medicine, delivered in the Jefferson Medical College, Philadelphia. 12mo.
- A Treatise of the Materia Medica and Therapeutics. 8vo.
- A Treatise on the Diseases, and Physical Education of Children. 8vo.
- Eclectic and General Dispensatory ; comprehending a System of Pharmacy, Materia Medica ; the Formulæ of the London, Edinburgh, and Dublin Pharmacopœias ; Prescriptions of many eminent Physicians, and Receipts for the most common Empirical Medicines. By an American Physician. 8vo.
- EDWARDS, H. M. A Manual of Surgical Anatomy, containing a minute description of the parts concerned in Operative Surgery, with the Anatomical effects of Accidents, and Instructions for the performance of operations. Translated by W. Coulson : with Notes by James Webster, M.D. 12mo.

- EDWARDS, H. M. *Outlines of Anatomy and Physiology*: translated by J. F. W. Lane, M.D. 8vo.
- EDWARDS, H. M., and P. VAVASSEUR. *A Manual of Materia Medica and Pharmacy*, comprising a concise description of the Articles used in Medicine; their Physical and Chemical properties; the Botanical character of the Medicinal Plants; and the Formulæ for the principal Official Preparations of the American, Parisian, London, Dublin, Edinburgh, &c., Pharmacopœiæ, &c. &c. Translated by J. Togno, M.D., and E. Durand. 8vo.
- EDWARDS, W. F. *On the Influence of Physical Agents on Life*. Translated by Drs. Hodgkin and Fisher. With an Appendix containing Observations on Electricity, by Dr. Edwards, M. Pouillet, and Luke Howard, F.R.S. 8vo.
- ELLIOTSON, J. *The Principles and Practice of Medicine*. Edited by N. Rogers, M.D., and A. C. Lee. With Additions by Thomas Stewardson, M.D. 8vo.
- Numerous Cases of Surgical Operations without pain, in the Mesmeric State: with remarks upon the opposition of many members of the Royal Medical and Chirurgical Society and others, to the reception of the inestimable blessings of Mesmerism. 8vo. *Pamphlet*.
- ELLIS, B. *The Medical Formulary*; being a collection of Prescriptions derived from the Writings and Practice of many of the most eminent Physicians in America and Europe. With an Appendix containing the usual Dietetic preparations and antidotes for Poisons, &c. Revised by S. G. Morton, M.D. 8vo.
- Encyclopedia Americana*: A Popular Dictionary of Arts, Sciences, Literature, History, Politics, and Biography. 13 vols. 8vo.
- ESQUIROL, E. *Mental Maladies: a Treatise on Insanity*. Translated by E. K. Huut, M.D. 8vo.
- EVANS, OLIVER. *The Young Millwright and Miller's Guide*. With Additions, by T. P. Jones. 28 plates. 8vo.
- EVANSON, R. T., and H. MAUNSELL. *A Practical Treatise on the Management and Diseases of Children*. With Notes by D. F. Condie, M.D. 8vo.
- EVEREST, T. R. *A Popular View of Homœopathy*. With annotations and a brief survey of the progress and present state of Homœopathy in Europe: by A. G. Hull, M.D. 8vo.
- EVERS, P. *Student's Compendium of Comparative Anatomy*. 8vo.
- EWBANK, T. *A Descriptive and Historical Account of Hydraulic and other Machines for Raising Water, ancient and modern*; with observations on various subjects connected with the Mechanic Arts; including the progressive development of the Steam Engine, &c. 286 engravings. 8vo.
- EWELL, T. *Statement of Improvements in the Theory and Practice of the Science of Medicine*. 8vo.
- FAITHORNE, J. *Facts and Observations on Liver Complaints, Bilious Disorders in general, and on those derangements of that important organ, and of its immediate or intimate connections, which most sensibly influence the Biliary Secretion*. 8vo.
- FARADAY, M. *Chemical Manipulations. Instruction to Students on the methods of performing experiments of demonstration or research with accuracy and success*. With Additions by J. R. Mitchell, M.D. Illustrations. 8vo.
- FAWDINGTON, T. *A Case of Melanosis, with general observations on the Pathology of this interesting disease*. 8 coloured plates. 8vo.
- FERGUSON, W. *A System of Practical Surgery*. With Notes, &c., by George W. Norris, M.D. 246 engravings. 8vo.
- FESSENDEN, T. G. *The Complete Farmer and Rural Economist*; containing a compendious epitome of the most important branches of Agriculture and Rural Economy. Engravings. 12mo.
- *New American Gardener, containing practical directions on the Culture of Fruits and Vegetables*; including Landscape and Ornamental Gardening. 12mo.
- FISHER, J. D. *Description of the Distinct, Confluent, and Inoculated Small Pox, Varioloid Disease, Cow Pox, and Chicken Pox*. 13 coloured plates. 4to.
- FISHER, J. *Observations on the Character and Culture of the European Vine, du-*

- ring a residence of five years in the Vine-growing districts of France, Italy, and Switzerland. With the Manual of the Swiss Vigneron. 12mo.
- FITCH, S. S. A Treatise on the Uses of the Lungs, and on the Causes and Cure of Pulmonary Consumption. 12mo.
- FLOOD, S. Exposition of the Pathology and Treatment of Tubercular Phthisis. 12mo.
- FLOURENS, PROF. Cours sur la Generation, l'Ovologie, et l'Embryologie fait au Museum d'Histoire Naturelle en 1836. Recueilli par M. Deschamps. 10 plates. 4to.
- FORD, E. Observations on Disease of the Hip Joint: with Remarks on White Swellings of the Knee, Caries of the Joint of the Wrist, and other similar complaints. 8 plates. 8vo.
- FORDYCE, G. Five Dissertations on Fever. 8vo.
- Formulæ Selectæ; or a Collection of Prescriptions of eminent Physicians, and the most celebrated Patent Medicines: with a Syllabic view of Mineral Waters and Poisons. 8vo.
- FORRY, S. Climate of the United States, and its Endemic Influences. 2 plates. 8vo.
- FOTHERGILL, J. The Works of John Fothergill, M.D., with an account of his Life by John Coakley Lettsom. 11 plates. 4to.
- FOWNES, G. Elementary Chemistry, Theoretical and Practical. With Additions by Robert Bridges, M.D. 162 illustrations. 12mo.
- Chemistry, as exemplifying the Wisdom and Beneficence of God. 12mo.
- FOX, JOSEPH. The Natural History and Diseases of the Human Teeth: containing the History of the Formation and Structure of the Teeth; the Symptomatic Diseases incidental to the first Dentition; the changes which take place during the second Dentition; and the treatment to prevent and remedy Irregularities in the arrangement of the Teeth. Also the History and Treatment of the Diseases of the Teeth; the Gums; and the Alveolar Processes; with the operations which they respectively require. To which are added Observations on other Diseases of the Mouth, and on the mode of fixing Artificial Teeth. 23 plates. 4to.
- FRECKLETON, G. Outlines of General Pathology. 8vo.
- FRESENIUS, C. R. Elementary Instruction in Chemical Analysis: with a Preface by Professor Liebig. Edited by J. L. Bullock. 12mo.
- FRICK, G. A Treatise on Diseases of the Eye; including the Doctrines and Practice of the most eminent modern Surgeons, and particularly those of Professor Beer. Plate. 8vo.
- GALL, F. J. Manual of Phrenology; being an Analytical summary of the system of Dr. Gall, on the Faculties of Man and the Functions of the Brain. 5 plates. 12mo.
- Analytical View of the System of Dr. Gall, on the Faculties of Man and the Functions of the Brain; or, Cranioscopy—in one plate, 32 inches long.
- GALL, VIMONT, and BROUSSAIS, on the Functions of the Cerebellum; translated by Combe. Engravings. 8vo.
- GALLUP, J. A. Outlines of the Institutes of Medicine; founded on the Philosophy of the Human Economy, in Health and in Disease. 2 vols. 8vo.
- GALT, JOHN M. Practical Medicine; illustrated by cases of the most important Diseases. 8vo.
- GAMAGE, W., JR. Account of the Fever which existed in Boston during the Autumn and Winter of 1817 and 1818. With Remarks on Typhus Fever. 8vo.
- GANNAL, J. N. History of Embalming, and of Preparations in Anatomy, Pathology, and Natural History; including an account of a new process for Embalming. With Notes, &c., by R. Harlan, M.D. 8vo.
- GERHARD, W. W. Lectures on the Diagnosis, Pathology, and Treatment of the Diseases of the Chest. Plate. 8vo.
- GIBERT, M. Changes of the Blood in Disease. Translated by John H. Dix, M.D. 8vo.
- GIBSON, J. M. A Condensation of Matter upon the Anatomy, Surgical Operations, and Treatment of Diseases of the Eye. 12 plates. 4to.

- GIBSON, W. Institutes and Practice of Surgery; being Outlines of a Course of Lectures. 34 plates. 2 vols. 8vo.
- Rambles in Europe: with Sketches of prominent Surgeons, Physicians, Medical Schools, Hospitals, Literary Personages, Scenery, &c. 12mo.
- GODDARD, P. B. Plates of the Arteries, with References; for the use of Medical Students. 12 coloured plates. 4to.
- Plates of the Cerebro-Spinal Nerves, with References; for the use of Medical Students. 12 plates. 4to.
- The Anatomy, Physiology, and Pathology of the Human Teeth; with the most approved methods of Treatment; including Operations, and the method of making and setting Artificial Teeth. 30 plates. 4to.
- GODMAN, J. D. Addresses delivered on various Public Occasions: with an Appendix on the injurious effects of Tight Lacing upon the Organs and Functions of Respiration, Circulation, Digestion, &c. 8vo.
- Anatomical Investigations; comprising descriptions of various Fasciæ of the Human Body: with an account of some irregularities of Structure and Morbid Anatomy, &c. 8 plates. 8vo.
- American Natural History: to which is added the "Rambles of a Naturalist," with a Biographical Sketch of the author. 51 plates. 2 vols. 8vo.
- GOLDSMITH, O. A History of the Earth, and Animated Nature. 85 plates. 4 vols. 8vo.
- GOOCH, R. A Practical Compendium of Midwifery. 8vo.
- An Account of the most important Diseases peculiar to Women. 8vo
- GOOD, J. M. The Study of Medicine. Improved from the Author's Manuscripts, and by reference to the latest advances in Physiology, Pathology, and Practice: by Samuel Cooper, M.D. With Notes by A. S. Doane, M.D. Also a Sketch of the History of Medicine from its origin to the commencement of the 19th century, by J. Bostock, M.D. 2 vols. 8vo.
- The Book of Nature. To which is prefixed a Sketch of the Author's Life. 8vo.
- GORMAN, J. B. Philosophy of Animated Existence; or Sketches of Living Physics: with discussions of Physiology Philosophical. To which is added a brief Medical Account of the Middle Regions of Georgia. 8vo.
- GOUPIL, J. M. A. An Exposition of the Principles of the New Medical Doctrine, with an Analysis of the Theses sustained on its different parts. To which is appended a short Essay on Leeches, by J. C. Nott, M.D. 8vo.
- GRAHAM, S. A Lecture to Young Men on Chastity. 18mo.
- Lectures on the Science of Human Life. Many engravings. 2 vols. 12mo
- GRAHAM, T. Elements of Chemistry: including the application of the Science in the Arts. With Notes, &c., by Robert Bridges, M.D. 133 illustrations. 8vo.
- GRAHAM, T. J. The Cold-Water System: an Essay exhibiting the real merits, and most safe and effectual employment, of this excellent system in Indigestion, Costiveness, Asthma, Cough, Consumption, Rheumatism, &c. 8vo.
- A Treatise on Indigestion; with observations on some painful complaints originating in Indigestion, as Tic Douloureux, Nervous Disorder, &c. 8vo.
- GRANVILLE, A. B. Counter Irritation, its principles and practice illustrated by one hundred cases of the most painful and important Diseases effectually cured by external applications. 8vo.
- GRAVES, ROBERT J. Clinical Lectures: with Notes and a Series of Lectures, by W. W. Gerhard, M.D. 8vo.
- Sketch of the origin and progress of Asiatic Cholera. 8vo.
- GRAY, ALONZO. Elements of Chemistry: containing the Principles of the Science, both experimental and theoretical. 169 engravings. 12mo.
- GRAY, ASA. Botanical Text Book: comprising an Introduction to Structural and Physiological Botany, and the Principles of Systematic Botany; with an account of the chief natural families of the Vegetable Kingdom, and notices of the principal officinal or otherwise useful Plants. Numerous engravings. 12mo.

- GRAY, S. F. Elements of Pharmacy, and of the Chemical History of the Materia Medica. 8vo.
- Supplement to the Pharmacopœia. 8vo.
- GREEN, J. A Practical Compendium of the Diseases of the Skin, including a particular consideration of the more frequent and intractable forms of these affections: with cases. 8vo.
- GREENHOW, T. M. Cholera, as it recently appeared in the towns of Newcastle and Gateshead; including cases illustrative of its Physiology and Pathology, with a view to the establishment of sound principles of practice. 8vo.
- GREGORY, G. Elements of the Theory and Practice of Physic: with Notes and Additions adapted to the Practice of the United States, by N. Potter, M.D., and S. Colhoun, M.D. 2 vols. 8vo.
- GREGORY, J. Conspectus Medicinæ Theoreticæ. 2 vols. 8vo.
- Dissertation on the Influence of a Change of Climate in Curing Diseases. With Notes, &c., by W. P. C. Barton, M.D. 8vo.
- GRIFFITH, W. A Treatise on Hydrocephalus, or Water in the Brain; with the most successful modes of Treatment. 12mo.
- GRINDRON, R. B. Bacchus: an Essay on the Nature, Causes, Effects, and Cure of Intemperance. With Notes, &c., by Charles A. Lee, M.D. 12mo.
- GRISCOM, J. H. Animal Mechanism and Physiology; being a plain and familiar exposition of the Structure and Functions of the Human System. Engravings. 18mo.
- GROSS, S. D. Anatomy, Physiology, and Diseases of the Bones and Joints. 8vo.
- An Experimental and Critical Inquiry into the Nature and Treatment of Wounds of the Intestines; illustrated by engravings. 8vo.
- Elements of Pathological Anatomy; illustrated by numerous engravings—part coloured. 2 vols. 8vo.
- GUNN, J. C. Domestic Medicine, or Poor Man's Friend in the hours of affliction, pain, and sickness; pointing out in plain language, free from Doctors' terms, the Diseases of Men, Women, and Children, with the most approved means for their cure, &c., for the use of Families. With Descriptions of the Medicinal Roots and Herbs of the United States, and how they are to be used in the cure of Diseases. 8vo.
- GUTHRIE, G. J. Diseases and Injuries of the Arteries, with the Operations required for their cure. 8vo.
- Clinical Lectures on Compound Fractures of the Extremities, on Excision of the Head of the Thigh Bone, the Arm-Bone, and the Elbow-Joint. On the Diseases of the Peninsula, &c. 8vo.
- Lectures on the Operative Surgery of the Eye; or a Historical and Critical Inquiry into the methods recommended for the Cure of Cataract, for the Formation of Artificial Pupil, &c. &c. 7 plates. 8vo.
- Anatomy and Diseases of the Urinary and Sexual Organs: containing the Anatomy of the Bladder and of the Urethra, and the Treatment of the Obstructions to which these passages are liable. 8vo.
- GUY, W. A. Principles of Forensic Medicine. With Notes and Additions by Charles A. Lee, M.D. 8vo.
- HAHNEMANN, S. Organon of Homœopathic Medicine. 8vo.
- On Chronic Diseases, their specific nature and Homœopathic treatment. Edited by C. F. Hempel, M.D.
- HALE E., JR. History and Description of an Epidemic Fever, commonly called Spotted Fever, which prevailed at Gardiner, Maine, in 1814. 8vo.
- HALE, S. J. Flora's Interpreter; or the American Book of Flowers and Sentiments. 2 coloured plates. 12mo.
- HALFORD, SIR HENRY. Essays and Orations; with an Account of the opening of the Tomb of King Charles I. Contains Essays on the Climacteric Disease,—Necessity of Caution in estimating Symptoms in last Stages of some Diseases,—Tic Douloureux,—Popular and Classical Illustrations of Insanity,—Influence of some Diseases of the Body on the Mind,—Brain Fever of Aretæus,—Treatment of

- Gout,—Phlegmasia Dolens,—Moral Treatment of Insanity,—Deaths of Illustrious Persons of Antiquity, &c. 8vo.
- HALL, J. C. Interesting Facts connected with the Animal Kingdom; with some remarks on the Unity of our Species. 4 plates. 8vo.
- HALL, MARSHALL. The Principles of Diagnosis. With Notes by John A. Swett, M.D. 8vo.
- On the Mimoses; or Essay on Disorders of the Digestive Organs and General Health, and particularly of their numerous forms and complications, contrasted with some Acute and Insidious Diseases. 8vo.
- Commentaries, principally on those Diseases of Females which are Constitutional. 7 coloured plates. 8vo.
- A Critical and Experimental Essay on the Circulation of the Blood; also, Researches principally relative to the Morbid and Curative Effects of the Loss of Blood. 8vo.
- Lectures on the Nervous System and its Diseases. 8vo.
- Principles of the Theory and Practice of Medicine. Edited by J. Bigelow, M.D., and O. W. Holmes, M.D. Many engravings. 8vo.
- HALLER, A. VON. First Lines of Physiology. 8vo.
- Anatomical Description of the Arteries of the Human Body. Edited by John C. Warren, M.D. 15 coloured plates. 8vo.
- HALSTED, O. Account of the New Method of Curing Dyspepsia: with observations on Diseases of the Digestive Organs. 4 plates. 12mo.
- HAMILTON, J. Observations on the Utility and Administration of Purgative Medicines in several Diseases. 8vo.
- Observations on the use and abuse of Mercurial Medicines in various Diseases. With Notes by Ansel W. Ives, M.D. 8vo.
- HAMILTON, W. The History of Medicine, Surgery, and Anatomy, from the Creation of the World to the commencement of the 19th century. 2 vols. 8vo.
- HARE, ROBERT. A Compendium of the Course of Chemical Instruction in the Medical Department of the University of Pennsylvania. Numerous engravings. 8vo.
- HARRIS, C. A. The Principles and Practice of Dental Surgery. 69 engravings. 8vo.
- A Physiological and Pathological Inquiry concerning the Physical characteristics of the Human Teeth and Gums, the Salivary Calculus, the Lips and Tongue, and the Fluids of the Mouth, &c. 8vo.
- Dissertation on the Diseases of the Maxillary Sinus. 8vo.
- HARRISON, R. Surgical Anatomy of the Arteries of the Human Body.
- The Dublin Dissector, or Manual of Anatomy; comprising a description of the Bones, Muscles, Vessels, Nerves, and Viscera; also the Relative Anatomy of the different Regions of the Human Body, together with the Elements of Pathology. With Additions by Robert Watts, Jr., M.D. 12mo.
- HARRISON, J. An Essay towards a Correct Theory of the Nervous System. 8vo.
- HARRISON, J. P. Elements of Materia Medica and Therapeutics. 8vo.
- HASTINGS, C. An Experimental Inquiry of the general nature of Inflammation. 8vo.
- HASLAM, J. Illustrations of Madness; exhibiting a singular case of Insanity, and a no less remarkable difference in medical opinion; developing the nature of the assault, and the manner of working events; with a description of the tortures experienced by bomb-bursting, lobster-cracking, and lengthening the brain. Plate. 8vo.
- HATIN, J. A Manual of Practical Obstetrics; arranged so as to afford a concise and accurate description of the Management of Preternatural Labours; preceded by an account of the Mechanism of Natural Labour. Translated by S. D. Gross, M.D. With an Appendix containing a Physiological Memoir upon the Brain; from the French of M. Magendie, by J. Gardner, M.D. 12mo.
- HAWKINS, F. Rheumatism, and some Diseases of the Heart and other internal Organs: considered in the Gulstonian Lectures at the Royal College of Physicians, London. 8vo.
- HEBERDEN, W. Commentaries on the History and Cure of Diseases. 8vo.
- Epitome of the Diseases incident to Children. 12mo.

- HECKER, J. F. C. Epidemics of the Middle Ages: the Black Death in the 14th century, and the Dancing Mania in Germany and the Netherlands. Translated by B. G. Babington, M.D. 8vo.
- HENNEN, J. Principles of Military Surgery; comprising observations on the arrangement, police, and practice of Hospitals, and on the history, treatment, and anomalies of Variola and Syphilis. With a Life of the Author by his son. 8vo.
- HENRY, W. Elements of Experimental Chemistry. Many engravings. 2 vols. 8vo.
- HENRY, C. W. Report on the Physiology of the Nervous System. 8vo.
- HERSCHEL, SIR J. F. W. A Treatise on Astronomy. Many illustrations. 12mo.
- HEURTELOUP, BARON. Principles of Lithotomy: or a treatise on the art of extracting the Stone without incision. 5 plates. 8vo.
- HEUSTIS, J. W. Physical Observations, and Medical Tracts and Researches, on the Topography and Diseases of Louisiana. 8vo.
- HEWSON, W. An Experimental Inquiry into the Properties of the Blood: with remarks on some of its Morbid Appearances; and an Appendix relating to the Discovery of the Lymphatic System in Birds, Fish, and Amphibious Animals. 12mo.
- HEY, W. A Treatise on Puerperal Fever: illustrated by cases which occurred in Leeds and its vicinity, in 1809—1812. 8vo.
- HIBBARD, D. R. A Treatise on Cow Pox: in which the existence of Small Pox, or Varioloid in any form, subsequent to Vaccination, is shown to arise from some imperfection in its performance, and not the result of inefficacy on the part of the Vaccina to shield the system entirely from these diseases. 12mo.
- HILDENBRAND, J. V. DE. A Treatise on the Nature, Cause, and Treatment of Contagious Typhus. Translated by S. D. Gross, M.D. 8vo.
- HILLARY, W. Observations on the Changes of the Air, and the concomitant Epidemical Diseases in the Island of Barbadoes. With a Treatise on the Putrid Bilious Fever, commonly called the Yellow Fever; and such other Diseases as are indigenous or endemical in the West India Islands, or in the Torrid Zone. With Notes by Benjamin Rush, M.D. 8vo.
- HINDS, J. Veterinary Surgeon: or Farriery taught on a new and easy plan; being a Treatise on all the Diseases and Accidents to which the Horse is liable; the causes and symptoms of each, and the most approved remedies employed for the cure in every case, &c. With additions by Thomas M. Smith, V.S. Engravings. 12mo.
- The Groom's Oracle, and Pocket Stable Directory: in which the management of Horses generally, as to health, dieting, and exercise, are considered. 12mo.
- HIPPOCRATES. Aphorisms, from the Latin Version of Verhoofd, with a literal translation and explanatory Notes: by Elias Marks, M.D. 12mo.
- Prognostics and Crises. With critical and explanatory Notes, by H. W. Duchacet, M.D. 12mo.
- Hippocrates Contractus, in quo Magni Hippocratis, Medicorum Principis, Opera omnia in brevem epitomen summa diligentia redacta habentur. Studio et opera T. Burnet, M.D. 12mo.
- HITCHCOCK, EDWARD. Final Report on the Geology of Massachusetts 55 plates and 275 engravings. 2 vols. 4to.
- HOARE, C. A Practical Treatise on the Cultivation of the Grape Vine on open walls. 12mo.
- HOBLYN, R. E. A Dictionary of Terms used in Medicine and the Collateral Sciences. Edited by Isaac Hays, M.D. 12mo.
- HOCKEN, E. A. A Treatise on Amaurosis and Amaurotic Affections. 8vo.
- HODGKIN, T. Catalogue of the Preparations in the Anatomical Museum of Guy's Hospital. 8vo.
- HOLLAND, HENRY. Medical Notes and Reflections. 8vo.
- HOLMES, O. W. Boylston Prize Dissertations for years 1836 and 1837: containing, Facts and Traditions respecting the existence of Indigenous Intermittent Fever

- in New England, with a Map; Nature and Treatment of Neuralgia; and Utility and Importance of Direct Exploration in Medical Practice. Svo.
- HOME, E. On the Treatment of Ulcers on the Legs: with observations on Varicose Veins and Piles. 12mo.
- Practical Observations on the Treatment of Strictures in the Urethra, and in the Esophagus. 3 plates. Svo.
- HOME, F. Clinical Experiments, Histories, and Dissections. Svo.
- HOMIE Book of Health and Medicine: a Popular Treatise on the means of avoiding and curing Diseases, and of preserving the health and vigour of the Body to the latest period; including an account of the nature and properties of Remedies, the treatment of the Diseases of Women and Children, and the management of Pregnancy and Parturition. Svo.
- HOOD, S. Analytic Physiology, treating of the cure of Nervous Diseases, by external applications to the Spine. Svo.
- HOOPER, R. Lexicon Medicum; or Medical Dictionary; containing an explanation of the Terms in Anatomy, Botany, Chemistry, Materia Medica, Midwifery, Mineralogy, Pharmacy, Physiology, Practice of Physic, Surgery, and the various branches of Natural Philosophy, connected with Medicine. With Additions by Samuel Akerly, M.D. Svo.
- HOPE, J. A Treatise on Diseases of the Heart and Great Vessels, and on the Affections which may be mistaken for them. With Notes, &c., by C. W. Pennock, M.D. 9 plates. Svo.
- Principles and Illustrations of Pathological Anatomy; being a complete Series of Coloured Drawings, from originals, by the Author; with Descriptions, Symptoms, Treatment, &c. Edited by L. M. Lawson, M.D. 48 plates. Svo.
- HORNER, G. R. B. Medical and Topographical Observations upon the Mediterranean; and upon Portugal, Spain, and other countries. 8 plates. Svo.
- Medical Topography of Brazil and Uruguay, with illustrations. Svo.
- HORNER, W. E. Lessons in Practical Anatomy, for the use of Dissectors. Svo.
- Special Anatomy and Histology. 2 vols. Svo.
- HOSACK, DAVID. Lectures on the Theory and Practice of Physic, delivered in the College of Physicians and Surgeons, New York. Edited by Henry W. Ducachet, M.D. With Portrait. Svo.
- Inaugural Discourse, delivered at the opening of Rutgers Medical College, in the city of New York, 1826. Svo.
- Essays on Various Subjects of Medical Science. Plates. 3 vols. Svo.
- A System of Practical Nosology: to which is prefixed a Synopsis of the Systems of Sauvages, Linnæus, Vogel, Sagar, Macbride, Cullen, Darwin, Crichton, Pinel, Parr, Swediaur, Young, and Good: with references to the best authors on each Disease. Svo.
- HOUSTON, J. A. New York Lancet. Svo.
- HOVEY, C. M. Magazine of Horticulture, Botany, and all useful discoveries and improvements in Rural Affairs. Published monthly: 10 vols. now ready. Many engravings. Svo.
- HOWARD, J. Practical Observations on Cancer. Svo.
- HOWSHIP, J. Practical Remarks on the Discrimination and Appearances of Surgical Disease. Svo.
- HUFELAND, C. G. A Treatise on Scrofulous Disease. Translated by Charles D. Meigs, M.D. 12mo.
- HUFELAND, C. W. Enchiridion Medicum: or the Practice of Medicine; the result of 50 years' experience. Translated by C. Bruchhausen, M.S., and revised by R. Nelson, M.D. 12mo.
- HUGHES, H. M. Observations on Fibrinous Concretions in the Heart. Svo.
- HUNTER, JOHN. Observations on certain parts of the Animal Economy. With Notes by Richard Owen, F.R.S. Svo.

- HUNTER, JOHN. A Treatise on the Blood, Inflammation, and Gunshot Wounds. With Notes by James F. Palmer, M.D. 4 plates. 8vo.
- A Treatise on the Natural History and Diseases of the Human Teeth. With Notes by Thomas Bell, F.R.S. 5 plates. 8vo.
- Lectures on the Principles of Surgery. With Notes by James F. Palmer, M.D. 3 plates. 8vo.
- A Treatise on the Venereal Disease. With Notes by George G. Babington, M.D. 6 plates. 8vo.
- Complete Works. With an Account of his Life, by D. Ottley. 4 vols. Plates. 8vo.
- HUNTER, WILLIAM. The Anatomy of the Human Gravid Uterus exhibited in Figures. 34 plates, large folio.
- HUTIN, P. Manual of the Physiology of Man; or a concise description of the Phenomena of his Organization. With Notes by Joseph Tognio. 12mo.
- Illustrations of the Principal Regions of the Human Body, in relation to Surgical Anatomy. 12 plates, folio.
- Influenza or Epidemic Catarrh, Origin, Symptoms, and Cure of,—also Hints respecting Common Colds, and Incipient Pulmonary Consumption. 8vo.
- INNES, J. A Short Description of the Human Muscles, arranged as they appear on Dissection; also their several Uses, and the Synonyma of the best authors. 17 plates. 12mo.
- JACKSON, R. A Treatise on the Fevers of Jamaica; with some observations on the Intermitting Fever of America, &c. 12mo.
- JACKSON, S. Principles of Medicine, founded on the Structure and Functions of the Animal Organism. 8vo.
- JAHR, G. H. G. New Homœopathic Pharmacopœia and Posology; or the Preparation of Homœopathic Medicines and the Administration of Doses. With Additions by F. Kitchen, M.D. 8vo.
- Manual of Homœopathic Medicine. With Additions by C. Hering, M.D.
- JAMES, B. A Treatise on the Management of the Teeth. 12mo.
- JARDINE, SIR WILLIAM. The Naturalist's Library. Complete in 40 vols. 12mo., viz.
- | | | | |
|--------------|----------|--------------|---------|
| Mammalia. | 13 vols. | Ichthyology. | 6 vols. |
| Ornithology. | 14 vols. | Entomology. | 7 vols. |
- Illustrated by upwards of 1300 elegant coloured plates.
- "Perhaps the most interesting, the most beautiful, and the cheapest series yet offered to the public."—*Athenæum*.
- JEFFRAY, J. Observations on the Heart, and on the Peculiarities of the Fœtus. 15 plates. 8vo.
- JENNER, E. Inquiry into the Causes and Effects of the Variolæ Vaccinæ, a Disease discovered in some of the western counties of England, and known by the name of Cow-Pox. 4 coloured plates. 4to.
- The Life of Edward Jenner, M.D., with illustrations of his doctrines, and selections from his correspondence, by John Baron, M.D. Portraits. 2 vols. 8vo.
- JOBSON, D. W. A Treatise on the Anatomy and Physiology of the Teeth; their Diseases and Treatment, &c. 4 plates. 8vo.
- JOHNSON, C. W. Farmer's Encyclopedia, and Dictionary of Rural Affairs; embracing all the most recent Discoveries in Agricultural Chemistry. Adapted to the United States, by G. Emerson. Many illustrations of Animals, Implements, &c. 8vo.
- JOHNSON, C. T. A Practical Essay on Cancer. 12mo.
- JOHNSON, E. The Theory, Principles, and Practice of the Water Cure shown to be in accordance with Medical Science, and the Teachings of Common Sense. 9 engravings. 12mo.
- JOHNSON, JAMES. Medico-Chirurgical Review, and Journal of Practical Medicine. Published Quarterly. 50 vols. now complete. See Notice at the end of Catalogue.
- Influence of Civic Life, Sedentary Habits, and Intellectual Refinement on Human Health, and Human Happiness; including an Estimate of the balance of enjoyment and suffering in the different gradations of Society. 8vo.

- JOHNSON, JAMES. Essay on Indigestion ; or Morbid Sensibility of the Stomach and Bowels, as the proximate cause or characteristic condition of Dyspepsy, Nervous Irritability, Mental Despondency, Hypochondriasis, and many other ailments of Body and Mind. With Observations on the Diseases and Regimen of Invalids on their return from hot and unhealthy climates. 12mo.
- Change of Air, or the Philosophy of Travelling ; being Autumnal Excursions through France, Switzerland, Italy, Germany, and Belgium ; with observations and reflections on the Moral, Physical, and Medicinal Influence of Travelling-Exercise, Change of Scene, Foreign Skies, and Voluntary Expatriation. To which is prefixed, Wear and Tear of Modern Babylon. 8vo.
- Treatise on Derangements of the Liver, Internal Organs, and Nervous System. 12mo.
- Economy of Health ; or the Stream of Human Life from the Cradle to the Grave. With Reflections, Moral, Physical, and Philosophical, on the Septennial Phases of Human Existence. 18mo.
- JOHNSON, JAMES, AND J. R. MARTIN. The Influence of Tropical Climates on European Constitutions. 8vo.
- JOHNSTON, H. Practical Observations on Urinary Gravel and Stone ; on Disease of the Bladder and Prostate Gland ; and on Strictures of the Urethra. 12mo.
- JOHNSTON, J. F. W. Elements of Agricultural Chemistry and Geology. 18mo.
- Lectures on the Application of Chemistry and Geology to Agriculture. 12mo.
- JOHNSTONE, J. A Therapeutic Arrangement and Syllabus of Materia Medica. 8vo.
- JONES, J. F. D. A Treatise on the Process employed by Nature in Suppressing Hæmorrhage from divided and punctured Arteries ; on the Use of the Ligature ; and on Secondary Hæmorrhages. 15 plates. 8vo.
- JONES, T. W. Manual of Pharmacology ; or Compendium of Materia Medica, Conspicuous of the London, Edinburgh, and Dublin Pharmacopœias, and Formulary of Extemporaneous Prescriptions. 18mo.
- KANE, R. Elements of Chemistry, including the most recent discoveries and applications of the Science to Medicine and Pharmacy, and to the Arts. With Additions, &c., by John William Draper, M.D. Numerous engravings. 8vo.
- KAY, J. P. Physiology, Pathology, and Treatment of Asphyxia : including Suspended Animation in new-born Children, and from Drowning, Hanging, Wounds of the Chest, Mechanical Obstructions of the Air-Passages, Respiration of Gases, Death from Cold, &c. Plate. 8vo.
- KENNEDY, E. Observations on Obstetric Auscultation, with an Analysis of the Evidences of Pregnancy, and an Inquiry into the proofs of the life and death of the Fœtus in Utero. With Notes by J. E. Taylor, M.D. 17 plates. 12mo.
- KENRICK, W. New American Orchardist ; or an Account of the most valuable Fruit of all climates, adapted to cultivation in the United States ; with their History, modes of Culture, &c. 12mo.
- KEY, C. A. Report of Primary Syphilitic Cases. 8vo.
- KIRBY, J. Cases with Observations on Wry Neck ; on the Reduction of Luxations of the Shoulder Joint ; on the Operation for Hare Lip ; on Cartilaginous substances of the Knee Joint ; on Aneurism ; on the Use of Extract of Stramonium, &c. 8vo.
- KIRBY, W. On the Power, Wisdom, and Goodness of God, as manifested in the Creation of Animals, and in their History, Habits, and Instincts. A Bridgewater Treatise. 20 plates. 8vo.
- KIRTLAND, G. The Anatomy of the Horse's Foot. 11 coloured figures.
- KITCHENER, W. Economy of the Eyes. Plain Rules which will enable all to judge exactly when and what Spectacles are best calculated for their Eyes, and Precepts for the improvement and preservation of the Sight. 12mo.
- Directions for Invigorating and Prolonging Life ; or, the Invalid's Oracle With Improvements by T. S. Barrett, M.D. 18mo.
- KNOX, G. F. The Art of Cupping ; being a brief History of the Operation, from its origin to the present time ; its utility ; rules for its performance ; a list of dis-

eases in which it is most beneficial; description of the various instruments employed, &c. 8vo.

KNOX, J. F. *The Anatomist's Instructor, and Museum Companion*; being practical directions for the formation and subsequent management of Anatomical Museums. Plate. 12mo.

KOECKER, L. *Essays on Diseases of the Jaws, and their Treatment*; with observations on the Amputation of a part or the whole of the Inferior Maxilla; tending to prove such operation is seldom, if ever, necessary. 2 plates. 8vo.

——— *An Essay on Artificial Teeth, Obturators, and Palates, with the Principles for their Construction and Application.* 21 plates. 8vo.

KRAMER, W. *Nature and Treatment of Diseases of the Ear.* Translated by J. R. Bennett, M.D. Engravings. 8vo.

Lady's Medical Pocket-Book, and Nursery Adviser. 32mo.

LAENNEC, R. T. H. *A Treatise on Diseases of the Chest, and on Mediate Auscultation.* With copious Notes, a Sketch of the Author's Life, and an extensive Bibliography of the different Diseases, by John Forbes, M.D. To which are added the Notes of Professor Andral, translated and accompanied with Observations on Cerebral Auscultation, by John D. Fisher, M.D. 2 plates. 8vo.

LAENNEC, M. *Manual of Percussion and Auscultation.* Translated by J. B. Sharpe, M.R.C.S. 18mo.

LALLEMAND, M. *On Involuntary Seminal Discharges*: translated by W. Wood, M.D. 8vo.

LARREY, BARON D. J. *Surgical Memoirs of the Campaigns of Russia, Germany, and France.* Translated by John C. Mercer. 6 plates. 8vo.

——— *Observations on Wounds, and their complications by Erysipelas, Gangrene, and Tetanus, and on the principal Diseases and Injuries of the Head, Ear, and Eye.* Translated by E. F. Rivinus, M.D. 2 plates. 8vo.

LATHAM, P. M. *Lectures on subjects connected with Clinical Medicine.* 8vo.

LAURIE, J. *Homœopathic Domestic Medicine, with the Treatment and Diseases of Females, Infants, Children, and Adults.* With Additions by A. G. Hull, M.D.

LAVATER, J. C. *Essays on Physiognomy*; designed to promote the Knowledge and the Love of Mankind. With Memoirs of the Life of the Author, and upwards of 400 profiles and other engravings. 8vo.

LAWRENCE, W. *A Treatise on Diseases of the Eye.* With Numerous Additions, and 67 illustrations, by Isaac Hays, M.D. 8vo.

——— *Lectures on Physiology, Zoology, and the Natural History of Man.* 7 plates. 8vo.

——— *A Treatise on the Venereal Diseases of the Eye.* 8vo.

——— *A Treatise on Ruptures.* 8vo.

LAYCOCK, T. *An Essay on Hysteria*; being an Analysis of its irregular and aggravated forms, including Hysterical Hæmorrhage, and Hysterical Ischuria. 8vo.

LEE, A. *Chemical Diagrams, accompanied with a concise description of each Decomposition, the Vegetable Alkalies, the Urine and Urinary Calculi, Tables of Chemical Equivalents, &c.* 12mo.

LEE, E. *Observations on the Principal Medical Institutions and Practice of France, Italy, and Germany*; with Notices of the Universities, and Cases from Hospital Practice. With an Appendix on Animal Magnetism and Homœopathy. 8vo.

LEE, R. *Lectures on the Theory and Practice of Midwifery.* 67 engravings. 8vo.

LE GALLOIS, M. *Experiments on the Principle of Life, and particularly on the Principle of the Motions of the Heart, and on the Seat of this Principle, &c.* Translated by N. C. and J. G. Nancrede, M.D. Plate. 8vo.

LEO WOLF, W. *Remarks on the Abracadabra of the Nineteenth Century*; or, on Dr. S. Hahnemann's Homœopathic Medicine, with particular reference to Dr. C. Hering's "Concise View of the Rise and Progress of Homœopathic Medicine." 8vo.

Lessons on Shells. 10 plates. 18mo.

Letters to a Mother on the Watchful Care, of her Infant. 18mo.

- LEVER, J. C. W. A Practical Treatise on Organic Diseases of the Uterus. 8vo.
- LIEBIG, J. Familiar Letters on Chemistry, and its relation to Commerce, Physiology, and Agriculture. Edited by J. Gardner, M.D. 18mo.
- Animal Chemistry, or Organic Chemistry in its application to Physiology and Pathology. With Additions, &c., by W. Gregory, M.D., and J. W. Webster, M.D. 12mo.
- LIEUTAUD, J. Synopsis of the Universal Practice of Medicine; exhibiting a concise view of all Diseases, both internal and external, illustrated with complete commentaries. Translated by E. A. Atlee, M.D. 8vo.
- LINCOLN, ALMIRA H. Familiar Lectures on Botany, Practical, Elementary, and Physiological. With an Appendix containing Descriptions of the Plants of the United States, and Exotics, &c. Many engravings. 12mo.
- LIND, J. Essay on Diseases incidental to Europeans in Hot Climates; with the method of preventing their fatal consequences. With an Appendix on Intermittent Fevers, &c. 8vo.
- LINDLEY, J. Theory of Horticulture: or an Attempt to explain the principal operations of Gardening upon Physiological Principles. With Notes by A. J. Downing and A. Gray. 35 engravings. 12mo.
- An Introduction to the Natural System of Botany; or a Systematic View of the organization, natural affinities, and geographical distribution of the whole Vegetable Kingdom; with the Uses of the most important species in Medicine, the Arts, and Rural or Domestic Economy. With an Appendix by John Torrey, M.D. 8vo.
- LISFRANC, M. Diseases of the Uterus, a Series of Clinical Lectures delivered at the Hospital La Pitié. Edited by H. Pauly, M.D., and translated by G. H. Lodge, M.D. 8vo.
- LISTON, R. Elements of Surgery. With Notes and Additions by S. D. Gross, M.D. Numerous engravings. 8vo.
- Practical Surgery. With Notes, &c., by G. W. Norris, M.D. 150 engravings. 8vo.
- Lectures on the Operations of Surgery, and on Diseases and Accidents requiring Operations. With Additions by T. D. Mutter, M.D. Numerous illustrations. 8vo.
- Literary Men. Remarks on the Disorders of Literary Men, or an inquiry into the means of preventing the evils usually incident to sedentary and studious habits. 12mo.
- LITTELL, JR., S. A Manual of Diseases of the Eye. 12mo.
- LITTLE, W. J. Ankylosis, or Stiff Joint: a Practical Treatise on the Contractions and Deformities resulting from Diseases of Joints. 32 engravings. 8vo.
- LIZARS, J. A System of Anatomical Plates of the Human Body; accompanied with Descriptions, and Physiological, Pathological, and Surgical Observations. 101 beautifully coloured plates, folio.
- LOEBSTEIN, J. F. D. A Treatise upon the Semeiology of the Eye, for the Use of Physicians; and of the Countenance, for Criminal Jurisprudence. 8vo.
- A Treatise on the Structure, Functions, and Diseases of the Human Sympathetic Nerve. Translated by J. Pancoast, M.D. 6 plates. 8vo.
- London Dissector, or Guide to Anatomy, for the use of Students: comprising a Description of the Muscles, Vessels, Nerves, Lymphatics, and Viscera of the Human Body, as they appear on Dissection; with Directions for their Demonstration. Revised by E. J. Chaisty, M.D.
- London Practice of Midwifery; including the Treatment during the Puerperal state, and the principal Infantile Diseases; with engravings, and Aphorisms of Dr. Denman. 8va.
- London Surgical Pocket-Book; (Medical, Operative, and Mechanical;) founded on the popular Lectures and Works of Abernethy, Sir A. Cooper, Lawrence, &c.; subdigested in the order of Causes, Symptoms, Chirurgical and Medical Treatment; Diagnoses, Prognoses, Modes of Operation, and other agents employed in Hospital and private Practice, &c. &c. 2 plates. 12mo.

- LOVDON, J. C. *Encyclopedia of Agriculture*: containing the Theory and Practice of the Valuation, Transfer, Laying out, Improvement, and Management of Landed Property; and the Cultivation and Economy of the Animal and Vegetable Productions of Agriculture, including all the latest improvements; a General History of Agriculture in all countries, &c. &c. 1272 engravings. 8vo.
- *Encyclopedia of Gardening*; comprising the Theory and Practice of Horticulture, Floriculture, Arboriculture, and Landscape Gardening, including all the latest improvements; a General History of Gardening in all countries, &c. &c. 981 engravings. 8vo.
- LOVDON, Mrs. *Gardening for Ladies, and Companion to the Flower Garden*; being an Alphabetical arrangement of all the Ornamental Plants usually grown in Gardens and Shrubberies; with full directions for their Culture. Edited by A J Downing. 61 engravings. 12mo.
- LOUIS, P. C. A. *Researches on the effects of Blood-letting in some Inflammatory Diseases, and on the Influence of Tartarized Antimony, and Vesication in Pneumonitis*. Translated by C. G. Putnam, M.D. With an Appendix, &c., by James Jackson, M.D. 8vo.
- *Anatomical, Pathological, and Therapeutic Researches upon the Disease known under the name of Gastro-Enterite, Putrid, Adynamic, Ataxic, or Typhoid Fever, &c., compared with the most common Acute Diseases*. Translated by Henry I. Bowditch, M.D. 2 vols. 8vo.
- *Pathological Researches on Phthisis*. With Notes, Additions, &c., by Charles Cowan, M.D., and revised by H. I. Bowditch, M.D. 8vo.
- *Anatomical, Pathological, and Therapeutic Researches on the Yellow Fever of Gibraltar, of 1828*. Translated by G. C. Shattock, Jr., M.D. 8vo.
- LOWRIE, W. F. *Toxicologia, or a Treatise on Internal Poisons, in their relation to Medical Jurisprudence, Physiology, and the Practice of Physic*. 8vo.
- LUCAS, P. B. A *Concise Anatomical Description of the Arteries of the Human Body*; with full directions for cutting down upon and securing the several Arterial Trunks. 12mo.
- LUDLOW, J. L. A *Mannual of Examinations upon Anatomy and Physiology, Surgery, Practice of Medicine, Chemistry, Materia Medica, Obstetrics, &c.* 12mo.
- LUGOL, J. G. A. *Researches on Scrofulous Diseases*. Translated by A. S. Doane, M.D. 12mo.
- LYELL, C. *Elements of Geology*. 294 illustrations. 12mo.
- *Principles of Geology*: being an Inquiry how far the former changes of the Earth's surface are referable to causes now in operation. 241 illustrations. 2 v. 8vo.
- MACARTNEY, J. A *Treatise on Inflammation*. 8vo.
- MACAULAY, A. A *Dictionary of Medicine, designed for Popular Use*: containing an Account of Diseases and their Treatment, including those most frequent in Warm Climates; with directions for administering the various substances used as Medicines; the Regulation of Regimen and Diet; and the management of Diseases of Women and Children. 8vo.
- MACCULLOCH, J. An *Essay on the Remittent and Intermittent Diseases, including, generically, Marsh Fever and Neuralgia*: comprising, under the former, various anomalies, obscurities, and consequences, and under a new systematic view of the latter, treating of Tic Douloureux, Sciatica, Headache, Ophthalmia, Toothache, Palsy, &c. &c. 8vo.
- *Malaria; an Essay on the Production and Propagation of this Poison, and on the nature and localities of the places by which it is produced*. 8vo.
- M'DOWELL, W. A *Demonstration of the Curability of Pulmonary Consumption, in all its stages*: comprising an Inquiry into the nature, causes, symptoms, treatment, and prevention of Tuberculous Diseases in general. 8vo.
- M'INTOSH, C. *The Orchard*; including the management of Wall and Standard Fruit Trees, and the Forcing Pit; with selected lists and synonyms of the choicest varieties. 18 coloured plates, and many wood engravings. 12mo.

- MACKENZIE, W. A Practical Treatise on the Diseases of the Eye. 8vo.
- MACKENZIE'S Five thousand Receipts in all the Useful and Domestic Arts. 8vo
- MACKINTOSH, J. Principles of Pathology and Practice of Medicine. With Notes and Additions by S. G. Morton, M.D. 8vo.
- MACLEAN, C. Evils of Quarantine Laws, and Non-existence of Pestilential Contagion; deduced from the phenomena of the Plague of the Levant, the Yellow Fever of Spain, and the Cholera Morbus of Asia. 8vo.
- MACLEAN, L. Inquiry into the Nature, Causes, and Cure of Hydrothorax. 8vo.
- MACNISH, R. The Philosophy of Sleep. 12mo.
- MACROBIN, J. An Introduction to the Study of Practical Medicine; being an Outline of the leading facts and principles of the Science. 8vo.
- MAGENDIE, F. An Elementary Treatise on Human Physiology. Translated and enlarged by John Revere, M.D. 61 engravings. 8vo.
- Lectures on the Blood; and on the changes which it undergoes during Disease. 8vo.
- Physiological and Chemical Researches on the Use of Prussic or Hydrocyanic Acid in the treatment of Diseases of the Breast, and particularly in Phthisis Pulmonalis. With Notes by J. G. Percival, M.D. 12mo.
- Formulary for the Preparation and Employment of several New Remedies. Edited by J. Houlton, F.L.S. 12mo.
- MALCOLMSON, J. G. Clinical Remarks on some Cases of Liver Abscess presenting externally. 8vo.
- MALE, G. E. Elements of Juridical or Forensic Medicine; for the use of Medical Men, Coroners, and Barristers. 8vo.
- MANEC, J. P. Great Sympathetic Nerve: a coloured plate, 30 inches long, on Roller.
- Cerebro-Spinal System in Man, together with the Origin and Primary Divisions of the Nerves which arise from it. 32 inches long, on Roller.
- MANN, J. Medical Sketches of the Campaigns of 1812, 1813, 1814: with Surgical Cases; Observations on Hospitals; a Dissertation on Dysentery; Observations on Peripneumonia Notha, &c. 8vo.
- MARKHAM, W. O. Remarks on the Surgical Practice of Paris. 8vo.
- MARSHAL, A. Morbid Anatomy of the Brain in Mania and Hydrophobia; with the Pathology of these two Diseases. Edited by S. Sawrey.
- MARSHALL, J. Practical Observations on Diseases of the Heart, Lungs, Stomach, Liver, &c., occasioned by Spinal Irritation: and on the Nervous System in general, as a source of Organic Disease. 8vo.
- MARTINET, L. Manual of Therapeutics. With Additions, by R. Norton, M.D. 18mo.
- Manual of Pathology; containing the Symptoms, Diagnosis, and Morbid Characters of Diseases, &c. With Notes by Jones Quain, M.D. 12mo.
- MASON, R. The Gentleman's New Pocket Farrier, comprising a general Description of the Horse; with the quickest and simplest mode of fattening; necessary treatment whilst undergoing excessive fatigue, or on a journey; the construction and management of Stables; different marks for ascertaining the Age of a Horse; Diseases to which he is subject, with Remedies, &c., &c. Also a Prize Essay on Mules; and Observations and Recipes for the cure of the most common distempers incident to Oxen, Cows, Calves, Sheep, Lambs, Swine, Dogs, &c. Engravings. 12mo.
- MASSE, J. N. Petit Atlas complet d'Anatomie descriptive du Corps Humain, destiné à compléter tous les Traités d'Anatomie descriptive. 112 exquisitely coloured plates. 12mo.
- MAUNSELL, H. The Dublin Practice of Midwifery. With Notes and Additions by C. R. Gilman, M.D. 12mo.
- MAURY, F. Treatise on the Dental Art. With Notes and Additions by J. B. Savier 20 plates and 54 wood engravings. 8vo.

- MAYGRIER, J. P. Anatomist's Manual; or a Treatise on the manner of preparing all the parts of Anatomy, with a complete description of the parts. Translated by G. S. Bedford, M.D. 12mo.
- Midwifery Illustrated. With Notes, &c., by A. S. Doane, M.D. 84 plates. 8vo.
- MAYO, H. Management of the Organs of Digestion, in Health and in Disease. 8vo
- Outlines of Human Pathology. 8vo.
- MAYO, T. Elements of the Pathology of the Human Mind. 8vo.
- MEASE, J. On the Causes, Cure, and Prevention of the Sick-Headache. 8vo.
- MECKEL, J. F. Manual of General, Descriptive, and Pathological Anatomy: with Additions and Notes by A. J. L. Jourdan and G. Breschet, M.D. Translated by A. S. Doane, M.D. 3 vols. 8vo.
- Medical Magazine, by Drs. Peirson, Flint, and Bartlett—Boston. Plates. 3 vols. 5vo
- Medical and Surgical Register: consisting chiefly of Cases in the New York Hospital. By John Watts, Jr., M.D., Valentine Mott, M.D., and Alexander H. Stevens, M.D. Plates. 2 vols. 8vo.
- Medicina Flagellata; or, the Doctor Scarified: laying open the vices of the Faculty; the Insignificance of a great part of their Materia Medica; with certain rules to discern the true Physician from the Empiric, and the Useful Medicine from the Noxious and Trading Physic, &c., &c. 12mo. 1721.
- Medico-Chirurgical Transactions: a collection of the most valuable Memoirs read to the Medico-Chirurgical Societies of London and Edinburgh, &c., &c. Edited by Isaac Hays, M.D. 8vo.
- MEIGS, C. D. The Philadelphia Practice of Midwifery. 36 engravings. 8vo.
- MENDENHALL, G. Medical Student's Vade Mecum; containing Examinations upon Anatomy, Chemistry, Materia Medica, Surgery, Practice of Medicine, Obstetrics, and Poisons. 18mo.
- MERRIAM, S. Dissertation on Retroversion of the Womb, including observations on Extra Uterine Gestation. 7 plates. 8vo.
- Synopsis of the various kinds of Difficult Parturition, with Practical remarks on the Management of Labours. With Notes by T. C. James, M.D. 5 plates. 8vo.
- MICHELL, W. On Difficult Cases of Parturition; and the Use of Ergot of Rye. 8vo.
- MIDDLEMORE, R. Treatise on Diseases of the Eye and its Appendages. 2 vols. 8vo.
- MILLER, E. Medical Works of Edward Miller, M.D., late Professor of the Practice of Physic in the University of New York: with a Biographical Sketch of the author, by Samuel Miller, D.D. 8vo.
- MILLER, J. The Principles of Surgery, by James Miller, F.R.S.E., Professor of Surgery in the University of Edinburgh. 8vo.
- MILLENGEN, J. G. Aphorisms on the Treatment and Management of the Insane; with considerations on Public and Private Lunatic Asylums, pointing out the errors in the present system. 8vo.
- Curiosities of Medical Experience. 8vo.
- MINER, T., AND W. TULLY. Essays on Fevers, and other Medical subjects. 8vo.
- MONDAT, C. V. On Sterility in the Male and Female, its Causes and Treatment. 50 illustrations. 12mo.
- MONEY, W. A Vade Mecum of Morbid Anatomy, Medical and Chirurgical; with Pathological Observations and Symptoms. Illustrated by upwards of 250 drawings. 8vo.
- MONRO, A. Anatomy of the Human Bones and Nerves, with a description of the Human Lacteal Sac and Duct. With Notes, by J. Kirby, M.D. 12mo.
- MONTGOMERY, W. F. Exposition of the Signs and Symptoms of Pregnancy, the Period of Human Gestation, and the Signs of Delivery. 8vo.
- MORAND, M. Memoir on Acupuncture, embracing a series of cases, drawn up under the inspection of M. Julius Cloquet: translated by F. Bache, M.D. 12mo.
- MOREAU, F. J. Practical Treatise on Midwifery; exhibiting the present advanced state of the Science. Edited by P. B. Goddard, M.D. 80 plates. 4to.

- MORGAGNI, J. B. Seats and Causes of Diseases, investigated by Anatomy; containing a great variety of Dissections, and accompanied with Remarks. With Notes, &c., by W. Cooke, M.D. 2 vols. 8vo.
- MORGAN, G. T. Outline of Inflammation and its Effects; being First Principles of Surgery. 8vo.
- MORISON, A. Outlines of Lectures on Mental Diseases. 13 plates. 8vo.
- MORTON, S. G. Illustrations of Pulmonary Consumption, its Anatomical Characters, Causes, Symptoms, and Treatment. 12 plates, drawn and coloured from Nature. 8vo.
- *Crania Americana*; or a Comparative View of the Skulls of various Aboriginal Nations of North and South America; to which is prefixed an Essay on Varieties of the Human Species. 78 Plates. Folio.
- MORTON, T. Engravings illustrating the Surgical Anatomy of the Head and Neck, Axilla Bend of the Elbow and Wrist, with Descriptions. 8 coloured plates. 8vo.
- The Surgical Anatomy of the Groin, the Femoral and Popliteal Regions. 8 coloured plates and 11 wood engravings. 8vo.
- The Surgical Anatomy of Inguinal Herniæ, the Testis and its Coverings. 5 coloured plates and 11 wood engravings. 8vo.
- The Surgical Anatomy of the Perineum. 4 coloured plates and 3 wood engravings. 8vo.
- MOSELEY, B. Treatise on Tropical Diseases, and on the climate of the West Indies. 8vo.
- MOSS, W. Essay on the Management and Feeding of Infants. 12mo.
- MOTT, VALENTINE. Travels in Europe and the East. 8vo.
- MUEHRY, A. Observations on the Comparative State of Medicine in France, England, and Germany. Translated by E. G. Davis, M.D. 8vo.
- MULLER, J. Elements of Physiology. Translated by Wm. Baly, M.D. Edited by John Bell, M.D. 8vo.
- MUNRO, H. Compendious System of the Theory and Practice of Modern Surgery, arranged in a New Nosological and Systematic Method, different from any yet attempted in Surgery,—in the form of a Dialogue. 8vo.
- MURPHY, P. J. Practical Observations, showing that Mercury is the sole cause of what are termed Secondary Symptoms. 8vo.
- MURRAY, A. Description of the Arteries of the Human Body; reduced to Tables. Translated by A. Scott. 8vo.
- MURRAY, J. System of Materia Medica and Pharmacy. With Notes by N. Chapman, M.D. 2 vols. 8vo.
- MUTER, R. Practical Observations on the Lateral Operation of Lithotomy, and on the various improved and new modes of performing this operation; with remarks on the Recto Vesical Operation. 2 plates. 8vo.
- MUTTER, T. D. Lecture on Loxarthrus, or Club Foot. Engravings. 8vo.
- Naturalist. Manual of the Practical Naturalist; or Directions for Collecting, Preparing, and Preserving Subjects of Natural History. Plate. 12mo.
- NEILL, J. Outlines of the Arteries, with short Descriptions, for the use of Medical Students. 6 coloured plates.
- Outlines of the Nerves, with short Descriptions, for the use of Medical Students. Coloured plates.
- NELIGAN, J. M. Medicines, their Uses and Mode of Administration; including a complete Conspectus of the Three British Pharmacopœias, an Account of all the New Remedies, and an Appendix of Formulæ. With Notes, &c., by D. M. Reese, M.D. 8vo.
- Nervous System. Documents and Dates of Modern Discoveries in the Nervous System, by Walker, Bell, Magendie, Whytt, Prochaska, Hall, Muller, Copland, &c. 8vo.
- NEVILLE, W. B. On Insanity; its Nature, Causes, and Cure. 8vo.
- NEWNHAM, W. Human Magnetism; its Claims to Dispassionate Inquiry: being an attempt to show the utility of its application for the Relief of Human Suffering. 12mo.

- NICHOLLES, J. The Teeth in relation to Beauty, Voice, and Health. 8vo.
- NICOL, W. Villa-Garden Directory, or Monthly Index of Work to be done in Town and Villa Gardens, Shrubberies, and Parterres, &c. 12mo.
- NORTH, E. Outlines of the Science of Life; which treats Physiologically of both Body and Mind; designed only for Philosophers and other candid persons. With Essays on other subjects. Plate. 8vo.
- A Treatise on a Malignant Epidemic, commonly called Spotted Fever; with remarks on the nature of Fever in general, &c. 12mo.
- NORTH, J. Practical Observations on the Convulsions of Infants. 8vo.
- NORTON, W. A. An Elementary Treatise on Astronomy; containing a systematic and comprehensive exposition of the Theory, and the more important Practical Problems, with Solar, Lunar, and other Astronomical Tables. 132 illustrations. 8vo.
- NORWOOD, J. G. Treatise on the Prevention and Cure of Diseases, by Regimen and Simple Medicines. With the addition of a Vegetable Materia Medica, pointing out the virtues, preparations, and doses, of our most valuable native Medicinal Plants; and an Outline of Anatomy and Physiology. 100 engravings. 8vo.
- NUNNELEY, T. Treatise on the Nature, Causes, and Treatment of Erysipelas. 8vo.
- OLIVER, D. First Lines of Physiology. 8vo.
- OLMSTED, D. Introduction to Astronomy: with a Supplement by E. P. Mason, containing Rules for the adjustment and use of Astronomical Instruments, Calculation of Eclipses and Occultations, and methods of finding the Latitude and Longitude. 8 plates and 74 engravings. 8vo.
- Compendium of Astronomy, containing the Elements of the Science, familiarly explained and illustrated, with the latest discoveries. 54 engravings. 12mo.
- Introduction to Natural Philosophy. 305 illustrations. 8vo.
- Compendium of Natural Philosophy; adapted to the use of the general reader. With a Supplement containing instructions to Young Experimenters, with a copious list of experiments and directions for performing them. 207 engravings. 12mo.
- Rudiments of Natural Philosophy and Astronomy, for the Younger Classes in Academies, &c. 122 engravings. 18mo.
- Oracle of Health and Long Life: containing plain and practical instructions for the preservation of Sound Health, and the attainment of vigorous Old Age, &c., by Medicus. 12mo.
- ORFILA, M. P. Popular Treatise on the Remedies to be employed in cases of Poisoning and Apparent Death; including the means of detecting Poisons, of distinguishing Real from Apparent Death, and on ascertaining the Adulteration of Wines. Translated by W. Price, M.D. 8vo.
- OSBORNE, J. Nature and Treatment of Dropsical Diseases. 8vo.
- PAINE, MARTYN. Medical and Physiological Commentaries. 3 vols. 8vo.
- A Therapeutical Arrangement of the Materia Medica; or the Materia Medica arranged on Physiological principles. 12mo.
- Letters on the Cholera Asphyxia, as it has appeared in the City of New York. 8vo.
- PAMBOUR, CHEV. F. M. G. DE. Practical Treatise on Locomotive Engines upon Railways. 4 plates. 8vo.
- PANCOAST, J. Treatise on Operative Surgery: comprising a description of the various Processes of the Art, including all the New Operations; exhibiting the State of Surgical Science in its present advanced condition. 80 plates. 4to.
- PARIS, J. A. Pharmacologia; being an extended Inquiry into the operations of Medicinal Bodies, upon which are founded the Theory and Art of Prescribing. With Notes by Charles A. Lee, M.D. 8vo.
- Same work, with Notes and Additions by John B. Beck, M.D. Plate. 8vo.
- Treatise on Diet; with a view to establish, on practical grounds, a system of Rules for the Prevention and Cure of the Diseases incident to a disordered state of the Digestive Functions. 8vo.

- PARIS, J. A. Elements of Medical Chemistry ; embracing only those branches of Chemical Science which are calculated to illustrate or explain the different objects of Medicine ; and to furnish a Chemical Grammar to the Author's Pharmacologia. Numerous illustrations. 8vo.
- PARIS, J. A., AND J. S. M. FONBLANQUE. Medical Jurisprudence. 3 vols. 8vo.
- PARK, R. Pantology ; or a Systematic Survey of Human Knowledge ; proposing a classification of all its branches, and illustrating their history, relations, uses, and objects ; with a synopsis of their leading facts and principles ; and a select catalogue of Books on all subjects, suitable for a Cabinet Library. 12 plates. 8vo.
- PARKER, L. The Stomach in its Morbid States, and the Modern Treatment of Syphilitic Diseases. 8vo.
- PARMLEY, L. S. A Practical Guide to the Management of the Teeth ; comprising a discovery of the origin of Caries, or Decay of the Teeth ; with its prevention and cure. Plate. 18mo.
- PARNELL, E. A. Applied Chemistry ; in Manufactures, Arts, and Domestic Economy. 45 illustrations. 8vo.
- PARRISH, J. Practical Observations on Strangulated Hernia, and some of the Diseases of the Urinary Organs. 4 plates. 8vo.
- PARSONS, H. Directions for Making Anatomical Preparations, formed on the Basis of Pole, Marjolin, and Breschet, and including the new mode of Swan. 4 plates. 8vo.
- Boylston Prize Dissertations on Inflammation of the Periosteum, Eneuresis Irritata, Cutaneous Diseases, Cancer of the Breast ; also remarks on Animal and Vegetable Decomposition. 8vo.
- PASCALIS, F. Exposition of the Dangers of Interment in Cities ; illustrated by an account of the Funeral Rites and Customs of the Hebrews, Greeks, Romans, and Primitive Christians ; by ancient and modern Ecclesiastical Canons, Civil Statutes, and Municipal Regulations ; and by Chemical and Physical Principles. 8vo.
- PAUL, A. Essay on Ringworm, cured by a Medicated Soap, without the use of Mercury : with Hints on the subject to Parents, &c. 2 coloured plates. 8vo.
- PAXTON, J. Introduction to the Study of Human Anatomy. With Additions by W. Lewis, Jr., M.D. 303 engravings. 8vo.
- PEMBERTON, C. R. Practical Treatise on various Diseases of the Abdominal Viscera. Engravings. 8vo.
- PEREIRA, J. Elements of Materia Medica and Therapeutics. With Notes and Additions by Joseph Carson, M.D. 279 illustrations. 2 vols. 8vo.
- Treatise on Food and Diet ; with observations on the Dietetical Regimen suited for Disordered States of the Digestive Organs ; and an account of the Dietaries of some of the principal Metropolitan and other Establishments for Paupers, Lunatics, Criminals, Children, the Sick, &c. Edited by Charles A. Lee, M.D. 8vo.
- PETTIGREW, T. J. On Superstitions connected with the History and Practice of Medicine and Surgery. 12mo.
- Medical Portrait Gallery ; or Biographical Memoirs of the most celebrated Physicians, Surgeons, &c., who have contributed to the advancement of Medical Science. 60 portraits. 4 vols. 8vo.
- Pharmacopœia of the United States of America. By authority of the National Medical Convention of 1840. 8vo.
- PHILIP, A. P. W. Experimental Inquiry into the Laws of the Vital Functions. 8vo.
- Treatise on the Nature and Cure of those Diseases, either Acute or Chronic, which precede Change of Structure ; with a view to the preservation of Health, and particularly the prevention of Organic Diseases. With Notes, &c., by J. H. Miller, M.D. 8vo.
- Treatise on Febrile Diseases, including the various species of Fever, and all Diseases attended with Fever. With Additions by Nathan Smith, M.D. 2 v. 8vo.
- Treatise on Indigestion and its Consequences, called Nervous and Biliary Complaints ; with observations on the Organic Diseases in which they sometimes terminate. 8vo.

- PHILIP, A. P. W.** Treatise on Protracted Indigestion and its Consequences ; being the application to the Practical department of Medicine of the results of an inquiry into the Laws of the Vital Functions, &c. 8vo.
- PHILLIPS, B.** A Treatise on the Urethra ; its Diseases, especially Stricture, and their cure. 2 plates. 8vo.
- PHILLIPS, W.** Elementary Treatise on Mineralogy : containing the latest Discoveries in American and Foreign Mineralogy, &c., by Francis Alger. 600 illustrations. 8vo.
- Phrenology**, Epitome of ; being an Outline of the Science as taught by Gall, Spurzheim, and Combe. Engravings. 18mo.
- PILCHER, G.** Treatise on the Structure, Economy, and Diseases of the Ear. 16 plates. 8vo.
- PINNEY, J.** The Alternative ; Disease and premature Death, or Health and Long Life : being an exposure of the prevailing misconception of their respective sources, and showing what is, and what is not according to those laws which alone can insure sound Health and Length of Life. 8vo.
- PLENCK, J. J.** Hygology, or Chemico-Physiological Doctrine of the Fluids of the Human Body. Translated by R. Hooper, M.D. 8vo.
- PLUMBE, S.** A Practical Treatise on Diseases of the Skin, their Constitutional Causes and Local Character, &c. Coloured plates. 8vo.
- Pocket Anatomist** : a summary Description of the Muscles ; with a Tabular View of the Arteries and Nerves. 32mo.
- POE, E. A.** Conchologist's First Book ; a System of Testaceous Malacology, for the use of Schools, in which the Animals, according to Cuvier, are given with the shells, &c. 12 plates. 12mo.
- PORTAL, A.** Observations sur la Nature et le Traitement de l'Apoplexie, et sur les moyens de la prevenir. 8vo.
- PORTER, A. L.** Chemistry of the Arts ; being a practical display of the Arts and Manufactures which depend on Chemical Principles. On the Basis of Gray's Operative Chemist, adapted to the United States ; with Treatises on Calico Printing, Bleaching, &c. 67 plates. 2 vols. 8vo.
- POST, A. C.** Observations on the Cure of Strabismus ; with an Appendix on the New Operation for the Cure of Stammering. 7 coloured plates. 18mo.
- POTT, PERCIVAL.** Treatise on the Hydrocele or Water Rupture, and other Diseases of the Testicle, its Coats, and Vessels. Plates. 8vo.
- **Chirurgical Works** : containing Observations on Injuries of the Head,—Fistula Lachrymalis,—Fractures and Dislocations,—Ruptures,—Hydrocele,—Fistula in Ano,—Polypus of the Nose,—Mortification of the Toes and Feet,—Palsy of the Lower Limbs,—Tumours,—Hæmorrhoidal Excrescences, &c., &c. With Notes, &c., by Sir James Earle, F.R.S. Plates. 2 vols. 8vo.
- POWERS, G.** Essay upon the Influence of the Imagination on the Nervous System, contributing to a False Hope in Religion. 12mo.
- Prescriber's Pharmacopœia** : containing all the Medicines in the London Pharmacopœia, arranged in classes according to their Action, with their Composition and Doses. Revised and improved by an American Physician. 32mo.
- PRESTWICH'S** Dissertation on Mineral, Animal, and Vegetable Poisons ; containing a description of Poisons in general, their manner of Action, Effects on the Human Body, and respective Antidotes. Plates. 8vo.
- PRICHARD, J. C.** Treatise on Insanity and other Disorders affecting the Mind. 8vo.
- **Researches into the Physical History of Mankind.** Plates, Maps, &c. 4 vols. 8vo.
- PRIESSNITZ, V.** Hydriatics ; or Manual of the Water Cure, especially as practised by Vincent Priessnitz in Græffenberg. Translated by F. Græter. Plate. 18mo.
- PRINGLE, SIR JOHN.** Observations on the Diseases of the Army. With Notes by Benjamin Rush, M.D. 8vo

- PROUT, W. Nature and Treatment of Stomach and Renal Diseases; being an inquiry into the connection of Diabetes, Calculus, and other affections of the Kidney and Bladder, with Indigestion. 6 plates. 8vo.
- Same work, with Notes and Additions by S. Colhoun, M.D. Plate. 8vo.
- PUERPERAL Fever. History, Pathology, and Treatment of Puerperal Fever and Cru-
ral Phlebitis. By Drs. A. Gordon, W. Hey, J. Armstrong, and R. Lee. With an
Introductory Essay by Charles D. Meigs, M.D. 8vo.
- PUGH, B. Treatise on Midwifery, chiefly with regard to the Operation, &c. 11
plates. 8vo.
- PUGH, J. A Physiological, Theoretic, and Practical Treatise on the Utility of the
Science of Muscular Action, for restoring the Power of the Limbs. 15 plates. 4to.
- QUAIN, J. Elements of Anatomy. 8vo.
- QUAIN, J., AND W. J. E. WILSON. A Series of Anatomical Plates, with References
and Physiological Comments, illustrating the Arteries of the Human Body. With
Notes and Additions by J. Pancoast, M.D. 34 coloured plates. 4to.
- A Series of Anatomical Plates with References and Physiological Comments
illustrating the Arteries and Veins of the Human Body. With Notes and Ad-
ditions by J. Pancoast, M.D. 50 coloured plates. 4to.
- A Series of Anatomical Plates with References and Physiological Comments
illustrating the Structure of the different parts of the Human Body. With Notes
and Additions by J. Pancoast, M.D. 200 plates. 4to.
- RACIBORSKI, A. Elementary Treatise on Auscultation and Percussion, or the Ap-
plication of Acoustics to the Diagnosis of Diseases, with a Synoptical Table.
With Notes, by M. Post, M.D. Plate. 8vo.
- RAMADGE, F. H. Consumption Curable: and the manner in which Nature as well
as remedial Art operates in effecting a Healing Process in cases of Consumption,
&c. Also a Mode of Treatment by which the development of Tubercles may
be prevented in persons liable thereto, &c. 8vo.
- RANSAY, A. A Series of Plates of the Heart, Cranium, and Brain, in imitation of
Dissections. 15 coloured plates. 4to.
- RAMSEOTHAM, F. H. Principles and Practice of Obstetric Medicine and Surgery, in
reference to the process of Parturition. 55 plates. 8vo.
- RAMSEOTHAM, J. Practical Observations in Midwifery. With Notes by W. P.
Dewees, M.D. 8vo.
- RANSDEN, T. Practical Observations on the Slerocele and other Morbid enlarge-
ments of the Testicle; also on the Cause and Cure of the Acute, the Spurious,
and the Chronic Hydrocele, &c. 2 plates. 8vo.
- RASPAIL, F. V. A New System of Organic Chemistry. With Additions by W.
Henderson, M.D. 12 plates. 8vo.
- RATIER, F. S. Practical Formulary of the Parisian Hospitals; exhibiting the Pre-
scriptions employed by the Physicians and Surgeons of those Establishments; re-
marks illustrative of their Doses, Mode of Administration, and Appropriate Ap-
plication, &c., &c. With Notes by R. D. M'Lellan, M.D. 18mo.
- Medical Guide to Paris; or a Description of the principal Hospitals of Paris,
with an account of the Practice of the most eminent Physicians and Surgeons
attached to the different Hospitals. With Additions by J. R. Alcock, M.D. 18mo.
- RAY, J. Treatise on the Medical Jurisprudence of Insanity. 12mo.
- RAY, P. A Theoretical and Practical Treatise on the Diseases of the Skin.
With Notes and Additions by John Bell, M.D. 40 coloured plates. 4to.
- REESE, D. M. Medical Lexicon of Modern Terminology; being a Complete Vo-
cabulary of Definitions, including all the Technical Terms employed by Writers
and Teachers of Medical Science at the present day, &c. 32mo.
- REID, T. Essay on the Active Powers of the Human Mind; an Inquiry into the
Human Mind on the Principles of Common Sense; and an Essay on Quantity. 8vo
- REID, J. Essays on Hypochondriacal and other Nervous Affections. 8vo.

- RENNIE, J. New Supplement to the latest Pharmacopœias of London, Edinburgh, Dublin, and Paris, forming a complete Dispensatory, Conspectus, and Dictionary of Medical Chemistry, giving all the old and new names, including the New French and American Medicines and Poisons; with Symptoms, Treatment, and Tests, &c., &c. 8vo.
- RICHARDSON, C. J. A Popular Treatise on the Warming and Ventilation of Buildings; showing the advantages of the improved system of Heated Water Circulation, &c. 18 plates. 8vo.
- RICHERAND, A. Elements of Physiology. Translated by G. J. M. De Lys, M.D. With Notes, &c., by James Copland, M.D. 8vo.
- RICORD, P. Practical Treatise on Venereal Diseases; or critical and experimental Researches on Inoculation, applied to the study of these Affections. with a Therapeutical Summary and Special Formulary. 8vo.
- RIGBY, E. A System of Midwifery. Numerous engravings. 8vo.
 ——— Essay on the Theory of the Production of Animal Heat, and on its application in the Treatment of Cutaneous Eruptions, Inflammations, and some other Diseases. 8vo.
- RILLIET AND BARTHEZ's Treatise on the Pneumonia of Children. Translated by S. Parkman, M.D. 8vo.
- ROBERTSON, A. Conversations on Anatomy, Physiology, and Surgery. 18mo.
- ROBERTSON, W. Practical Treatise on the Human Teeth: showing the causes of their destruction, and the means of their preservation. 6 plates. 8vo.
- ROGET, P. M. Outlines of Physiology: with an Appendix on Phrenology. 8vo.
 ——— Animal and Vegetable Physiology, considered with reference to Natural Theology. (A Bridgewater Treatise.) 463 engravings. 2 vols. 8vo.
- ROKITANSKY, C. Treatise on Pathological Anatomy. With Additions by Dr. John C. Peters. 8vo.
- ROOT, J. The Horrors of Delirium Tremens. 8vo.
- ROSE, J. S. Consumption Curable, a Practical Treatise to prove Consumption a manageable Disease. 12mo.
- ROUPEL, G. L. A Short Treatise on Typhus Fever. 8vo.
- ROWLAND, R. Treatise on Neuralgia. 8vo.
- ROYLE, J. F. A Manual of Materia Medica and Therapeutics, with illustrations. Revised and adapted to the Pharmacopœia of the United States.
- RUCCO, J. Introduction to the Science of the Pulse, as applied to the Practice of Medicine. 2 vols. 8vo.
- RUOFF's Repertory of Homœopathic Medicine, Nosologically arranged. Translated by A. H. Okie, M.D. With Additions, &c., by G. Humphrey, M.D. 12mo.
- RUSH, BENJAMIN. Medical Inquiries and Observations upon the Diseases of the Mind. 8vo.
 ——— Sixteen Introductory Lectures to Courses of Lectures upon the Institutes and Practice of Medicine, with a Syllabus of the latter: with two Lectures upon the Pleasures of the Senses and of the Mind, and an inquiry into their proximate cause. 8vo.
 ——— Medical Inquiries and Observations on Cause of Animal Life—Natural History of Medicine among the Indians—Influence of Physical Causes upon the Moral Faculty—Effects of Ardent Spirits upon the Human Mind and Body—Tetanus—Disease caused by drinking Cold Water—Worms and Anthelmintic Medicines—External Use of Arsenic in cure of Cancers—Sore Legs—State of Body and Mind in Old Age—Duties of a Physician—Salt in the cure of Hæmoptysis—Pulmonary Consumption—Dropsy—Dropsy in the Brain—Gout—Hydrophobia—Cholera Infantum—Cynanche Trachealis—Bilious Remitting Fever—Scarlatina Anginosa—Measles—Influenza—Yellow Fever, &c., &c. 2 vols. 8vo.
- RUSSELL, D. The Invalid's Friend, being a plain practical work, designed for the use of Families and Individuals, on Vegetable and Botanical Principles. 12mo.

- RYAN, M. A Manual of Midwifery, or Compendium of Gynæcology and Paidon-
ology ; comprising a new nomenclature of Obstetric Medicine, with a concise
account of the symptoms and treatment of the most important Diseases of Wo-
men and Children, &c. 8vo.
- A Manual of Medical Jurisprudence, compiled from the best Medical and
Legal Works. 8vo.
- Lectures on Population, Marriage, and Divorce, as Questions of State Med-
icine, comprising an account of the causes and treatment of Impotence and Ste-
rility, &c. 12mo.
- RYLAND, F. Treatise on the Diseases and Injuries of the Larynx and Trachea. 8vo
- SAISSY, J. A. Essay on the Diseases of the Internal Ear. Translated by N. R.
Smith, M.D. Plate. 8vo.
- SALMON, F. A Practical Essay on Stricture of the Rectum ; illustrated by cases,
showing the connexion of that Disease with affections of the Urinary Organs and
the Uterus, with Piles and various constitutional complaints. 8vo.
- Practical Observations on Prolapsus of the Rectum. 4 plates. 8vo.
- SARLANDIERE, J. Systematized Anatomy ; or Human Organography, in Synop-
tical Tables. Translated by W. C. Roberts, M.D. 15 plates, folio.
- SAUNDERS, J. C. Anatomy of the Human Ear, illustrated by a series of engravings
of the natural size : with a Treatise on the Diseases of that Organ, the Causes
of Deafness, and their proper Treatment. 8vo.
- Treatise on some practical points relating to the Diseases of the Eye. With
a short account of the Author's Life, and his method of curing the Congenital
Cataract, by J. R. Farre. 7 coloured plates. 8vo.
- SAUNDERS, W. Treatise on the Structure, Economy, and Diseases of the Liver :
with an Inquiry into the properties and component parts of the Bile and Biliary
Concretions. 12mo.
- SAWYER, M. E. Treatise on Primitive or Secondary Disguised or Misplaced Fever,
as a single Disease ; with the varieties, cause, and treatment, as it appears in
most of the particular forms of Fever recognized by Nosologists. 12mo.
- SAYERS, E. American Fruit Garden Companion, being a practical treatise on the
Propagation and Culture of Fruit. 18mo.
- Treatise on the Culture of the Dahlia and Cactus. 18mo.
- SCARPA, A. Treatise on the Principal Diseases of the Eyes. With Notes by James
Briggs. Plates. 8vo.
- Treatise on the Minute Anatomy of the Bones. 18mo.
- SCHILL, PROF. Outlines of Pathological Semeiology. With Notes by D. Spillan, M.D.
- SCOUTETTEN, H. Memoir of the Radical Cure of Club-Foot. Translated by F. C.
Stewart, M.D. 6 plates. 8vo.
- SCUDAMORE, C. Treatise on the Nature and Cure of Gout and Rheumatism, in-
cluding general considerations on Morbid States of the Digestive Organs ; re-
marks on Regimen and on Gravel. 8vo.
- SENAC, J. Treatise on the Hidden Nature, and the Treatment of Intermittent and
Remitting Fevers. Translated by C. Caldwell, M.D. Preface by Benj. Rush,
M.D. 8vo.
- SERNY, J. B. Spinal Curvature, its Consequences, and its Cure. 15 plates. 8vo.
- SEVERN, C. First Lines of the Practice of Midwifery ; with remarks on the Foren-
sic Evidence requisite in cases of Feticide and Infanticide. 2 plates. 8vo.
- SHAW, EDWARD B. L. Medical Remembrancer, or Book of Emergencies ; in
which are concisely pointed out the Immediate Remedies to be adopted in the first
moments of danger from Poisoning, Drowning, Apoplexy, Burns, and other ac-
cidents ; with the Tests for the principal Poisons, and other useful information. 32mo.
- SHAW, J. Manual for the Student of Anatomy ; containing Rules for Displaying
the Structure of the Body, so as to exhibit the elementary views of Anatomy, and
their application to Pathology and Surgery. With Notes by W. Anderson, M.D.
2 plates. 12mo.

- SHERWOOD, H. H. Motive Power of Organic Life, and Magnetic Phenomena of Terrestrial and Planetary Motions, with the application of the ever-active and all-pervading agency of Magnetism, to the nature, symptoms, and treatment of Chronic Diseases. Many engravings. 8vo.
- Manual for Magnetizing with the Rotary and Vibrating Magnetic Machine, in the Duodynamic Treatment of Diseases. Engravings. 32mo.
- SHEW, J. Hand Book of Hydropathy ; or a Popular Account of the treatment and prevention of Diseases, by means of Water. Plate. 12mo.
- Hydropathy ; or the Water-Cure : its principles, modes of treatment, &c. Plate. 12mo.
- SHEW, M. L. Water-Cure for Ladies ; a popular work on the Health, Diet, and Regimen of Females and Children, and the prevention and cure of Diseases ; with a full account of the Processes of Water-Cure. Revised by J. Shew. 2 plates. 12mo.
- SKINNER, J. S. The Dog and the Sportsman, embracing the Uses, Breeding, Training, Diseases, &c., of Dogs, and an account of the different kinds of Game, with their habits, &c. Engravings. 12mo.
- SMELLIE, W. Treatise on the Theory and Practice of Midwifery. 3 vols. 12mo.
- Anatomical Plates of Midwifery, with concise explanations, for the use of Students. 14 plates. 8vo.
- Philosophy of Natural History. With Additions, &c., by John Ware, M.D. 12mo.
- SMETHURST, T. Hydrotherapia ; or the Water Cure : being a practical view of the Cure in all its bearings, exhibiting the great utility of Water as a preservative of Health and remedy for Disease : with a description of Græffenberg, and the system there, as practised by Priessnitz ; History of the Water Cure from the remotest Antiquity, &c. 8vo.
- SMITH, DANIEL B. Principles of Chemistry. 12mo.
- SMITH, HENRY H. Anatomical Atlas, illustrative of the Structure of the Human Body. Under the supervision of W. E. Horner, M.D. 636 engravings. 8vo.
- Minor Surgery ; or, Hints on the every-day duties of the Surgeon. 189 engravings. 18mo.
- SMITH, SIR JAMES EDWARD, M.D. Memoir of his Life and Correspondence. 2 plates. 2 vols. 8vo.
- SMITH, JEROME V. C. Class Book of Anatomy, explanatory of the first principles of Human Organization as the Basis of Physical Education. 110 engravings. 12mo.
- SMITH, JOHN AUGUSTINE. Select Discourses on the Functions of the Nervous System, in opposition to Phrenology, Materialism, and Atheism ; with a Lecture on the Diversities of the Human Character, arising from Physiological peculiarities. 12mo.
- SMITH, NATHAN. Practical Essay on Typhus Fever. 8vo.
- Medical and Surgical Memoirs : containing Essay on Typhus Fever, — Necrosis, — Fractures of the Femur, — Apparatus for Fractures of the Thigh and Leg, — Dislocation of Hip Joint, — Operation for Hydrocele, — Amputation, — Extirpation of Ovarian Sac, &c. Edited, with Addenda, by N. R. Smith. 7 plates. 8vo.
- SMITH, NATHAN R. Surgical Anatomy of the Arteries. 18 coloured plates. 4to.
- SMITH, SAMUEL STANHOPE. An Essay on the Causes of the Variety of Complexion and Figure in the Human Species. 8vo.
- SMITH, SOUTHWOOD. Treatise on Fever. 8vo.
- SMYTH, J. C. Effect of Nitrous Vapour in preventing and destroying Contagion with an Introduction respecting the nature of the Contagion which gives rise to Jail or Hospital Fever, &c. 8vo.
- SNELL, J. Practical Guide to Operations on the Teeth. With a historical sketch of the Rise and Progress of Dental Surgery. 5 plates. 8vo.
- SPALLANZANI, L. Memoirs on Respiration. Edited by J. Senebier. 8vo.
- Experiments upon the Circulation of the Blood, throughout the Vascular System ; on Languid Circulation ; on the Motion of the Blood, independent of

- the action of the Heart; and on the Pulsations of the Arteries. With Notes by R. Hall, M.D. 8vo.
- SPENDER, J. C. Observations on the Causes and Treatment of Ulcerous Diseases of the Leg. 8vo.
- SPOONER, S. Guide to Sound Teeth, or a Popular Treatise on the Teeth, illustrating the whole judicious management of these Organs from Infancy to Old Age, &c. 12mo.
- SPURZHEIM, J. G. Anatomy of the Brain, with a general view of the Nervous System. Translated by R. Willis, M.D., and revised by C. H. Stedman, M.D. 18 plates. 8vo.
- Observations on the Deranged Manifestations of the Mind, or Insanity. With an Appendix by A. Brigham, M.D. 4 plates. 8vo.
- Phrenology, or the Doctrine of the Mental Phenomena. 15 plates. 2 vols. 8vo.
- Phrenology, in connexion with the Study of Physiognomy. With a Biography of the Author, by N. Capen. 35 plates. 8vo.
- STAFFORD, R. A. An Essay upon the Treatment of the Deep and Excavated Ulcer: with Cases. 8vo.
- STEEL, J. H. Analysis of the Mineral Waters of Saratoga and Ballston, with practical remarks on their Medical properties, &c.
- STEVENS, R. A New Synopsis, or the Natural Order of Diseases; containing their Definition, Principles, and Treatment, with a new Pathology of Fever and Inflammation. 8vo.
- STEVENSON, J. Cataract; a familiar description of its Nature, Symptoms, and ordinary modes of treatment. 12mo.
- A Simplified Practical Guide to the Cure of Venereal Complaints. 18mo.
- STEWART, J. Practical Treatise on the Diseases of Children. 8vo.
- STEWART, F. C. Hospitals and Surgeons of Paris: a Historical and Statistical Account of the Civil Hospitals of Paris; Biographical Notices of some of the most eminent of the living Parisian Surgeons, &c. 8vo.
- STOKES, W. Treatise on the Diagnosis and Treatment of Diseases of the Chest. 8vo.
- Researches on the State of the Heart, and the Use of Wine in Typhus Fever. 8vo.
- STROTHER, E. Essay on Sickness and Health; containing all necessary cautions and directions for the regulation of Diseased and Healthy persons. 8vo.
- STRUVE, C. A. Practical Essay on the Art of Recovering Suspended Animation: with a review of the most proper and effectual means to be adopted in cases of imminent danger. 18mo.
- SULLY, H. Observations on, and Plain Directions for, all classes of People, to prevent the fatal effects of the Bites of Animals labouring under Hydrophobia. Plate. 8vo.
- Surgery, a Synopsis of Modern Practical: by a Member of the Royal College of Surgeons, London. 12mo.
- SWEDIAUR, F. Complete Treatise on the Symptoms, Effects, Nature, and Treatment of Syphilis. Translated by T. T. Hewson, M.D. 8vo.
- SWEETSER, W. Treatise on Consumption; embracing an inquiry into the influence exerted upon it by Journeys, Voyages, and Change of Climate: with Directions for the Consumptive visiting the South of Europe, and remarks upon its Climate. 8vo.
- Mental Hygiene, or an Examination of the Intellect and Passions, designed to illustrate their influence on Health and the Duration of Life. 12mo.
- A Treatise on Digestion, and the Disorders incident to it which are comprehended under the term Dyspepsia. 12mo.
- SYDENHAM, T. Works of Thomas Sydenham, M.D., on Acute and Chronic Diseases; wherein their histories and modes of cure, as recited by him, are delivered with accuracy and perspicuity, with Notes from the most eminent Medical Writers, &c., &c. Edited by George Wallis, M.D. 2 vols. 8vo.

- SYME, J. Principles of Surgery. 8vo.
 — Diseases of the Rectum. 8vo.
- TAVERNIER, A. Elements of Operative Surgery. With Notes, &c., by S. D. GROSS, M.D. 8vo.
- TAYLOR, A. S. On Perforations of the Stomach, from Poisoning and Disease. 8vo.
 — Medical Jurisprudence. With Notes and Additions by R. E. Griffith, M.D. 8vo.
- TEALE, T. P. Treatise on Neuralgic Diseases, dependent upon irritation of the Spinal Marrow and Ganglia of the Sympathetic Nerve. 12mo.
- TESTE, A. Practical Manual of Animal Magnetism; containing an exposition of the methods employed in producing the Magnetic Phenomena; with its application to the treatment and cure of Diseases. Translated by D. Spillan, M.D. 12mo.
- THACHER, J. Observations on Hydrophobia, produced by the bite of a mad Dog, or other rabid animal: with an examination of the various theories and methods of cure; and an inquiry into the merit of specific remedies, &c. 8vo.
 — American Modern Practice; or a simple method of Prevention and Cure of Diseases, according to the latest improvements and discoveries, &c. 8vo.
 — American Medical Biography; or Memoirs of eminent Physicians who have flourished in America: with a succinct History of Medical Science in the United States, from the first settlement of the Country. 15 portraits. 8vo.
- THACKRAH, C. T. The Effects of the Principal Arts, Trades, and Professions, and of Civic States and Habits of Living, on Health and Longevity, &c. 18mo.
- THENARD, L. J. Essay on Chemical Analysis. With Additions by J. G. Children, F.R.S. Plates. 8vo.
- THOMAS, R. Modern Practice of Physic, exhibiting the Characters, Causes, Symptoms, Prognostics, Morbid Appearances, and improved method of treating the Diseases of all Climates. With an Appendix, by David Hosack, M.D. 8vo.
- THOMSON, J. Lectures on Inflammation, exhibiting a view of the general doctrines, Pathological and Practical, of Medical Surgery. 8vo.
- THOMSON, W. Historical Notices on the occurrence of Inflammatory Affections of the Internal Organs, after External Injuries and Surgical Operations. 8vo.
 — Practical Treatise on the Diseases of the Liver and Biliary Passages. Also Clinical Illustrations of Diseases of the Liver and Spleen; by W. Twining, M.D. 8vo.
- THOMSON, A. T. Conspectus of the Pharmacopœias of the London, Edinburgh, and Dublin Colleges of Physicians, and of the United States Pharmacopœia; being a practical Compendium of Materia Medica and Pharmacy. Edited by Charles A. Lee, M.D. 18mo.
 — The Domestic Management of the Sick-Room, necessary, in aid of Medical Treatment, for the cure of Diseases. With Additions by R. E. Griffith, M.D. Engravings. 12mo.
- THOMSON, T. A System of Chemistry of Inorganic Bodies: with illustrations. 2 vols. 8vo.
- THORBURN, J. S. Elements of Bedside Medicine and General Pathology, or General Disease-Discourse, with a sketch of the origin, progress, and prospects of Clinical Medicine and Surgery, &c. 8vo.
- TICKNOR, C. Popular Treatise on Medical Philosophy; or an Exposition of Quackery and Imposture in Medicine. 12mo.
 — Same work, with Notes by W. Wright. 8vo.
 — The Philosophy of Living; or the Way to enjoy Life and its Comforts. Engravings. 18mo.
- TIEDEMANN, F. Anatomy of the Fœtal Brain; with a comparative exposition of its Structure in Animals. Translated by W. Bennett, M.D. 14 plates. 12mo.
- TIMBREL, W. H. Practical Observations on the Management of Ruptures. With engravings. 12mo.
- TORREY, J., AND A. GRAY. A Flora of North America, containing abridged Descrip-

tions of all the known Indigenous and Naturalized Plants, growing north of Mexico; arranged according to the Natural System. 8vo.

TOWNSEND, C. H. Facts in Mesmerism, with reasons for a dispassionate Inquiry into it. Engravings. 12mo.

TOWNSEND, P. S. Account of the Yellow Fever, as it prevailed in the city of New York, in 1822. 8vo.

TRAILL, T. S. Outlines of a Course of Lectures on Medical Jurisprudence. 8vo.

Transactions of the Medical Society of the State of New York. 3 vols. 8vo.

TRAVERS, B. Inquiry concerning that disturbed state of the Vital Functions usually denominated Constitutional Irritation. 8vo.

——— Synopsis of Diseases of the Eye, and their Treatment: to which are prefixed a short Anatomical Description, and a sketch of the Physiology of that organ. With Notes and Additions by Edward Delafield, M.D. Coloured Plates. 8vo.

TRAVERS, B., AND J. H. GREEN. Principles and Practice of Ophthalmic Surgery: comprising the Anatomy, Physiology, and Pathology of the Eye, with the treatment of its Diseases. Edited by A. C. Lee. Coloured plate. 18mo.

TREDGOLD, T. Principles of Warming and Ventilating Public Buildings, Dwelling Houses, Manufactories, Hospitals, Hot-Houses, Conservatories, &c., and of constructing Fire-Places, Boilers, Steam-Apparatus, Grates, and Drying-Rooms, &c. With an Appendix by T. Bramah, comprising Observations on Heating by means of Warm Water, &c. 12 plates. 8vo.

TRIMMER, J. Practical Geology and Mineralogy; with instructions for the Qualitative Analysis of Minerals. 212 illustrations. 8vo.

TROTTER, T. A View of the Nervous Temperament; being a practical inquiry into the increasing prevalence, prevention, and treatment of those Diseases commonly called Nervous, Bilious, Stomach, and Liver Complaints; Indigestion; Low Spirits; Gout, &c. 8vo.

TROUSSEAU, A., AND H. BELLOC. A Practical Treatise on Laryngeal Phthisis, Chronic Laryngitis, and Diseases of the Voice. Translated by J. A. Warder, M.D. 8vo.

TURNBULL, A. Treatise on Painful and Nervous Diseases, and on a New Mode of Treatment for Diseases of the Eye and Ear. Plate. 8vo.

——— An Investigation into the remarkable Medicinal effects resulting from the external application of Veratria. 8vo.

——— Medical Properties of the Natural Order Ranunculacæ; and more particularly on the Uses of Sabadilla Seeds, Delphinium Staphisagria, and Aconitum Napellus, and their Alcaloids, Veratria, Sabadilline, Delphinia, and Aconitine. 8vo.

TURNBULL, W. A Popular Treatise on Health, and the means of preserving it; or the Dyspeptic's Pocket Companion. 18mo.

TURNER, E. Elements of Chemistry, including the recent discoveries and doctrines of the Science. With Notes and Additions by Franklin Bache, M.D. 8vo.

TUSON, E. W. The Dissector's Guide; or Student's Companion: illustrated by numerous engravings, clearly exhibiting and explaining the Dissection of every part of the Human Body. With Additions by W. Lewis, Jr., M.D. 12mo.

TWEEDIE, A. A System of Practical Medicine, comprised in a series of original Dissertations. With Notes and Additions by W. W. Gerhard, M.D. 3 vols. 8vo.

——— Dissertations on Nervous Diseases, by Drs. Hope, Prichard, Bennett, Taylor, and Thomson. Edited by A. Tweedie, M.D. With Notes and Additions by W. W. Gerhard, M.D. 8vo.

——— Dissertations on Diseases of the Organs of Respiration, by Drs. Williams, Thomson, Carpenter, and Joy. Edited by A. Tweedie, M.D. With Notes and Additions by W. W. Gerhard, M.D. 8vo.

——— Clinical Illustrations of Fever. 8vo.

UNDERWOOD, M. Treatise on the Diseases of Children: with Directions for the Management of Infants. With Notes by S. Merriman, M.D., Marshall Hall, M.D., and John Bell, M.D. 8vo.

- URE, A. Practical Compendium of the Materia Medica, adapted to the treatment of the Diseases of Infancy and Childhood; with numerous Prescriptions. 18mo.
 ——— A Dictionary of Arts, Manufactures, and Mines; containing a clear exposition of their principles and practice. 1240 engravings. 8vo.
 ——— Recent Improvements in Arts, Manufactures, and Mines; being a Supplement to his Dictionary. 190 engravings. 8vo.
- UTERUS. Lectures on the Functions and Diseases of the Womb, by C. Waller, M.D.
 Lectures on Diseases of the Uterus and its Appendages, by M. Lisfranc.
 Clinical Lectures on Diseases of the Puerperal State, by J. T. Ingleby. 8vo.
- VAUGHAN, W. An Essay on Head Aches, and on their Cure. 8vo.
- VELPEAU, ALF. A. L. M. A Treatise on Surgical Anatomy; or the Anatomy of Regions, considered in its relations with Surgery. With Notes by John W. Sterling, M.D. 14 plates. 2 vols. 8vo.
 ——— New Elements of Operative Surgery: with a Treatise on Minor Surgery, illustrated by over 300 engravings. Translated by P. S. Townsend, M.D., under the supervision of Valentine Mott, M.D. 3 vols. 8vo., and 1 quarto vol. of 22 plates
 ——— Elementary Treatise on Midwifery; or, Principles of Tokology and Embryology. With Notes by C. D. Meigs, M.D., and Additions by W. Harris, M.D. 8vo.
- VELPEAU, M. A Treatise on Diseases of the Breast. Translated by S. Parkman, M.D. 8vo.
- VOGHT, J. G. Treatise on Bowel Complaints. 12mo.
- WADD, W. Nugæ Chirurgicæ; or a Biographical Miscellany, illustrative of a collection of Professional Portraits. 8vo.
- WAGNER, R. Elements of the Comparative Anatomy of the Vertebrate Animals Edited by Alfred Tulk. 8vo.
- WAITE, G. The Gums; with late discoveries on their Structure, Growth, Connections, Diseases, and Sympathies. 8vo.
 ——— Surgeon Dentist's Anatomical and Physiological Manual. 18mo.
- WALKER, A. Pathology, founded on the Natural System of Anatomy and Physiology. 12mo.
 ——— The Nervous System, Anatomical and Physiological: in which the Functions of the various parts of the Brain are for the first time assigned, &c. 8vo.
- WALKER, J. The Philosophy of the Eye; being a familiar exposition of its Mechanism, and of the phenomena of Vision, with a view to the evidence of design. Numerous illustrations. 12mo.
- WALLACE, W. C. Wonders of Vision, a Treatise on the Eye; containing discoveries of the causes of near and far-sightedness, and of the affections of the retina, &c. 55 engravings. 12mo.
- WALLER, J. A Treatise on the Incubus, or Night-Mare, Disturbed Sleep, Terrific Dreams, and Nocturnal Visions; with the means of removing these distressing complaints. 12mo.
- WALSH, W. H. Anatomy, Physiology, Pathology, and Treatment of Cancer. With Additions by J. M. Warren, M.D. Engravings. 12mo.
 ——— Physical Diagnosis of Diseases of the Lungs. 18mo.
- WARRE, J. Chirurgical Observations relative to the Eye: with an Appendix on the Introduction of the Male Catheter, and the treatment of Hæmorrhoids. Plates. 2 vols. 8vo.
- WARREN, G. A Commentary, with Practical Observations on Disorders of the Head, in which is particularly considered the propriety of Bleeding. 8vo.
- WARREN, J. C. Surgical Observations on Tumours, with Cases and Operations. 16 coloured plates. 8vo.
- WARRINGTON, J. The Nurse's Guide: containing a series of Instructions to Females who wish to engage in the important business of Nursing Mother and Child in the Lying-in Chamber. 18mo.
 ——— Obstetric Catechism. 12mo.
- WATERHOUSE, B. Essay concerning Tussis Convulsiva, or Whooping-Cough: with Observations on Diseases of Children. 8vo.

- WATSON, T. Lectures on the Principles and Practice of Physic. With Additions by D. F. Condie, M.D. 8vo.
- WEATHERHEAD, G. H. Practical Treatise on the principal diseases of the Lungs, considered especially in relation to the particular Tissues affected, illustrating the different kinds of Cough. 8vo.
- WEBB, E. Philosophy of Medicine, deduced from analyzing the Constitution of Man, and its Legitimate Structure and Action. 8vo.
- WEBER, M. J. Anatomical Atlas of the Human Body, in Natural Size. 90 plates, 23 by 30 inches. Forty-four of these plates, forming eleven full-length figures, are mounted on canvass with Roller, and are very appropriate and useful ornaments for the Lecture-Room or Office. They are,
- | | |
|----------------------------|------------------------------|
| Male Adult Skeleton, | front view. |
| Do. do. do. | posterior view. |
| Ligaments, | front do. |
| Do. | posterior do. |
| Four Bodies of Muscles, | front, back, and side views. |
| Vascular and Nervous Body, | coloured. |
| Blood-Vessels and Nerves, | front view, coloured. |
| Do. do. do. | posterior view, do. |
- WEBSTER, J. W. Manual of Chemistry, containing the principal Facts of the Science, in the order in which they are discussed and illustrated in the Lectures of Harvard University and other Medical Schools in the United States. 200 illustrations. 8vo.
- WEBSTER, NOAH. A Brief History of Epidemic and Pestilential Diseases; with the principal Phenomena of the Physical World, which precede and accompany them, and Observations deduced from the Facts stated. 2 vols. 8vo.
- WELLER, C. H. Manual of the Diseases of the Human Eye, intended for Surgeons commencing Practice, from the best National and Foreign Works, and in particular those of Professor Beer. Translated by G. C. Monteath, M.D. Coloured plates. 2 vols. 8vo.
- WELLS, H. An Essay on Teeth; comprising a brief description of their Formation, Diseases, and proper Treatment. 12mo.
- WELLS, W. C. An Essay on Dew, and several appearances connected with it. 8vo
- WHITE, E. L. A Popular Essay on the Disorder familiarly termed a Cold; in which the means of obviating the various causes are explained; with a collection of approved Receipts, and observations on the most popular Remedies, &c. With Notes by J. Stewart, M.D. 12mo.
- WHITLAW, C. New Medical Discoveries, with a Defence of the Linnæan Doctrine, and a Translation of his Vegetable Materia Medica. 8vo.
- Treatise on the Causes and Effects of Inflammation, Fever, Cancer, Scrofula, and Nervous Affections; observations on the correctness of Linnæus's Classification of Diseases, with remarks on the specific action of his Patent Medicated Vapour Bath, and Rules for Diet and Regimen. 8vo.
- WILLAN, R. On Cutaneous Diseases. 8vo.
- WILLIAMS, C. J. B. Practical Treatise on the Diseases of the Respiratory Organs including Diseases of the Larynx, Trachea, Lungs, and Pleura. With Additions, &c., by M. Clymer, M.D. Engravings. 8vo.
- Principles of Medicine: comprising General Pathology and Therapeutics, and a brief general view of Etiology, Nosology, Semeiology, Diagnosis and Prognosis. 8vo.
- A Rational Exposition of the Physical Signs of the Diseases of the Lungs and Pleura; illustrating their Pathology, and facilitating their Diagnosis. 2 plates. 8vo.
- WILLIAMS, S. W. Catechism of Medical Jurisprudence; being principally a Compendium of the opinions of the best Writers upon the subject. 18mo.
- American Medical Biography; or Memoirs of eminent Physicians who

- have died since the publication of Dr. Thacher's work on the same subject. 9 plates. 8vo.
- WILLIS, R. Urinary Diseases and their Treatment. 8vo.
- Illustrations of Cutaneous Disease : a series of Delineations of the Affections of the Skin in their most interesting and frequent forms, with a practical summary of their Symptoms, Diagnosis, and Treatment. 94 beautiful coloured plates. Folio.
- WILSON, ANDREW. Practical Observations on the Action of Morbid Sympathies ; as included in the Pathology of certain Diseases. 8vo.
- WILSON, CHARLES. Observations on Gout and Rheumatism, including an Account of a speedy, safe, and effectual Remedy for those Diseases. 8vo.
- WILSON, ERASMUS. A System of Human Anatomy, General and Special. Edited by P. B. Goddard, M.D. 170 engravings. 8vo.
- The Dissector, or Practical and Surgical Anatomy. Modified and re-arranged by P. B. Goddard, M.D. 106 engravings. 12mo.
- A Practical and Theoretical Treatise on the Diagnosis, Pathology, and Treatment of Diseases of the Skin, arranged according to a Natural System of Classification, and preceded by an Outline of the Anatomy and Physiology of the Skin. 8vo.
- WISTAR, C. A System of Anatomy, for the use of Students. With Additions by W. E. Horner, M.D. New edition, entirely remodelled and illustrated by more than 200 engravings, by J. Pancoast, M.D. 2 vols. 8vo.
- WOOD, G. B., AND F. BACHE. Dispensary of the United States of America. 8vo.
- WOODVILLE, W. Medical Botany : containing Systematic and General Descriptions, with Plates of all the Medicinal Plants, indigenous and exotic, comprehended in the Catalogues of the *Materia Medica*, as published by the Royal Colleges of London and Edinburgh, &c. The Botanical Descriptions arranged and corrected by Dr. W. J. Hooker, F.R.S., and the New Medico-Botanical portion supplied by G. Spratt, Esq. 310 coloured plates. 5 vols. 4to.
- WORCESTER, N. Synopsis of the Symptoms, Diagnosis, and Treatment of the more common and important Diseases of the Skin. 60 coloured figures. 8vo.
- WRIGHT, W. A Few Minutes' Advice to Deaf Persons ; comprising also useful information for the Professional World. 18mo.
- On the Varieties of Deafness, and Diseases of the Ear, with proposed methods of relieving them. 8vo.
- WYATT, T. A Synopsis of Natural History : embracing the Natural History of Animals, with Human and General Animal Physiology, Botany, Vegetable Physiology and Geology. Translated from the French of C. Lemmonnier, with Additions from the Works of Cuvier, Dumaril, Lacepede, &c. 49 plates. 8vo.
- YEATS, G. D. A Statement of the Early Symptoms which lead to the Disease termed Water in the Brain ; with observations on the necessity of a watchful attention to them, and on the fatal consequences of their neglect. 8vo.
- YOUATT, W. The Horse, with numerous illustrations ; also a General History of the Horse, and an Essay on the Ass and the Mule, by J. S. Skinner. 8vo.
- Cattle ; their Breeds, Management, and Diseases. Numerous engravings. 8vo.

S. S. & W. W. have an extensive assortment of SCHOOL, THEOLOGICAL, AND MISCELLANEOUS BOOKS, STATIONERY, &c., at low prices, catalogues of which may be had on application at 261 PEARL STREET.

SUPPLEMENTARY CATALOGUE.

- ACCUM, F. Treatise on Adulterations of Food, and Culinary Poisons, with methods of detecting them. 18mo.
- ACTON, W. A Complete Practical Treatise on Venereal Diseases, and their immediate and remote consequences; with observations on certain affections of the Uterus attended with Discharges. 6 colored plates. 8vo.
- ADAIR, J. M. Philosophical and Medical Sketch of the Natural History of the Human Body and Mind; with an Essay on the Difficulties of attaining Medical Knowledge, &c. 8vo.
- ADAMS, W. Practical Observations on Ectropium, or Eversion of the Eye-Lids, with description of a new Operation for the cure of that disease; on the modes of forming an Artificial Pupil; and on Cataract. 3 plates. 8vo.
- ALIBERT, J. L. Clinique de l'Hopital Saint Louis, ou traité complet des maladies de la Peau, &c. 63 colored plates. Folio.
- ALLEN, —. The Heart and Arteries, including the Fœtal and Pulmonary Circulation. 10 colored plates.
- ANDRY, F. Manual of Diagnosis of Diseases of the Heart; preceded by Clinical Researches for the study of these affections: translated by Samuel Kneeland, Jr., M. D. 12mo.
- ARMSTRONG, JOHN. Morbid Anatomy of the Stomach and Bowels, with a summary of the Symptoms of the Acute and Chronic Affections of these organs. 16 colored plates. 4to.
- ARNOTT, J. Cases illustrative of the Treatment of Obstructions in the Urethra, &c., by the new instrument, the Dilator; with account of improvements of the method of Dissolving Stone by Injection, &c. Plate. 8vo.
- ARTHUR, R. A Popular Treatise on the Diseases of the Teeth; with description of their structure and modes of treatment: usual mode of inserting Artificial Teeth, &c. 28 illustrations. 12mo.
- BALLARD, E. and A. B. GARROD. Elements of Materia Medica and Therapeutics. With additions by R. E. Griffith, M. D. 8vo.
- BARD, SAMUEL. Domestic Narrative of the Life of Samuel Bard, M. D. LL. D., late President of the College of Physicians and Surgeons of the University of the State of New York, &c. By J. M'Vickar, A. M. 8vo.
- BARTH, M. and M. H. ROGER. A Manual of Auscultation and Percussion. Translated by F. G. Smith, M. D. 12mo.
- BATCHELDER, J. P. Thoughts on the Connection of Life, Mind, and Matter, in respect to Education. 8vo. stitched.
- BATEMAN, T. Account of the Contagious Fever of England, exemplified in the Epidemic prevailing in London in 1818; with modes of treatment, observations on the Nature and Properties of Contagion, &c. 8vo.
- BECK, LEWIS C. Adulterations of various substances used in Medicine and the Arts, with the means of detecting them; intended as a Manual for the Physician, the Apothecary, and the Artisan. 12mo.
— A Manual of Chemistry; containing a condensed view of the present state of the science, copious references to more extensive treatises, original papers, &c. 136 illustrations. 12mo.
- BELL, JOHN. Letters on Professional Character and Manners; on the Education of a Surgeon, and the Duties and Qualifications of a Physician: addressed to James Gregory, M. D., Prof. of the Practice of Medicine in the University of Edinburgh. 8vo.
- BELL, LUTHER V. An Attempt to investigate some obscure and undecided doctrines in relation to Small Pox, Varioloid, and Vaccination. 8vo.

- BEULLAE, J. P. *Memoir on Mutual Instruction, applied to the study of the elementary principles of Medicine.* 18mo. stitched.
- BLOXAM, W. *Surgical Anatomy of the Arteries.* 12 colored plates. Folio.
- BOERHAAVE'S Aphorisms concerning the Knowledge and Cure of Diseases. 8vo.
— Opera Omnia Medica. 4to.
- BOGUE, T. *A Treatise on the Structure, Color, and Preservation of the Human Hair.* 2 plates. 18mo.
- BOIVIN et DUGES. *Traité Pratique des Maladies de l'Uterus et de ses annexes, fondé sur un grand nombre d'observations cliniques.* 2 vols. 8vo., avec un Atlas de 41 planches in folio colorées, représentant les principales alterations morbides des organes genitaux de la femme.
- BOLTON, J. *A Treatise on Strabismus, with description of new instruments designed to improve the operation for its cure, in simplicity, ease, and safety.* Plate. 12mo.
- BOURGERY et JACOB. *Anatomie Elementaire en 20 tableaux, representant chacun un sujet dans son entier a la proportion de demi-nature, et quelquefois de grandeur naturelle.* Colored. Beautiful specimens of art: sold in sets or separately.
— *Traité Complet de l'Anatomie de l'Homme, comprenant la medecine operationaire, avec planches lithographiées d'apres nature.* Ouvrage divisé en quatre parties: anatomie descriptive; anatomie chirurgicale et operationaire; anatomie generale; et anatomie philosophique. 8 vols. in folio, publiés en 90 livraisons. Chaque livraison est composee de 3 ou 4 feuilles de texte et de 8 planches, avec leur explication en regard. Colored.
- This superb work, in course of publication in Paris, is now nearly completed, 75 Nos. being published, and will be imported to order. Specimen Nos. are in store for exhibition.
- BOWDITCH, HENRY I. *The Young Stethoscopist; or, the Student's Aid to Auscultation.* 42 illustrations. 18mo.
- BOYER, L. A. H. *Recherches sur l'operation du Strabisme.* 12 colored plates. 8vo.
- BRADFORD, D. *The Wonders of the Heavens: being a Popular View of Astronomy, including a full illustration of the Mechanism of the Heavens; embracing the Sun, Moon, and Stars, with descriptions of the Planets, Comets, Fixed Stars, Double Stars, the Constellations, the Galaxy or Milky Way, the Zodiacal Light, Aurora Borealis or Northern Lights, Meteors, Clouds, Falling Stars, Aerolites, &c.* Numerous maps and engravings. 4to.
- BRESCHET, G. *Memoires Chirurgicaux sur differentes especes d'Aneurysmes.* 6 plates. 4to.
— *Essai sur les Veines du Rachis: Recherches historiques et experimentales sur la formation du Cal: Considerations et observations Anatomiques et Pathologiques sur la Hernie Femorale, ou Merocce: de la Dessiccation et des autres moyens de conservation de Pieces Anatomiques.* 3 plates. 4to.
- BRIDGEMAN, T. *The Florist's Guide; containing practical directions for the cultivation of Annual, Biennial, and Perennial Flowering Plants, &c., with instructions for the management of Greenhouse Plants throughout the year.* 12mo.
— *The Kitchen Gardener's Instructor: containing practical directions for the cultivation of Culinary Vegetables and Herbs, &c.* 12mo.
- BRODIE, SIR B. C. *Clinical Lectures on Surgery.* 8vo.
- BROWNE, D. J. *The Trees of America, Native and Foreign, pictorially and botanically delineated, and scientifically and popularly described; being considered principally with reference to their geography and history, soil and situation, propagation and culture, accidents and diseases, properties and uses, economy in the arts, introduction into commerce, and their application in useful and ornamental plantations.* Numerous engravings. 8vo.
- BRYANT, T. S. *Examinations in Anatomy and Physiology.* 12mo.
- BUDD, G. *On Diseases of the Liver.* 2 colored plates and 20 wood engravings. 8vo.
- BUDD, R. H. *A Practical Treatise on the Diseases of the Foot of the Horse, their nature, causes, and prevention; also Rules for Shoeing, &c.* 8vo.

- BURKE, W. The Mineral Springs of Western Virginia, with remarks on their use, and the diseases to which they are applicable; with a Notice of the Fauquier White Sulphur Spring, &c. Map. 18mo.
- CAMPERI PETRI Icones Herniarum. Editæ a S. T. Soemmering. 14 plates. Folio.
- CARPENTER, W. B. Elements of Physiology, including Pathological Anatomy, for the use of the Medical Student. 180 illustrations. 8vo.
- CAZENAVE, A. Traite des Syphilides, ou Maladies Veneriennes de la Peau. 8vo. with folio atlas of 12 colored plates.
- CAZENAVE and SCHEDEL. Manual of Diseases of the Skin; with Notes and additions by T. H. Burgess, M. D. Revised and corrected, with additional Notes, by H. D. Bulkley, M. D. 12mo.
- CHAPMAN, H. A Brief Description of Surgical Apparatus, intended to accompany a series of delineations of the most important mechanical auxiliaries of Surgery. 8vo. with 4to. Atlas of 20 plates.
- CHAPMAN, N. A Compendium of Lectures on the Theory and Practice of Medicine, delivered by Professor Chapman in the University of Pennsylvania. Prepared by N. D. Benedict, M. D. 8vo.
- CHEYNE, J. An Essay on Cynanche Trachealis, or Croup. 12mo.
- An Essay on Hydrocephalus Acutus, or Dropsy in the Brain. 12mo.
- CHURCHILL, F. Observations on the Diseases incident to Pregnancy and Childbed. 8vo.
- CLEOBUREY, W. A Review of the different Operations performed on the Eyes, for the restoration of lost, and the improvement of imperfect Vision; with full account of the various structures and diseases of the Eyes and their appendages, mode of treatment, &c. 8vo.
- CLOQUET, H. A System of Human Anatomy; with Notes, &c., by R. Knox, M. D. 8vo.
- CLYMER, M. Fevers; their Diagnosis, Pathology, and Treatment. Prepared and edited, with large additions, from the Essays on Fever in Tweedie's Library of Practical Medicine. 8vo.
- COCK, M. R. The American Poultry Book; being a practical treatise on the management of Domestic Poultry. 18mo.
- COLEY, H. A Treatise on Poisons and Asphyxia in relation to Medical Jurisprudence. 8vo.
- COLEY, J. M. A Practical Treatise on the Diseases of Children. 8vo.
- COOPER, T. Tracts on Medical Jurisprudence, including Farr's Elements of Medical Jurisprudence, Dease's Remarks on Medical Jurisprudence, Male's Epitome of Juridical or Forensic Medicine, and Haslam's Treatise on Insanity; with Notes, and a Digest of the Law relating to Insanity and Nuisance. 8vo.
- COX, T. Observations on Acute Rheumatism, and its Metastasis to the Heart, &c. 8vo.
- COXE, J. R. The American Dispensatory, containing the Natural, Chemical, Pharmaceutical, and Medical History of the different substances employed in Medicine, &c. &c. 8vo.
- CRICHTON, A. Practical Observations on the Treatment and Cure of several varieties of Pulmonary Consumption, and on the effects of the Vapor of Boiling Tar on that disease. 8vo.
- CROWTHER, B. Practical Observations on the Disease of the Joints, commonly called White Swelling; with remarks on Caries, Necrosis, and Scrofulous Abscess, &c. 7 colored plates. 8vo.
- CRUIKSHANK, W. The Anatomy of the Absorbing Vessels of the Human Body. 5 plates. 4to.
- CRUVEILHIER, J. Anatomie Pathologique du Corps Humain, ou descriptions avec figures lithographiees et coloriees des diverses alterations morbides dont le corps humain est susceptible. 233 colored plates. 2 vols. folio.
- CURTIS, J. H. Observations on the Preservation of Hearing, and on the choice, use, and abuse of Hearing Trumets &c. 18mo.

- CUTTER, C. Anatomy and Physiology: designed for Academies and Families. 215 engravings. 12mo.
- DARBY, J. A Manual of Botany, adapted to the productions of the Southern States. Arranged on the Natural System, preceded by an Analysis. 145 engravings. 12mo.
- DAVIES, T. Lectures on the Diseases of the Lungs and Heart. 8vo.
- DELPECH, J. Chirurgie Clinique de Montpellier, ou observations et reflexions tirees destravaux de chirurgie clinique de cette ecole. 35 plates. 2 vols. 4to.
- Memorial des Hopitaux du Midi, et de la Clinique de Montpellier. 8 plates. 2 vols. 4to.
- De l'Orthomorphie, par rapport a l'Espece Humaine: ou recherches anatomico-pathologique sur les causes, les moyens de prevenir, ceux deguerir les principales difformites et sur les veritables fondemens de l'art appele orthopedique. 2 vols. 8vo., with 4to. atlas of 78 plates.
- DENDY, W. C. The Philosophy of Mystery. 12mo.
- DIXON, E. H. A Treatise on Diseases of the Sexual Organs; adapted to popular and professional reading, and the exposition of Quackery, professional and otherwise. 12mo.
- DONALDSON, P. A Review of the present Systems of Medicine and Chirurgery of Europe and America, &c. Portrait. 8vo.
- DRAPER, J. W. A Text-Book of Chemistry, for the use of Schools and Colleges. 272 engravings. 12mo.
- DURLACHER, L. A Treatise on Corns, Bunions, the Diseases of Nails, and the general management of the Feet. 12mo.
- ECCLLES, W. Observations on the Ulcerative Process and its Treatment, particularly when affecting the Leg. 8vo.
- EPPS, J. Domestic Homeopathy; or, Rules for the Domestic Treatment of the Maladies of Infants, Children and Adults, and for the conduct and the treatment during Pregnancy, Confinement, and Suckling. 18mo.
- Essays on Practical Medicine and Surgery, by Drs. Chapman, Bache, Coates, Geddings, Griffith, Mitchell, Wood, Dunglison, Condic, Dewees, Hays, Jackson, Paterson, Hodge, Horner, &c. 2 vols. 8vo.
- FALKNER, F. The Farmer's Manual: a Practical Treatise on the Nature and Value of Manures, founded from experiments on various crops, &c. 12mo.
- FELTHAM, J. A Popular View of the Structure and Economy of the Human Body; with reflections, moral, practical, and miscellaneous; including modern discoveries, &c. 12mo.
- FISHER, G. T. Photogenic Manipulation; containing the Theory and Plain Instructions in the Art of Photography, or the production of Pictures through the agency of Light: including Calotype, Chrysotype, Cyanotype, Chronotype, Energiatype, Anthotype, and Amphitype. 14 engravings. 18mo.
- FLETCHER, R. Medico-Chirurgical Notes and Illustrations, on some dangerous affections of the Throat, which induce sudden death by suffocation,—on Strictures of the Oesophagus, and the dangers of the Bougie,—on the cure of the falling down of the Bowel in grown persons,—Anomalies in Rupture Operations, &c. &c. 4 plates. 4to.
- FLOURENS, P. Phrenology Examined. Translated by C. D. Meigs, M. D. 18mo.
- FLOYER, J. The Physician's Pulse-Watch; or, an Essay to explain the old art of feeling the Pulse, and to improve it by the help of a Pulse-Watch. 12mo. 1707.
- FORBES, JOHN. Homeopathy, Allopathy, and "Young Physic." 18mo.
- FOWLER, O. S. Practical Phrenology; giving a concise elementary view of Phrenology, presenting some new and important remarks upon the temperament, and describing the primary mental powers in seven different degrees of development, and the location of the organs. Numerous engravings. 12mo.
- FOX, JOSEPH. The Natural History and Diseases of the Human Teeth. Remodelled, with an Introduction and numerous Additions, by Chapin A. Harris, M. D., D. D. S. 30 plates. 8vo.

- FYFE, A. A Compendium of Anatomy, Human and Comparative. 40 plates. 4 vols. 8vo.
- GARDENER and FLORIST. A Select Manual of Kitchen Gardening, and culture of Fruits; containing familiar directions for the most approved practice in each department, descriptions of many valuable fruits, and a Calendar of work to be performed each month in the year. Also, a Manual of Gardening, containing practical instruction for the management of Greenhouse Plants, and for the cultivation of the Shrubby, the Flower-Garden, and the Lawn, &c. &c. 12mo.
- GARDINER, J. An Inquiry into the Nature, Cause, and Cure of the Gout, and of some of the Diseases with which it is connected. 12mo.
- GERRISH, A. A Synopsis of the Prevention and Cure of Disease. 12mo.
- GIBNEY, J. Practical Observations on the Use and Abuse of Cold and Warm Sea-Bathing, in various diseases, particularly in Scrofulous and Gouty cases. 8vo.
- GLANDORPH (MATTHIÆ) Bremensis Medici Clarissimi Opera Omnia: viz. Speculum Chirurgorum, Methodus Medendæ Paronychiæ, Tractatus de Polypo Narium affectu gravissimo, Gazophylacium Polyplusium Fonticularum et Setonum reseratum, &c. 4to. 1729.
- GOOCH, B. Cases and Practical Remarks in Surgery, with sketches of Machines of simple construction, easy application, and approved use. 12 plates. 8vo. 1758.
- GOVE, MARY S. Lectures to Women on Anatomy and Physiology: with an Appendix on the Water Cure. 12mo.
- GRANVILLE, A. B. Graphic Illustrations of Abortion and the Diseases of Menstruation. 14 colored plates. 4to.
- GRIFFIN, W. and D. Medical and Physiological Problems, being chiefly researches for correct principles of treatment in disputed points of Medical Practice. 8vo.
- GRIFFITH, J. W. A Practical Manual, containing a description of the General, Chemical, and Microscopical characters of the Blood, and Secretions of the Human Body, as well as their components, &c. &c. 12mo.
- GRIFFITHS, T. Chemistry of the Four Seasons; Spring, Summer, Autumn, Winter. With engravings.
- GUENON, F. A Treatise on Milch Cows, whereby the Quality and Quantity of Milk which any Cow will give may be accurately determined by observing natural marks or external indications alone; the length of time she will give milk, &c. With remarks and observations on the Cow and the Dairy, by J. S. Skinner. Many engravings. 8vo.
- GULLY, J. M. The Water Cure in Chronic Diseases: an exposition of the causes, progress, and terminations of various Chronic Diseases of the Digestive Organs, Lungs, Nerves, Limbs, and Skin; and of their treatment by Water and other Hygienic means. 12mo.
- GUTHRIE, G. J. On some points connected with the Anatomy and Surgery of Inguinal and Femoral Herniæ. 3 plates. 4to.
- HAHNEMANN, S. *Materia Medica Pura*. Translated and edited by C. J. Hempel, M.D. 4 vols. 8vo.
- HARE, S. Practical Observations on the Prevention, Causes, and Treatment of Curvatures of the Spine, with a description of an apparatus for the correction of the deformity. 11 plates. 8vo.
- HASLAM, J. Observations on Madness and Melancholy; including practical remarks on those diseases, together with Cases, Morbid appearances on dissection, &c. 8vo.
- Sound Mind; or, Contributions to the Natural History and Physiology of the Human Intellect. 8vo.
- HASSALL, A. H. The Microscopic Anatomy of the Human Body in Health and Disease, with colored plates.
- HAYES, T. A Serious Address on the dangerous consequences of neglecting common Coughs and Colds, with ample directions for the prevention and cure of Consumption; with observations on Hooping Cough and Asthma. 12mo.
- HEMPEL, C. J. The Homœopathic Domestic Physician. 18mo.

- HENDERSON, W. An Inquiry into the Homœopathic Practice of Medicine. 8vo.
- HERING, C. Domestic Physician; comprising the former edition of the Homœopæthist, or Domestic Physician, with the additions of Drs. Goullon, Gross, and Stapf, and a chapter on Diseases of Women. Portrait. 12mo.
- HEWSON, T. Observations on the History and Treatment of the Ophthalmia accompanying the secondary forms of *Lucas Venerea*. Colored plate. 8vo.
- HEY, W. Practical Observations in Surgery, illustrated with cases and 10 plates. 8vo.
- HIND, G. W. A series of Twenty Plates illustrating the causes of Displacement in the various Fractures of the Bones of the Extremities. Large 4to.
- HIPPOCRATIS *coi Prolegomena et Prognosticorum libri tres: cum paraphrastica versione et brevibus commentariis Johannis Hevrnii Ultraiectini*. Small 4to.
- HIPPOCRATES and GALEN, the Writings of, epitomised from the original Latin translations by J. R. Coxe, M. D. 8vo.
- HODGKIN, T. Lectures on the Morbid Anatomy of the Serous and Mucous Membranes. 8vo.
- HOOD, P. Practical Observations on the Diseases most fatal to Children; with reference to the propriety of treating them as proceeding from Irritation, and not from Inflammation. 8vo.
- HOOPER, R. Examinations in Anatomy, Physiology, Practice of Physic, Surgery, Chemistry, Materia Medica, and Pharmacy: for the use of Students. 18mo.
- HORNER, W. E. The United States Dissector; or, Lessons in Practical Anatomy. Edited by H. H. Smith, M. D. 72 engravings. 12mo.
- HUMBERT et JACQUIER. *Essai et Observations sur la maniere de reduire les Luxations spontanees ou symptomatiques de l'articulation ilio-femorale; methode applicable aux luxations congenitales et aux luxations anciennes par cause externe*. 8vo., with 4to. atlas of 29 plates.
- HUTCHISON, A. C. Practical Observations in Surgery, illustrated by cases and various official documents. Plate. 8vo.
- HUXHAM, J. An Essay on Fevers; with a Dissertation on the Malignant Ulcerous Sore Throat. 8vo.
- INGPEN, A. Instructions for Collecting, Rearing, and Preserving British and Foreign Insects; also for collecting and preserving Crustacea and Shells. 3 colored plates. 18mo.
- Intellectual and Moral Qualities, Facts and Arguments on the Transmission of, from Parents to Offspring. 12mo.
- JACKSON, S. H. Cautions to Women respecting the state of Pregnancy, the progress of Labor and Delivery, the confinement of Child-bed, and some constitutional diseases, with directions to Midwives and Nurses, &c. &c. 12mo.
- JAHR, G. H. G. Short Elementary Treatise upon Homœopathia, and the manner of its Practice; with some of the most important effects of Ten of the principal Homœopathic Remedies. Translated by E. Bayard, M. D. 18mo.
- JAMES, J. H. Observations on the General Principles, and on the Particular Nature and Treatment of various species of Inflammation. 8vo.
- JOHNSON, E. Life, Health, and Disease. 12mo.
- Results of Hydropathy; or, Constipation not a Disease of the Bowels,—Indigestion not a Disease of the Stomach; with an Exposition of the True Nature of these ailments, explaining the reason why they are so certainly cured by the Hydropathic treatment, &c. 12mo.
- KENTISH, E. An Essay on Burns, principally upon those which happen to Workmen in Mines from the explosions of inflammable air, &c. 8vo.
- KINGLAKE, R. A Dissertation on Gout, exhibiting a new view of the origin, nature, cause, cure, and prevention of that afflicting disease, illustrated by cases. 8vo.
- KIRBY, W. and W. SPENCE. An Introduction to Entomology; or, Elements of the Natural History of Insects: comprising an account of Noxious and Useful Insects, of their Metamorphoses, Food, Stratagems, Habitations, Societies, Motions, Noises, Hybernation, Instinct, &c. &c. 5 plates. 8vo.

- KOECKER, L. Principles of Dental Surgery; exhibiting a new method of treating the Diseases of the Teeth and Gums, especially calculated to promote their health and beauty; with a general view of the present state of Dental Surgery, &c. 8vo.
- LANGENBECK, C. J. M. Plates of the Peritoneum. 12 plates. Folio.
- LARDNER, D. Popular Lectures on Science and Art; delivered in the principal cities and towns of the United States. Numerous engravings. 2 vols. 8vo.
- LATHAM, P. M. Lectures on subjects connected with Clinical Medicine; comprising Diseases of the Heart.
- LE DRAN. Operations in Surgery. Translated by M. Gataker. 21 plates. 8vo. 1757.
- LEGER, T. Animal Magnetism; or, Psychodunamy. 12mo.
- LEV, H. An Essay on the Laryngismus Stridulus, or Croup-like Inspiration of Infants; with illustrations of the general principles of the Pathology of the Nerves, and of the functions and diseases of the Par Vagum and its principal branches. 5 plates. 8vo.
- LIND, J. An Essay on the most effectual means of preserving the Health of Seamen in the Royal Navy. Also a Dissertation on Fevers and Infection, Jail Distemper, &c. 8vo.
- LION, H. A complete Treatise upon Spinæ Pedum, containing several important discoveries. 5 plates. 8vo.
- LOBSTEIN, J. F. D. A General Guide for Practising Physicians, in the Examination of the Sick. 18mo.
- LOBSTEIN, J. F. Traite d'Anatomie Pathologique. 2 vols. 8vo., with folio Atlas of 26 colored plates.
- MACKENZIE, W. The Physiology of Vision. 8vo.
- MACNISH, R. The Anatomy of Drunkenness. 12mo.
- MANEC, P. J. A Theoretical and Practical Treatise upon the Ligature of Arteries. Translated by J. W. Garlick, M.R.C.S., and W. C. Copperthwaite, M.R.C.S. 14 colored plates. 4to.
- MASSE, J. N. A Pocket Atlas of the Descriptive Anatomy of the Human Body. Edited by G. S. Pattison, M.D. 112 plates. 12mo.
- MEAD, R. Medical Works; containing a mechanical account of Poisons,—influence of the Sun and Moon upon Human Bodies,—on the Plague,—on Small-Pox and Measles,—on extracting Foul Air from Ships,—on Scurvy,—Medical Precepts and Cautions,—Medica Sacra,—on Worms of Human Bodies,—Hydrophobia, &c. &c. Plates. 3 vols. 18mo.
- Medecins et Chirurgiens Celebres. 57 portraits, finely executed on stone. 4to.
- Medical and Surgical Monographs, by Drs. Andral, Babington, Beek, Bright, Brodie, Burne, Carmichael, Clutterbuck, Cormack, Dubois, Farr, Itard, Louis, Maunoir, and Taylor. 8vo.
- Medical Faculty of the University of New York, and Founders of the Medical Department: comprising the Portraits of Chancellor Frelinghuysen, and Professors Mott, Pattison, Revere, Paine, Bedford, and Draper, on one sheet 30 by 24 inches.
- of the College of Physicians and Surgeons of the University of the State of New York: comprising the Portraits of Professors Stevens, Smith, Beck, Torrey, Watts, Parker, and Gilman, on one sheet 30 by 24 inches.
- MILLER, J. The Practice of Surgery. 8vo.
- MONRO, A. Traite d'Osteologie, traduit de l'Anglois de M. Monro, ou l'on a ajoute des planches en taille-douce, que representent au naturel tous les Os de l'Adulte et du Fœtus, avec leurs explications. 32 plates. 2 vols. folio.
- Observations on Crural Hernia; with a general account of the other varieties of Hernia. 5 plates. 8vo.
- Engravings of the Thoracic and Abdominal Viscera, and the Canals connected with them; representing the natural appearance of those important parts immediately after death, and without being affected by previous disease. 5 plates. 4to.

- MURPHY, E. W. Lectures on Natural and Difficult Parturition. 70 Engravings. 8vo.
- MURRAY, J. A Treatise on Pulmonary Consumption; its prevention and remedy. 8vo.
- A System of Materia Medica and Pharmacy; including translations of the Edinburgh, London, and Dublin Pharmacopœias. With Notes and Additions by John B. Beck, M.D. 8vo.
- NEALE, A. Researches respecting the Natural History, Chemical Analysis, and Medicinal Virtues of the Spur, or Ergot of Rye, when administered as a remedy in certain states of the Uterus. Colored plate. 8vo.
- NICHOL, J. P. The Phenomena and Order of the Solar System. 21 plates. 12mo.
- Views of the Architecture of the Heavens. 23 plates. 12mo.
- Nurse's Manual and Mother's Medical Adviser; a Guide to the Inexperienced. By a Practising Physician. 12mo.
- OWEN, R. Lectures on the Comparative Anatomy and Physiology of the Invertebrate Animals. 138 engravings. 8vo.
- Odontography; or, a Treatise on the Comparative Anatomy of the Teeth, their physiological relations, mode of development, and microscopic structure, in the Vertebrate Animals. 150 plates. 2 vols. 8vo.
- PALLETTA, J. B. Exercitationes Pathologicae. 12 plates. 4to.
- PALMER, S. A Pentaglot Dictionary of the Terms employed in Anatomy, Physiology, Pathology, Practical Medicine, Surgery, Obstetrics, Medical Jurisprudence, Materia Medica, Pharmacy, Medical Zoology, Botany, and Chemistry. Part I., with the leading term in French, followed by the synonymes in the Greek, Latin, German, and English; explanations in English. Part II. A German-English-French Dictionary, comprehending the Scientific German Terms of the preceding part. 8vo.
- PARSONS, U. Physician for Ships; containing Medical Advice for Seamen and other persons at Sea, on the treatment of diseases, and on the preservation of health in sickly climates. 12mo.
- PETZOLDT, A. Lectures to Farmers on Agricultural Chemistry. 8vo.
- PHILIP, A. P. W. A Treatise on the Means of Preserving Health; and particularly the prevention of Organic Diseases. 8vo.
- PHILLIPS, B. Scrofula; its nature, its causes, its prevalence, and the principles of treatment. Plate. 8vo.
- Physic and Physicians: a Medical Sketch-Book, exhibiting the Public and Private Life of the most celebrated Medical Men of former days; with Memoirs of eminent living London Physicians and Surgeons. 2 vols. 18mo.
- POLE, T. The Anatomical Instructor; or, an illustration of the modern and most approved methods of preparing and preserving the different parts of the Human Body, and of Quadrupeds, &c. 10 plates. 8vo.
- PROSSER, T. A Treatise on the Strangles and Fevers of Horses. 8vo.
- PURSH, F. Flora Americae Septentrionalis; or, a systematic arrangement and description of the Plants of North America. 24 colored plates. 2 vols. 8vo.
- RANDALL, S. S. Incentives to the Cultivation of the Science of Geology; designed for the use of the Young. Many engravings. 18mo.
- REES, G. O. On the Analysis of the Blood and Urine, in Health and Disease; and on the treatment of Urinary diseases. Plate. 8vo.
- REID, T. Directions for Warm and Cold Sea-Bathing; with observations on their application and effects in different diseases. 8vo.
- RENWICK, J. First Principles of Chemistry; being a familiar introduction to the study of that science. 154 illustrations. 18mo.
- REYBEARD, J. F. Memoires sur le traitement des Anus Artificiels, des plaies des intestins, et des plaies penetrantes de poitrine. 3 plates. 8vo.
- RICHARDSON, G. F. Geology for Beginners. 261 engravings. 12mo.
- RICORD, P. Clinique Iconographique de l'Hopital des Veneriens, recueil d'observations, suivies de considerations pratique sur les Maladies qui ont ete traites dans cet Hopital. *In course of publication—to contain about 50 colored plates.* 4to.

- RIGBY, E. Memoranda for Practitioners in Midwifery. With Additions by S. C. Foster, M. D. 32mo.
- RIVERS, H. W. Accidents: Popular Directions for their immediate treatment; with observations on Poisons and their Antidotes. 4 plates. 12mo.
- ROBINSON, J. The Surgical, Mechanical, and Medical treatment of the Teeth: including Dental Mechanics. 139 engravings. 12mo.
- RUSN, J. The Philosophy of the Human Voice, embracing its Physiological History; with a system of principles by which Criticism in the art of Elocution may be rendered intelligible, and instruction definite and comprehensive. 8vo.
- SAUNDERS, E. Advice on the care of the Teeth. Colored plate. 18mo.
- SCARPA, A. Sull 'Ernie Memorie Anatomico-Chirurgiche. 11 plates. Folio.
- SCOUTETTEN, H. La Methode Ovalaire, ou nouvelle methode pour Amputer dans les Articulations. 11 plates. 4to.
- SERRE, M. Traite de la Reunion Immediate, et de son influence sur les progres recens de la Chirurgie dans toutes les operations, &c. 3 plates. 8vo.
- SERRES, M. Recherches d'Anatomie Transcendante et Pathologique: theorie des Formations et des Deformations Organique, appliquee a l'Anatomie de Ritta-Christina, et de la duplice monstrueuse. 4to., with folio atlas of 20 plates.
- SEWALL, T. An Examination of Phrenology. 8 plates. 8vo.
- SHEARMAN, W. Observations illustrative of the History and Treatment of Chronic Debility; the prolific source of Indigestion, Spasmodic Diseases, and various Nervous Affections. 8vo.
- SHEPARD, C. U. A Treatise on Mineralogy. 500 illustrations. 12mo.
- SHERILL, H. An Essay on Epidemics, as they appeared in Dutchess County, N. Y., from 1809 to 1825: also on Diseases of the Jaw-Bones, Epidemic Cholera in Poughkeepsie, 1832, &c. 8vo.
- SHIPMAN, G. A Familiar Treatise on Disorders of the Stomach and Bowels, Bilious and Nervous Affections, &c. 8vo.
- SIMON, J. F. Animal Chemistry with reference to the Physiology and Pathology of Man. Edited by G. E. Day, M. A. and L. M. 3 plates. 8vo.
- SKINNER, J. S. The Dog and the Sportsman; embracing the Uses, Breeding, Training, Diseases, &c., of Dogs, and an account of the different kinds of Game, with their habits, &c. 5 plates, 12mo.
- The Monthly Journal of Agriculture, containing the best current productions in promotion of Agricultural improvement; including the choicest Prize Essays issued in Europe and America: with original contributions from eminent Farmers and Statesmen. Numerous engravings. 8vo.
- SMITH, JOSEPH A. Productive Farming; or, a familiar digest of the recent discoveries of Liebig, Johnston, Davy, and other celebrated writers on Vegetable Chemistry; showing how the results of Tillage might be greatly augmented. 12mo.
- SMITH, JOSEPH M. Elements of the Etiology and Philosophy of Epidemics. 8vo.
- SOLLY, S. The Human Brain, its Configuration, Structure, Development, and Physiology; illustrated by references to the Nervous System in the lower orders of Animals. 12 plates. 8vo.
- SPRATT, H. G. Obstetric Tables; comprising Graphic Illustrations, with descriptions and practical remarks, exhibiting, on dissected plates, many important subjects in Midwifery. 15 colored plates. 4to.
- STOKER, W. A Treatise on Fever, with observations on the practice adopted for its cure in the Fever Hospital and House of Recovery in Dublin: illustrated by cases. 8vo.
- SUE, M. Elemens d'Anatomie, a l'usage des Peintres, des Sculpteurs, et des Amateurs. 14 plates. 4to.
- SYER, J. A Treatise on the Management of Infants; containing the general principles of their domestic treatment, with history and method of cure of some of their most prevalent and formidable diseases. 8vo.
- SYME, J. Treatise on the Excision of Diseased Joints; with case of Osteo-Sarcoma of the Lower Jaw. 6 plates. 8vo.

- TAMPLIN, R. W. Lectures on the Nature and Treatment of Deformities. 64 illustrations. 12mo.
- THAER, A. D. The Principles of Agriculture. Translated by W. Shaw and C. W. Johnson, F.R.S. 8vo.
- THOMAS, J. J. The Fruit Culturist, adapted to the climate of the Northern States; containing directions for raising Young Trees in the Nursery, and for the management of the Orchard and Fruit Garden. 36 engravings. 18mo.
- THOMSON, J. An Account of the Varioloid Epidemic in Edinburgh and other parts of Scotland; with observations on the Identity of Chicken-Pox with Modified Small-Pox. 8vo.
- TODD, R. B. The Descriptive and Physiological Anatomy of the Brain, Spinal Cord, and Ganglions, and of their coverings. 38 engravings. 8vo.
- TRAVERS, B. An Inquiry into the Process of Nature in repairing Injuries of the Intestines; illustrating the treatment of Penetrating Wounds, and Strangulated Hernia. 7 plates. 8vo.
- The Physiology of Inflammation, and the Healing Process. 8vo.
- Observations on the Pathology of Venereal Affections. 8vo.
- TWEEDIE, A. Dissertations on Fevers, General Pathology, Inflammation, and Diseases of the Skin, by Drs. Symonds, Allison, Christison, Schedel, Locock, Gregory, Burrows, and Shapter. Edited by A. Tweedie, M. D.
- TYRRELL, F. A Practical Work on the Diseases of the Eye, and their treatment, Medically, Topically, and by Operations. 8 colored plates. 8vo.
- VAN SWIETEN's Commentaries Abridged, by Dr. Schomberg. 8vo.
- VAUGHAN, W. An Exposition of the Principles of Anatomy and Physiology; containing the Prælectiones Anatomicæ of F. Leber. 2 vols. 8vo.
- WADD, W. Observations on the best mode of relieving Strictures in the Urethra; with remarks on the frequent inefficacy and ill effects of Caustic applications. 8vo.
- Cases of Diseased Bladder and Testicle. 21 plates. 4to.
- WALKER, G. A. Gatherings from Graveyards, particularly those of London; with a concise history of the modes of interment among different nations, from the earliest periods, and a detail of dangerous and fatal results produced by the unwise and revolting custom of inhuming the Dead in the midst of the Living. 8vo.
- WALKER, C. V. Electrotypic Manipulation: being the theory and plain instructions in the art of Working in Metals, by precipitating them from their Solutions, through the agency of Galvanic or Voltaic Electricity. Engravings. 18mo.
- WALLER, J. A. A Treatise on Burns and Scalds; detailing the best methods of treating those accidents, as practised in the London Hospitals, &c. 24mo.
- WARREN, J. C. Physical Education, and the Preservation of Health. 18mo.
- WELLS, W. C. An Essay upon Single Vision with Two Eyes; also on Dew: account of a White Female, part of whose skin resembles that of a Negro, with observations on the causes of differences in color, &c. 8vo.
- WHITE, C. A Treatise on the management of Pregnant and Lying-in Women, and the means of curing and preventing the principal disorders to which they are liable, &c. &c. 2 plates. 8vo.
- WILLARD, E. On the Motive Powers which produce Circulation of the Blood. 12mo.
- WILSON, E. A Practical Treatise on Healthy Skin; with rules for the Medical and Domestic Treatment of Cutaneous Diseases. 6 plates. 12mo.
- WINSLOW, F. The Plea of Insanity in Criminal cases. 12mo.
- WOOD, A. A Class-Book of Botany, designed for Colleges, Academies, &c.; containing the Elements of Botanical Science; the Natural Orders, illustrated by a Flora of the Northern States, &c. 38 engravings. 12mo.
- WORMALD, T. and A. M. McWHINNIE. A Series of Anatomical Sketches and Diagrams, with descriptions and references. 28 plates. 4to.
- YOUNG, S. Minutes of Cases of Cancer and Cancerous tendency, successfully treated by Samuel Young, Surgeon. 8vo.
- BELL, BENJAMIN. A System of Surgery. 99 plates. 4 vols. 8vo.

JOHNSON'S
MEDICO-CHIRURGICAL REVIEW,
AND
JOURNAL OF PRACTICAL MEDICINE.
(PUBLISHED QUARTERLY.)

ARRANGEMENT.

- 1.—Extensive Analytical Reviews of English and Foreign Medical Works.
- 2.—Quarterly Periscope of Practical Medicine; or Spirit of the Medical Journals, Foreign and Domestic
- 3.—Periscope of Authenticated Hospital Reports, British and Continental, with Commentaries.
- 4.—Analecta; or, Gleanings from the Periodical Presses of Europe and America.
- 5.—Prize Hospital Reports, Metropolitan and Provincial.

The whole forming two large annual volumes of concentrated practical information in

MEDICINE, SURGERY, MIDWIFERY, PHARMACY, ETC

REPUBLISHED

BY RICHARD AND GEORGE S. WOOD,

No. 261 PEARL STREET, NEW YORK.

Price, Five Dollars Per Annum, Payable in Advance.

A Work devoted exclusively to the interests of any one Profession, if it possesses merit, strongly recommends itself to every member of that profession; and as most of the improvements and discoveries in Medicine and Surgery are communicated through the medium of the Periodical Press, works of this kind are nearly as indispensable to the Physician or Surgeon, as a hammer to the Artist. None, therefore, who are not indifferent to the interests of their profession, or who wish to keep pace with the progress of knowledge in their own department, can, without disadvantage, be without such a work. This Journal having been reprinted and extensively circulated in this country for more than twenty years, is too well known to make it necessary to enlarge much upon its merits. The acknowledged talents of the author—the ability, the impartiality, and the justness of its notices of new publications; which, forming in general, a complete abstract or synopsis of the work reviewed, supersede, in a measure, the necessity of purchasing the works themselves—the wide range taken in collecting its reports of cases, which are gleaned from the extensive hospitals of Great Britain, Ireland, and the continent of Europe; offering to the young practitioner the advantages of the most extensive practice—and its selections from American Journals, by which it combines in some degree the advantages of a foreign and domestic journal, give it a claim to patronage possessed perhaps by no other similar work. In confirmation of this opinion, the following testimonials, from gentlemen of the highest respectability in the profession are submitted.

RECOMMENDATIONS.

Sir,—I consider the Medico-Chirurgical Review among the most important additions to the Medical Literature of the day. I have no doubt it has contributed very much to the improvement of Medicine, both in Europe and in America. It certainly will not suffer by a comparison with any similar work in either hemisphere; and I am happy to learn that you are engaged in continuing the publication of it in this country. Wishing you every success in this enterprise, I am, very respectfully, your obedient servant.

JOHN WATTS, Jr., Pres. of Coll. of Phys. and Surg., N. Y.

I consider the Medico-Chirurgical Journal as decidedly the most valuable periodical Medical publication we receive from Great Britain. The republication of it in this country, which I am happy to learn you intend to continue, cannot fail to advance the science of medicine.

ALEX. H. STEVENS, M. D. Prof. of Surg.

I entirely concur in the above opinions respecting the character of the Medico-Chirurgical Review, believing there is no Medical Journal superior to it in point of erudition and science on either side of the Atlantic.

JOSEPH M. SMITH, M. D. Prof. of Theo. and Prac. of Phys.

I have been in the habit of reading the Medico-Chirurgical Review from its first publication in this country, and have derived more advantage from its perusal, than any other Medical Journal with which I am acquainted.

EDW. DELAFIELD, M. D. Prof. of Obstet., &c.

The public opinion has been so decidedly and favourably expressed in relation to the

merits of Dr. Johnson's Journal as to render almost unnecessary any additional testimony in favour of the spirit and ability with which it is conducted. Devoted as this Journal is to the analysis of works to which a great proportion of the Medical Profession in this country are necessarily denied access, I have always looked upon its republication here, as a substantial service rendered to the cause of Medical Literature in the United States; and I cannot therefore but wish the enterprising publisher every success in his endeavours to give it an extended circulation.

JOHN B. BECK, M. D. Prof. of Mat. Med.

The following Gentlemen have expressed their concurrence with the foregoing:

J. AUG. SMITH, M. D. Prof. of Anat. and Surg., J. D. JAKES, M. D., Treas. and Trustee of Coll. of Phys. and Surg., JAS. FREEMAN DANA, Prof. of Chemistry; J. R. RHINELANDER, M. D., Dem. of Anat.; N. H. DERING, Regist.

I state with great readiness, the high opinion I entertain of Johnson's Medico-Chirurgical Review. Among foreign periodicals none can be more useful to the American practitioner both for enlightened views and extensive information.

T. ROMEYN BECK, (Albany.)

Sir,—I consider Dr. Johnson's "Medico-Chirurgical Review" one of the best periodicals. Journals of Great Britain, and believe that your republication of it will prove a valuable accession to the Medical Literature of this country. You have my best wishes for its success.

DAVID HOSACK, M. D., Pres. of Med. Fac. of Rutgers Coll.

We fully concur in the opinion expressed by Dr. Hosack, of the value of Johnson's Journal. SAML. L. MITCHELL, M. D., Vice-Pres., WILLIAM J. MACNEVIN, M. D., Prof. of Ther. and Mat. Med.; VALENTINE MOTT, M. D., Prof. of Surg.; JOHN W. FRANCIS, M. D., Prof. of Obstet., &c.; JOHN D. GODMAN, M. D., Prof. of Anat. and Physiol.; JOHN GRISCOM, LL. D., Prof. of Chem.; P. S. TOWNSEND, M. D., Regist.

N. CHAPMAN, M. D., Prof. of Prac. Med. of University of Pennsylvania; W. E. HORNER, M. D., Prof. of Anat.; GEORGE B. WOOD, M. D., Prof. of Mat. Med.; WM. GIBSON, M. D., Prof. of Surg.; H. L. HODGE, M. D., Prof. of Obstets.; S. JACKSON, M. D., Prof. of Insts. of Med.

R. DUNGLISON, M. D., Prof. of Insts. of Med. &c., of Jefferson Medl. College of Phila.; R. M. HUSTON, M. D., Prof. of Mat. Med.; JOS. PANCOAST, M. D., Prof. of Surgl. Anat.; J. K. MITCHELL, M. D., Prof. of Prac. of Med.; THOMAS J. MUTTER, M. D., Prof. of Prac. and Surg.; CHAS. D. MEIGS, M. D., Prof. of Obstets.

S. G. MORTON, M. D., Prof. of Anat. and Phys., of Pennsylvania College; GEORGE M'CLELLAN, M. D., Prof. of Surg.; WM. RUSH, M. D., Prof. of Prac. Med.; R. M. BIRD, M. D., Prof. of Mat. Med.; SAMUEL M'CLELLAN, M. D., Prof. of Obstets.

Sir,—I am much gratified to learn, that you propose to continue the republication of Johnson's Medico-Chirurgical Review. This work I have taken from its commencement in 1820, and from a comparison of it with other periodical publications, I have no hesitation in expressing my belief, that it is the best Medical Journal now published in Europe. It is not like many works of the kind, a mere temporary journal, a repository only of the most recent intelligence, but it conveys to the Profession an inestimable treasure, a full analytical view of Medical Literature, serving as a book of study for the present time, and a most valuable record for future reference. It cannot be too strongly recommended, and particularly to those who, from extensive professional avocations, or removal from good libraries, have no means of consulting the original works which daily issue from the press.

Truly and respectfully yours, THOMAS SEWELL, (Washington, D. C.)

My Dear Sir,—In compliance with your request of my opinion of the Medico-Chirurgical Review, I make no hesitation in saying, that in the style and arrangement of its matter, the number and variety of subjects discussed; the impartiality and strictness of its critiques, it is superior to any other British journal with which I am acquainted, and not inferior to any journal, European or American. The impartiality it has evinced towards American authors is an additional claim to the regard of the American Physician. I do therefore hope, most sincerely, that the Medico-Chirurgical Review will meet with a most liberal patronage, and that it will occupy a place in the library of every physician.

Very respectfully yours, W. WILLOUGHBY, (Newport, N. Y.)

The London Medico-Chirurgical Review, conducted by Dr. Johnson and others, is one of the best periodical works on Medical Science. Its republication in this country must be considered very useful to the medical profession.

JOHN C. WARREN, (Prof. of Anat. and Surg., Boston.)

Sir,—I have been a reader of the Medico-Chirurgical Review from its first publication, and have found it at all times to contain a general view of the current medical literature of the day. This makes its pages very valuable, and enables the medical practitioner to judge what works he need to possess himself of. I hope that its republication in this country will be sufficiently encouraged. I am, Sir, your obedient servant,

JAMES JACKSON, (Boston.)

* * Subscribers in ordering this work, are requested to write their names legibly, and at full length, adding the names of the town, county, &c. of their residence.

One dollar allowed to any Subscriber for procuring a new Subscriber and remitting payment for the same.





